

Lecture 5 Gram-Schmidt Orthogonalization

MIT 18.335J / 6.337J
Introduction to Numerical Methods

Per-Olof Persson
September 21, 2006

1

Gram-Schmidt Projections

- The orthogonal vectors produced by Gram-Schmidt can be written in terms of projectors

$$q_1 = \frac{P_1 a_1}{\|P_1 a_1\|}, \quad q_2 = \frac{P_2 a_2}{\|P_2 a_2\|}, \quad \dots, \quad q_n = \frac{P_n a_n}{\|P_n a_n\|}$$

where

$$P_j = I - \hat{Q}_{j-1} \hat{Q}_{j-1}^* \text{ with } \hat{Q}_{j-1} = \begin{bmatrix} | & | & \dots & | \\ q_1 & q_2 & \dots & q_{j-1} \\ | & | & \dots & | \end{bmatrix}$$

- P_j projects orthogonally onto the space orthogonal to $\langle q_1, \dots, q_{j-1} \rangle$, and $\text{rank}(P_j) = m - (j - 1)$

2

The Modified Gram-Schmidt Algorithm

- The projection P_j can equivalently be written as

$$P_j = P_{\perp q_{j-1}} \cdots P_{\perp q_2} P_{\perp q_1}$$

where (last lecture)

$$P_{\perp q} = I - qq^*$$

- $P_{\perp q}$ projects orthogonally onto the space orthogonal to q , and $\text{rank}(P_{\perp q}) = m - 1$
- The *Classical Gram-Schmidt* algorithm computes an orthogonal vector by

$$v_j = P_j a_j$$

while the *Modified Gram-Schmidt* algorithm uses

$$v_j = P_{\perp q_{j-1}} \cdots P_{\perp q_2} P_{\perp q_1} a_j$$

3

Classical vs. Modified Gram-Schmidt

- Small modification of classical G-S gives modified G-S (but see next slide)
- Modified G-S is numerically stable (less sensitive to rounding errors)

Classical/Modified Gram-Schmidt

for $j = 1$ to n

$$v_j = a_j$$

for $i = 1$ to $j - 1$

$$\begin{cases} r_{ij} = q_i^* a_j & \text{(CGS)} \\ r_{ij} = q_i^* v_j & \text{(MGS)} \end{cases}$$

$$v_j = v_j - r_{ij} q_i$$

$$r_{jj} = \|v_j\|_2$$

$$q_j = v_j / r_{jj}$$

4

Implementation of Modified Gram-Schmidt

- In modified G-S, $P_{\perp q_i}$ can be applied to all v_j as soon as q_i is known
- Makes the inner loop iterations independent (like in classical G-S)

Classical Gram-Schmidt

for $j = 1$ to n

$$v_j = a_j$$

for $i = 1$ to $j - 1$

$$r_{ij} = q_i^* a_j$$

$$v_j = v_j - r_{ij} q_i$$

$$r_{jj} = \|v_j\|_2$$

$$q_j = v_j / r_{jj}$$

Modified Gram-Schmidt

for $i = 1$ to n

$$v_i = a_i$$

for $i = 1$ to n

$$r_{ii} = \|v_i\|$$

$$q_i = v_i / r_{ii}$$

for $j = i + 1$ to n

$$r_{ij} = q_i^* v_j$$

$$v_j = v_j - r_{ij} q_i$$

5

Example: Classical vs. Modified Gram-Schmidt

- Compare classical and modified G-S for the vectors

$$a_1 = (1, \epsilon, 0, 0)^T, \quad a_2 = (1, 0, \epsilon, 0)^T, \quad a_3 = (1, 0, 0, \epsilon)^T$$

making the approximation $1 + \epsilon^2 \approx 1$

- Classical:

$$v_1 \leftarrow (1, \epsilon, 0, 0)^T, \quad r_{11} = \sqrt{1 + \epsilon^2} \approx 1, \quad q_1 = v_1 / r_{11} = (1, \epsilon, 0, 0)^T$$

$$v_2 \leftarrow (1, 0, \epsilon, 0)^T, \quad r_{12} = q_1^T a_2 = 1, \quad v_2 \leftarrow v_2 - 1q_1 = (0, -\epsilon, \epsilon, 0)^T$$

$$r_{22} = \sqrt{2}\epsilon, \quad q_2 = v_2 / r_{22} = (0, -1, 1, 0)^T / \sqrt{2}$$

$$v_3 \leftarrow (1, 0, 0, \epsilon)^T, \quad r_{13} = q_1^T a_3 = 1, \quad v_3 \leftarrow v_3 - 1q_1 = (0, -\epsilon, 0, \epsilon)^T$$

$$r_{23} = q_2^T a_3 = 0, \quad v_3 \leftarrow v_3 - 0q_2 = (0, -\epsilon, 0, \epsilon)^T$$

$$r_{33} = \sqrt{2}\epsilon, \quad q_3 = v_3 / r_{33} = (0, -1, 0, 1)^T / \sqrt{2}$$

6

Example: Classical vs. Modified Gram-Schmidt

- Modified:

$$v_1 \leftarrow (1, \epsilon, 0, 0)^T, \quad r_{11} = \sqrt{1 + \epsilon^2}, \quad q_1 = v_1 / r_{11} = (1, \epsilon, 0, 0)^T$$

$$v_2 \leftarrow (1, 0, \epsilon, 0)^T, \quad r_{12} = q_1^T v_2 = 1, \quad v_2 \leftarrow v_2 - r_{12} q_1 = (0, -\epsilon, \epsilon, 0)^T$$

$$r_{22} = \sqrt{2}\epsilon, \quad q_2 = v_2 / r_{22} = (0, -1, 1, 0)^T / \sqrt{2}$$

$$v_3 \leftarrow (1, 0, 0, \epsilon)^T, \quad r_{13} = q_1^T v_3 = 1, \quad v_3 \leftarrow v_3 - r_{13} q_1 = (0, -\epsilon, 0, \epsilon)^T$$

$$r_{23} = q_2^T v_3 = \epsilon / \sqrt{2}, \quad v_3 \leftarrow v_3 - r_{23} q_2 = (0, -\epsilon/2, -\epsilon/2, \epsilon)^T$$

$$r_{33} = \sqrt{6}\epsilon/2, \quad q_3 = v_3 / r_{33} = (0, -1, -1, 2)^T / \sqrt{6}$$

- Check Orthogonality:

- Classical: $q_2^T q_3 = (0, -1, 1, 0)(0, -1, 0, 1)^T / 2 = 1/2$

- Modified: $q_2^T q_3 = (0, -1, 1, 0)(0, -1, -1, 2)^T / \sqrt{12} = 0$

7

Operation Count

- Count number of floating points operations – “flops” – in an algorithm
- Each +, −, *, /, or $\sqrt{\quad}$ counts as one flop
- No distinction between real and complex
- No consideration of memory accesses or other performance aspects

8

Operation Count - Modified G-S

- Example: Count all +, −, *, / in the Modified Gram-Schmidt algorithm (not just the leading term)

- (1) for $i = 1$ to n
- (2) $v_i = a_i$
- (3) for $i = 1$ to n
- (4) $r_{ii} = \|v_i\|$ m multiplications, $m - 1$ additions
- (5) $q_i = v_i / r_{ii}$ m divisions
- (6) for $j = i + 1$ to n
- (7) $r_{ij} = q_i^* v_j$ m multiplications, $m - 1$ additions
- (8) $v_j = v_j - r_{ij} q_i$ m multiplications, m subtractions

9

Operation Count - Modified G-S

- The total for each operation is

$$\begin{aligned} \#A &= \sum_{i=1}^n \left(m - 1 + \sum_{j=i+1}^n m - 1 \right) = n(m - 1) + \sum_{i=1}^n (m - 1)(n - i) = \\ &= n(m - 1) + \frac{n(n - 1)(m - 1)}{2} = \frac{1}{2}n(n + 1)(m - 1) \\ \#S &= \sum_{i=1}^n \sum_{j=i+1}^n m = \sum_{i=1}^n m(n - i) = \frac{1}{2}mn(n - 1) \\ \#M &= \sum_{i=1}^n \left(m + \sum_{j=i+1}^n 2m \right) = mn + \sum_{i=1}^n 2m(n - i) = \\ &= mn + \frac{2mn(n - 1)}{2} = mn^2 \\ \#D &= \sum_{i=1}^n m = mn \end{aligned}$$

10

Operation Count - Modified G-S

and the total flop count is

$$\begin{aligned} \frac{1}{2}n(n + 1)(m - 1) + \frac{1}{2}mn(n - 1) + mn^2 + mn = \\ 2mn^2 + mn - \frac{1}{2}n^2 - \frac{1}{2}n \sim 2mn^2 \end{aligned}$$

- The symbol \sim indicates asymptotic value as $m, n \rightarrow \infty$ (leading term)
- Easier to find just the leading term:
 - Most work done in lines (7) and (8), with $4m$ flops per iteration
 - Including the loops, the total becomes

$$\sum_{i=1}^n \sum_{j=i+1}^n 4m = 4m \sum_{i=1}^n (n - i) \sim 4m \sum_{i=1}^n i = 2mn^2$$

11

MIT OpenCourseWare
<https://ocw.mit.edu>

18.335J Introduction to Numerical Methods
Spring 2019

For information about citing these materials or our Terms of Use, visit: <https://ocw.mit.edu/terms>.