UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

International General Certificate of Secondary Education

MARK SCHEME for the June 2004 question papers

	0652 PHYSICAL SCIENCE							
0652/01	Paper 1 (Multiple Choice), maximum raw mark 40							
0652/02	Paper 2 (Core), maximum raw mark 80							
0652/03	Paper 3 (Extended), maximum raw mark 80							
0652/05	Paper 5 (Practical), maximum raw mark 30							
0652/06	Paper 6 (Alternative to Practical), maximum raw mark 60							

These mark schemes are published as an aid to teachers and students, to indicate the requirements of the examination. They show the basis on which Examiners were initially instructed to award marks. They do not indicate the details of the discussions that took place at an Examiners' meeting before marking began. Any substantial changes to the mark scheme that arose from these discussions will be recorded in the published *Report on the Examination*.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the *Report on the Examination*.

• CIE will not enter into discussion or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the June 2004 question papers for most IGCSE and GCE Advanced Level syllabuses.



	maximum	mir	minimum mark required for grade:				
	mark available	A	С	E	F		
Component 1	40	36	28	21	17		
Component 2	80	-	45	29	24		
Component 3	80	49	31	19	14		
Component 5	30	23	19	16	14		
Component 6	60	51	37	24	18		

Grade thresholds taken for Syllabus 0652 (Physical Science) in the June 2004 examination.

The threshold (minimum mark) for B is set halfway between those for Grades A and C. The threshold (minimum mark) for D is set halfway between those for Grades C and E. The threshold (minimum mark) for G is set as many marks below the F threshold as the E threshold is above it.

Grade A* does not exist at the level of an individual component.



INTERNATIONAL GCSE

MARK SCHEME

MAXIMUM MARK: 40

SYLLABUS/COMPONENT: 0652/01

PHYSICAL SCIENCE Paper 1 (Multiple Choice)



Page 1	Mark Scheme	Syllabus	Paper
	PHYSICAL SCIENCE – JUNE 2004	0652	1

Question Number	Key	Question Number	Key
1	С	21	D
2	D	22	Α
3	D	23	D
4	С	24	D
5	В	25	D
6	С	26	С
7	В	27	Α
8	Α	28	D
9	В	29	Α
10	Α	30	С
11	D	31	С
12	С	32	D
13	Α	33	С
14	Α	34	Α
15	D	35	С
16	С	36	Α
17	С	37	Α
18	D	38	D
19	Α	39	D
20	D	40	В

TOTAL 40

INTERNATIONAL GCSE

MARK SCHEME

MAXIMUM MARK: 60

SYLLABUS/COMPONENT: 0652/02

PHYSICAL SCIENCE Paper 2 (Core)



Pac	ge 1		Mark Scheme	Syllabus	Paper
		PHYSICA	L SCIENCE – JUNE 2004	0652	2
(a)		Points correctly plo	tted		2
()			d/incorrectly plotted)		_
		Good straight line o	· · <i>·</i>		1
					_
(b)			ures taken from graph		1
		Clear use of figures Correct answer = 0			1 1
			.75 611		1
(c)		930 +/-10 N			2
		(Accept 905 to 955	for 1 mark)		
				-	
				I	otal
(a)		Mark vertically:	8; 8; 2,6		1
(~)		· · · · · · · · · · · · · · · · · · ·	8; 10; 2,6		1
			(Repeated error penalise once	e only)	
/L-`		Dat areas distant	chaning pair of classes -		
(b)		And correct outer s	sharing pair of electrons		1
		(OR H-O-H with co			
		(,		
				Т	otal
		2			
(a)		3			1
(b)		12 + 3 + 16 + 1			
. ,		= 32			
		–			
(c)			olecules stronger in methanol rrect statements about hydrog	on honding	, nin
		methanol, not in ca			, 111
				Т	otal
(a)		Mention of surface			4
		Much greater for a	powaer		1 ·
(b)	(i)	Dilute the acid (acc	ept add water)		
. /		·			
	(ii)	Lower the tempera	ture		
				т	otal
				I	
(a)		(Current in the coil)	magnetises the core		
		Attracting the bolt	-		
/L-`		It is measured a			
(b)		It is magnetic And loses its magn	atism agaily		•
		And ioses its mayn	CUSITI COSILY		
(c)		No current can flow	1		
•		So bolt remains in s	situ		
				-	- 4- I
				T	otal

Pag	e 2	Mark Scheme	Syllabus	Paper
		PHYSICAL SCIENCE – JUNE 2004	0652	2
(a)		Potential energy is released As particles move together (Do not accept answers which refer to loss of KE/s particles)	lowing dow	1 1 n of
(b)	(i)	330°C +/- 5°C		1
	(ii)	P solidifies at one temperature Q solidifies over a range of temperatures		1 1
			т	otal
(a)		Potassium is more reactive than magnesium (OR is higher up the activity series)		1
(b)	(i)	Energy is released		1
	(ii)	Litmus paper/universal indicator Turns blue/green		1 + ⁻
	(iii)	Lighted splint Causes small explosion/pop		1 + ⁻
			т	otal
(a)		Elastic/strain Kinetic/movement Heat/thermal/internal Work		1 1 1
(b)		2.5 × 3 7.5 Ncm (-1 if no/incorrect unit)		1 2
(c)		48/16 3 m/s (-1 if no/incorrect unit)		1 2
			т	otal
(a)		Combines with haemoglobin (Accept blood) Preventing oxygen being absorbed		1 1
(b)		Combines with rain water To form acid (rain)		1 1
			т	otal

	e 3	Mark Scheme	Syllabı		per
		PHYSICAL SCIENCE – JUNE 2004	0652		2
(a)		H H Ethanol	с — он		
		H-C-C-OH	1		1
		H H fully com	ect		1
		H - c - c < 0 Ethanoic acid:	с он		1
		H OH fully con	rect		1
(b)		Any TWO from: Fuel, solvent, in drinks		- ()	1+
				Total	
(a)		Ammeter			1
		Voltmeter Variable resistor			1 1
(h)		By changing the registerion			1
(b)		By changing the resistance The current in the circuit can be changed			1 1
(c)		Straight line through the origin OR curve increasing current In both quadrants	so that R increase	es with	1
		in both quadrants			
				Total	
		Acidic			1
		Non-metal Right			1 1
		Ngin			•
				Total	
(a)		Filament gets very hot Must not be allowed to oxidise/burn Argon provides inert atmosphere	ANY TWO		1+
(b)		High density High melting point Transition part of the Periodic Table	ANY TWO		1 +
				Total	
	(i)	Negative			4
(a)	(i)	Attracted to positive collector			1 1
(a)		Floatron			1
(a)	(ii)	Electron			
	(ii)				1
(a) (b)	(ii)	Deflect rays* Horizontally			1
	(ii)	Deflect rays* Horizontally Deflect rays*			1 1
	(ii)	Deflect rays* Horizontally	e)		1 1 1 1
(b)		Deflect rays* Horizontally Deflect rays* Vertically (* can be scored in either part but only onc			1 1 1
	(ii) (i)	Deflect rays* Horizontally Deflect rays* Vertically	e same		1 1 1 1
(b)		Deflect rays* Horizontally Deflect rays* Vertically (* can be scored in either part but only onc Amplitude smaller but frequency (about) th	e same		1 1 1

INTERNATIONAL GCSE

MARK SCHEME

MAXIMUM MARK: 80

SYLLABUS/COMPONENT: 0652/03

PHYSICAL SCIENCE Paper 3 (Extended)



Page 1	Mark Scheme Syllabus Pap	er
	PHYSICAL SCIENCE – JUNE 2004 0652 3	
(a)	(average) mass of one atom (of element) (of normal isotopic mixture) compared to 1/12 mass of one atom of carbon-twelve	1
	OR on a scale on which one atom of carbon-twelve has a mass of 12 exactly	1
(b) (i)	n = m/ <i>M</i> _r OR 5.0 / 30 Accept 5 / 30 .	1
	number of moles = 0.167 Accept 1/6, 0.17, 0.16 but not 0.2.	1
(ii)	(2.0 / 24) number of moles = 0.083	
	Accept 1/12. Accept 0.08 only if 2/24 shown.	
(iii)	(answer from (i) ÷ answer from (ii)) number of moles = 2 Accept answer from errors carried forward.	
(iv)	$2M + O_2 \rightarrow 2MO$	
	Answer from (iii) must be used in front of M. correct formulae of elements M and O ₂ balanced using answer from (iii)	1 1
	Total	
(a)	put water into can up to spout place measuring cylinder under spout <u>and</u> lower object into can (until immersed)	1 1
	volume of water displaced into cylinder equals volume of object	1
(b) (i)	g/cm ³ OR kg/m ³ etc	1
	Symbols must be correct, as listed in the syllabus	
(ii)	density = mass / volume OR 15.4 / 0.8 density = 19.25 (g/cm ³) numerical answer only	
	Accept 19.3 or 19.2 (Also accept 19 because volume given only to 1 sig. fig.)	
(iii)	gold	
	Accept error forward from (ii)	
(iv)	ideas of uncertainty of experimental method uncertainty of experimental readings may not be pure metal	1 1 1
	Accept explanation in terms of significant figures for one mark.	

Accept explanation in terms of significant figures for one mark.

Page 2	Mark Scheme St	yllabus	Paper		
		0652	3		
(c)	85g → 0.085kg <i>OR</i> equivalent W = mg <i>OR</i> g = W/m			1 1	
	Accept with values inserted whether mass is in gram	ns or kild	ograms		
	g = 1.65 N/kg complete answer			1	[
	Accept unit m/s². Symbols in unit must be correct, as list Accept 1.6 but not 1.7 because 0.14 / 0.085 = 1.647	ted in sy	llabus.		
			Total		[′
(a)	increase to silicon then decrease			1	[
	Ignore P & S anomaly. Must mention silicon.				
(b)	strong (forces of attractions between atoms) due to covalent bonding <i>OR</i> giant (tetrahedral) struct	ure		1 1	ĺ
(c)	Any symbols used should be correct, as listed in syllabu	s			
	(i) sodium (ii) phosphorus			1 1	
	(iii) magnesium			1	
	(iv) argon			1	
(d)	<i>ideas of…</i> sodium ions have +1 charge <u>and</u> magnesium ions have ∴ forces of (attraction) in metallic bonding weaker in soc magnesium			1 1	
	Comparison must be clear.				
			Total		
(a)	wire connected across voltmeter			1	
	Accept, for this circuit, wire connected across ba Be tolerant with symbol or drawing to represent t				
(b)	R = V/I OR $4.3 / 2.1$ resistance = 2.05 Ω numerical value (1) unit (1)			1 2	
	Accept 2.0, 2.04 but not 2.1. The mark for the unit Ω is a separate mark.				
(c)	twice the answer from (b)			1	
(d)	Ignore unit. state resistance of <u>shorter</u> wire likely to be more than	expect	ed	1	
. /	explain shorter wire (less resistance) more current			1	
	∴ hotter than longer wire Comparison must be clear.			1	
(e)	large current			1	
	could overheat ammeter			1	

[Page 3	Mark Scheme Syllabus Paper	,	
		PHYSICAL SCIENCE – JUNE 2004 0652 3		
	(f)	oscilloscope OR c.r.o. OR multimeter	1	[1]
		Total		[11
5	(a) (i)	calcium 2,8,8,2 fluorine 2,7	1 1	[2]
	(ii)	transfer of electrons from calcium atoms to fluorine atoms forming positive ions (Ca ²⁺) and negative ions (F-) that attract	1 1	[2]
	(iii)	CaF ₂	1	[1]
		Do not accept F1 for fluorine.		
	(b)	solid calcium fluorideions ions are held in lattice OR cannot move aboutmolten calcium fluorideions ions are free to move aboutliquid fluorinemolecules are not charged	1 1 1	[3]
		Total		[8]
6	(a)	n = 8		[1]
	(b)	speed = distance/time OR time = distance/speed OR time = 80/340 ∴ time = 0.235 s complete answer (1) Accept 0.24 s or 0.23 s but not 0.2 s	1 1	[2]
	(c) (i)	<i>ideas of…</i> start: fast speed of light means negligible delay in seeing smoke stop: slow speed of sound gives enough time for observer to respond	1 1	[2]
	(ii)	decreases possibility of echoes which would confuse observer	1 1	[2]
	(d)	3.5 kHz \rightarrow 3500 Hz v = f λ OR λ = v/f (accept c = f λ or λ = c/f). Accept with values inserted whether frequency is in kHz or Hz.	1 1	
		wavelength = 0.097 m complete answer * (1)		
		Do not accept 0.1 m.	1	[3]
		* Only the first incorrect or missing unit is penalised Total		[10
7	(a)	yeast temperature less than 40 °C	1 1	[2]
		Do not accept 'warm' on its own.		
	(b) (i)	fractional distillation both words	1	[1]

Page 4	Mark Scheme Syllabus Pap PHYSICAL SCIENCE – JUNE 2004 0652 3 Iabelled sketch of laboratory apparatus to show fractionating column fractionating column	
(ii)		
	thermometer condenser workable arrangement *	1 1 1 1
	* showing flask of solution being heated, vapour rising up fractionating column, thermometer in the top of this column with its bulb opposite tube leading down through water-cooled condenser into collecting vessel; the condenser should have water entering and leaving the outer tube correctly.	
	Tota	I
(a)	thermometer changes do not accept 'expands' equal range	1 1 1 1
	sensitive do not accept 'accurate'	1
(b)	<i>examples…</i> liquid-in-glass thermometer volume of liquid depends on temperature	
	accept named liquid, mercury or alcohol.	
	OR thermocouple \checkmark e.m.f depends on temperature \checkmark	
	Tota	I
(a)	to remove impurities (from the ore)	1
	Do not accept 'to form slag' unless 'impurities' are mentioned.	
(b)	Symbols and subscripts should be written correctly.	
	(i) $CaCO_3 \rightarrow CaO + CO_2$	
	formulae (1) (then) balanced (1)	2
	(ii) $Fe_2O_3 + 3CO \rightarrow 2Fe + 3CO_2$ formulae (1) (then) balanced (1)	2
	Accept $2Fe_2O_3 + 3C \rightarrow 4Fe + 3CO_2$	
(c)	<i>ideas of…</i> zinc is more reactive than iron ∴ when zinc-coating is	

Page 5	Mark Scheme	Syllabus	Paper
	PHYSICAL SCIENCE – JUNE 2004	0652	3

- Total [7]
- TOTAL FOR PAPER [80]

INTERNATIONAL GCSE

MARK SCHEME

MAXIMUM MARK: 30

SYLLABUS/COMPONENT: 0652/05

PHYSICAL SCIENCE Practical



<u>⊢</u> •	age 1	Mark Scheme PHYSICAL SCIENCE – JUNE 2004	Syllabus	Pape
		PHYSICAL SCIENCE – JUNE 2004	0652	5
(a)	(i)	Value for h within 0.4 mm of supervisor		1
	(ii)	Brief description of how volume was found Volume within 10 cm ³ of supervisor sensible volume		2
		Table:		
		Six pairs of values Good spread to include a value equal to 150 cm ³ Values in mm and decreasing with volume of water (Penalise 1 mark when all intervals are exactly the same)	3
(b)		Graph:		
		Axes correctly labelled Sensible scales for plotted points Plotting correct for 4 values		
		Best straight line drawn		4
		Volume correctly read needs evidence of extrapolation Within 10% of recorded volume		2
(c)		Measure water level in cylinder Put in the block and record new level Volume of water displaced calculated is equal to the volu block	ime of	3
			Total	[15
(a)		Gas/vapour burns Limewater milky Brown or charring/smoke/smell		3
(b)		Goes out NOT 'nothing' Limewater milky		2
(c)	(i)	Decolourised		1
(-)				•
(-)	(ii)	UI goes red pH about 1-4 Acid present		3
(d)	(ii)	pH about 1-4 Acid present Blue/green pH about 8-10		3
(d)	(ii)	pH about 1-4 Acid present Blue/green pH about 8-10 No mark for conclusion		3
	(ii)	pH about 1-4 Acid present Blue/green pH about 8-10		3
(d)	(ii)	pH about 1-4 Acid present Blue/green pH about 8-10 No mark for conclusion		3

INTERNATIONAL GCSE

MARK SCHEME

MAXIMUM MARK: 60

SYLLABUS/COMPONENT: 0652/06

PHYSICAL SCIENCE Alternative to Practical



F	Page 1		Mark Scheme		Syllabus	Paper	
			PHYSICAL SCIENCE – JUNE 2	004	0652	6	
1	(a)		2.6 cm, 5.8 cm correctly entered in Fig. 1.2 (no tolerance)			[2]	
	(b)		displacement increases as load increases OWTTE			[1]	
	(c)		repeat experiment (and average)/us	e a ruler mark	ed in millim	etres	[1]
	(d)	(i)	thicker beam gives smaller displacement OWTTE			[1]	
		(ii)	shorter beam gives smaller displacement OWTTE				[1]
	(e)		hang object on beam [1] read displacement [1] compare result with data from the ex by plotting a graph of the data [1]	periment [1]			[4]
						Tota	[10]
2	(a)		1.8V [1], 150mA 2.4V [1], 250mA (1 mark for bo +/- 0.1V, +/- 10mA	th current rea	dings)		[3]
	(b)		2 points correctly plotted [2] line drawn (can be straight or curved	i) [1]			[3]
	(c)	(i)	the bulb becomes brighter as resista	ince decrease	es		[1]
		(ii)	the filament of the bulb melted OWT	TE			[1]
	(d)		no, since it is not a straight line/V an OR	d I are not pro	oportional		[1]
			yes, graph is a straight line/(they are	proportional))		
						Tota	al [9]
3	(a)	(i)	53.4g, 60.0g (must say 60.0	0), no tolerano	ce [2]		[3]
		(ii)	6.6g (ecf) [1]				
	(b)		blue litmus (U.I) paper turns red in th	ne gas (reject	add indicato	or)	[1]
	(c)	(i)	56.8g (no tolerance)				[1]
		(ii)	3.2g (ecf) (both correct f	or 1 mark)			
	(d)		evaporate to remove some water [1] leave the solution to cool [1] OR evaporate solution [1] over a boiling water bath [1]				[2]

F	Page 2		Mark Scheme	Syllabus	Paper		
			PHYSICAL SCIENCE – JUNE 2004	0652	6		
	(e)	(i)	62.9g, (no tolerance) [1]			[2]	
		(ii)	9.5g (ecf) [1]				
	(f)		some copper nitrate left in the solution during crystallisation/water of crystallisation was lost/copper nitrate decomposed/other suitable answer based on experimental details				
					Total	[10]	
4	(a)		gas C: 8s gas D: 3s gas E: 12s. (no tolerance)			[3]	
	(b)		gas C because it took the least time to fall OWTT	E		[1]	
	(c)		heavier (denser) gases fall, lighter (less dense) gases rise [1] gases less dense (lighter) than air rise [1] gases more dense (heavier) than air fall [1]				
	(d)		to keep the experiment fair/so that the results are	accurate		[1]	
	(e)	(i)	gas A rose more quickly/it has the least density			[1]	
		(ii)	test with a lighted spill/burn in air [1] gas explodes (pop!) [1]			[2]	
					Total	[10]	
5	(a)		box 1 colourless (clear) to cloudy/milky [1] carbon dioxide/carbonate [1] box 2(a) carbon dioxide (suspected)/gas will not s combustion/no oxygen/may be nitrogen [1] box 2(b) carbon dioxide confirmed [1] box 3 turned from green [1] to red [1] box 4 turned yellow/orange (reject orange) [1]	upport		[7]	
	(b)		reaction vessel with delivery tube [1] gas collected over water or in a syringe [1] means of measuring gas volume/graduations sho	wn [1]		[3]	
					Total	[10]	
6	(a)	(i)	use a pipette/dropper/burette			[1]	
		(ii)	103 (no tolerance) [1] 147 (ecf) [1]			[2]	
	(b)		28mm, 14mm (+/- 1mm)			[2]	

Page 3		Mark Scheme	Syllabus	Paper	
		PHYSICAL SCIENCE – JUNE 2004	0652	6	
(c)	(i)	axes labelled and scale correctly shown [1] all points from Fig. 6.3 plotted correctly [1] straight line drawn extended to cut horizontal axis	[1]		[3]
	(ii)	from candidates' own graph (approx 147)			[1]
	(iii)	it will sink OWTTE			[1]
(d)		yes/comparison of (a) and (c)(ii) shows that numerically similar to (or greater than) its volume OR no/cup sank before its mass (g) exceeded to (depends on candidate's graph) (mark for explanation)			[1]

Total [11]