

Data table for Figure 9. Current asthma among children under age 18 years, by race and Hispanic origin: United States, 2007–2017

Excel and PowerPoint: https://www.cdc.gov/nchs/hus/contents2018.htm#Figure_009

Year	Race and Hispanic origin							
	Total ¹		Hispanic		Not Hispanic or Latino			
	Percent	SE	Percent	SE	White only		Black only	
					Percent	SE	Percent	SE
2007	9.1	0.4	9.3	0.7	7.3	0.5	15.4	1.1
2008	9.4	0.4	6.7	0.6	8.8	0.5	15.7	1.2
2009	9.6	0.4	7.7	0.6	8.5	0.5	17.0	1.3
2010	9.4	0.3	8.1	0.6	8.2	0.5	15.9	1.0
2011	9.5	0.3	9.6	0.6	7.8	0.4	16.3	1.1
2012	9.3	0.3	8.8	0.6	7.9	0.4	16.0	1.0
2013	8.3	0.3	7.4	0.5	7.5	0.5	13.4	0.9
2014	8.6	0.3	8.5	0.6	7.6	0.5	13.4	1.0
2015	8.4	0.3	8.0	0.5	7.4	0.5	13.4	1.0
2016	8.3	0.3	6.7	0.6	7.1	0.4	15.7	1.3
2017	8.4	0.4	7.7	0.7	7.7	0.5	12.6	1.3

¹Includes all other races not shown separately.

NOTES: SE is standard error. Data are for the civilian noninstitutionalized population. Based on parent or knowledgeable adult responding yes to both questions, “Has a doctor or other health professional ever told you that your child had asthma?” and “Does your child still have asthma?” Adults of Hispanic origin may be of any race. Race-specific estimates are tabulated according to the 1997 Revisions to the *Standards for the Classification of Federal Data on Race and Ethnicity*. The single-race categories plus multiple-race category shown in the table conform to the 1997 Standards. Starting with 2003 data, race responses of other race and unspecified multiple race were treated as missing, and then race was imputed if these were the only race responses. Almost all persons with a race response of other race were of Hispanic origin. See [Appendix II](#), Hispanic origin; Race.

SOURCE: NCHS, National Health Interview Survey. See [Appendix I](#), National Health Interview Survey (NHIS).