Author Year of systematic review	Time period and databases searched in systematic review	Eligibility criteria in systematic review	Study designs of eligible studies	Countries included in eligible studies	Sample size in eligible studies	Population in eligible studies	Interventions in eligible studies	Main results of eligible studies
Abbass 2009 <sup>102</sup>	CCDANCTR-Studies, CCDANCTR-References; CENTRAL, MEDLINE, CIHAHL, EMBASE, PsycINFO, DARE, Biological Abstracts: database inception-April 2005	All RCTs in which short-term psychodynamic psychotherapies was compared with wait-list controls, minimal treatment controls which had been designated as psychological "placebo treatments," and treatments as usual; adult outpatients with common mental disorders (excluding psychotic disorders)	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs
Binks 2011 <sup>104</sup>	Medline: 1966-January 2003; PsycINFO: 1872-December 2002; CCRCT: to October 2002; EMBASE: 1980-January 2003; and 21 additional specialist databases	Clinical RCTs with or without blinding involving psychological treatments (behavioral, cognitive-behavioral, psychodynamic, and psychoanalytic) for adults with a diagnosis of borderline personality disorder	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs
Dieterich 2010 <sup>106</sup>	Cochrane Schizophrenia Group Trials Registry: database inception-February 2009	Randomized clinical trials focused on people with severe mental illness ages 18-65 years; community care setting; intensive case management (ICM) compared to non-ICM or standard care	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs
Dubicka 2010 <sup>87</sup>	PsycINFO, MEDLINE, Cochrane databases: January 1980-March 2009	RCTs predominantly including adolescents aged 11-18 years with a DSM-IV defined episode of depression where CBT was combined with a newer generation antidepressant and compared with antidepressant treatment without CBT	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs
Irving 2010 <sup>107</sup>	Cochrane Schizophrenia Group's Register of Trials: 1998-January 2006; additional searches in past versions of this review	RCTs and quasi-RCTs; patients with schizophrenia or other serious mental illness presenting to or referred to a social/psychiatric/ nursing service because they were experiencing a psychosocial crisis, however defined (excluded people in crisis with drug-induced psychosis or in a depressive crisis)	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs
Kavanagh 2009 <sup>108</sup>	MEDLINE, CINAHL, EMBASE, The Cochrane Library, PSYCINFO, ERIC, SOCIAL SCIENCE CITATION INDEX, ASSIA, Trials Register of Public Health Interventions (TROPHI), Database of Public Health Effectiveness Reviews (DOPHER), C2 SPECTR, PSITRI: Time period not reported	RCTs published from 1996 onwards in the English language that at least measured depression, anxiety, or suicidality following an intervention based on cognitive behavioral techniques delivered within secondary schools to young people aged 11-19	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs
Muralidharan 2009 <sup>111</sup>	CINAHL, CENTRAL, Schizophrenia Groups Register, EMBASE, MEDLINE, PsycINFO: database inception-January 2006	Relevant RCTs and quasi-RCTs; people with the diagnosis of serious mental illness (including "serious/chronic mental illness" or "psychotic illness"), however diagnosed		No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs

Author Year of systematic review	Time period and databases searched in systematic review	Eligibility criteria in systematic review	Study designs of eligible studies	Countries included in eligible studies	Sample size in eligible studies	Population in eligible studies	Interventions in eligible studies	Main results of eligible studies
National Collaborating Centre for Mental Health 2005 <sup>100</sup>	MEDLINE, EMBASE, PsycINFO, Cochrane Library: Inception to September 2004	RCTs of depressed participants aged 5-18 treated with CBT, CBT+separate parenting sessions, interpersonal psychotherapy, psychoanalytic/psychodynamic child psychotherapy, self-modeling, relaxation, social skills training, family therapy, guided self-help, or control enhancement training, and that reported remission, symptom levels, functional status or discontinuation from treatment for any reason outcomes.	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs
Newton 2010 <sup>112</sup>	MEDLINE, EMBASE, CCRCT, CDSR, Health Technology Assessment Database, DARE, Academic Search Elite, PsycINFO: 1985-October 2009; and 4 additional specialist databases	Experimental or quasi-experimental designs; mental health-based, suicide prevention-focused intervention initiated in the ED or immediately after ED discharge through direct referral/enrollment; children and adolescents (≤18 years), or their parents or ED personnel; ≥1 clinically relevant primary outcome	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs
Pharaoh 2010 <sup>113</sup>	Cochrane Schizophrenia Group Trials Register: to September 2008	Relevant RCTs or quasi-RCTs; studies where most (>75%) families included at least one member with a diagnosis of schizophrenia and/or schizoaffective disorder	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs
Shek 2010 <sup>114</sup>	Cochrane Schizophrenia Group Trials Register: to May 2009	Relevant RCTs; people aged 18-65 years and suffering from illness such as schizophrenia, schizophrenia-like disorders, and bipolar disorder (excluding acutely ill patients)	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs
Shekelle 2009 <sup>14</sup> & Bagley 2010 <sup>103</sup>	MEDLINE, Cochrane Library, PsycINFO: June 2005-May 2008	English language; suicide or suicide attempt outcomes; no mental health interventions such as psychotherapy or pharmacotherapy unless they included Veterans	One RCT: Unutzer 2006	US	1,801	Non-Veteran/ military; no other data reported	Collaborative care model including case management in a primary care setting	No suicides in either treatment or control group
State of Victoria Department of Health 2010 <sup>115</sup>	MEDLINE, EMBASE, AMED, PsycINFO: January 1997-February 2009	English language; human; suicide related outcome; sample size ≥6; no duplication; emergency department or other acute care setting	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs
Williams 2009 <sup>98</sup> & Williams 2009 <sup>99</sup>	DARE, CDSR, MEDLINE, PsycINFO: 1998-May 2006	Patients aged 7-18 years with MDD or depression NOS; primary care setting, school-based clinics; English language only; excluded poor quality studies	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs	No eligible RCTs