

**Table SU04.7 Types of Illicit Drug Use in the Past Month among Individuals Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Comparison of Percentages and Standard Errors from Restricted-Use File and Public Use File Estimates, 2004 NSDUH**

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH																								
	Cigarettes and Alcohol						Cigarettes But Not Alcohol						Alcohol But Not Cigarettes						Neither Alcohol Nor Cigarettes						
	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	
<b>ILLICIT DRUGS<sup>1</sup></b>	25.0	25.0	1.00	0.63	0.59	0.95	9.6	9.9	1.03	0.57	0.59	1.03	6.6	6.6	1.00	0.27	0.31	1.17	1.7	1.8	1.01	0.10	0.09	0.89	
Marijuana and Hashish	20.8	21.0	1.01	0.56	0.51	0.93	6.9	7.0	1.01	0.44	0.53	1.22	5.0	5.0	0.99	0.23	0.29	1.27	0.8	0.8	1.01	0.06	0.08	1.19	
Cocaine	3.6	3.5	0.95	0.24	0.28	1.16	0.7	1.08	0.14	0.15	1.03	0.5	0.5	1.08	0.07	0.09	1.24	0.0	0.0	1.06	0.02	0.01	0.84		
Crack	0.9	0.9	1.01	0.14	0.14	0.98	0.3	0.3	1.03	0.10	0.11	1.01	0.0	0.0	0.75	0.02	0.01	0.89	0.0	0.0	0.98	0.01	0.01	0.98	
Heroin	0.2	0.2	0.94	0.06	0.06	1.05	0.3	0.3	1.12	0.11	0.12	1.13	0.0	0.0	0.80	0.01	0.01	0.87	*	0.0	*	*	*	0.00	*
Hallucinogens	1.5	1.4	0.96	0.11	0.12	1.11	0.3	0.4	1.05	0.09	0.08	0.96	0.3	0.3	1.10	0.04	0.05	1.19	0.1	0.1	1.08	0.01	0.02	1.53	
LSD	0.2	0.2	0.96	0.04	0.05	1.16	0.1	0.1	0.82	0.06	0.05	0.80	0.0	0.0	0.89	0.01	0.01	0.90	0.0	0.0	1.33	0.00	0.00	1.49	
PCP	0.1	0.1	0.97	0.03	0.03	1.01	*	0.0	*	*	0.01	*	*	0.0	*	*	0.00	*	0.0	0.0	1.19	0.01	0.02	1.18	
Ecstasy	0.7	0.7	1.00	0.08	0.09	1.17	0.1	0.2	1.23	0.05	0.06	1.17	0.2	0.2	1.09	0.04	0.05	1.30	0.0	0.0	1.15	0.00	0.01	1.37	
Inhalants	0.6	0.6	0.98	0.06	0.07	1.09	0.1	0.1	1.01	0.03	0.04	1.18	0.2	0.2	1.00	0.04	0.04	1.09	0.2	0.2	0.97	0.05	0.04	0.82	
Nonmedical Use of Psychotherapeutics <sup>2</sup>	7.5	7.4	0.98	0.31	0.32	1.03	3.3	3.4	1.03	0.38	0.43	1.13	1.9	1.9	0.99	0.15	0.18	1.23	0.8	0.8	1.03	0.06	0.06	0.98	
Pain Relievers	5.6	5.5	0.98	0.25	0.29	1.15	2.3	2.4	1.04	0.33	0.36	1.10	1.3	1.3	0.99	0.11	0.14	1.23	0.6	0.7	1.03	0.06	0.06	1.14	
OxyContin®	0.6	0.6	0.96	0.07	0.07	0.99	0.2	0.1	0.85	0.06	0.06	0.96	0.0	0.0	0.49	0.02	0.01	0.47	0.0	0.0	1.08	0.01	0.01	0.79	
Tranquilizers	2.1	2.2	1.02	0.17	0.20	1.21	1.0	1.1	1.05	0.21	0.26	1.21	0.6	0.5	0.96	0.07	0.09	1.14	0.1	0.1	0.94	0.03	0.03	1.09	
Stimulants	1.7	1.7	0.96	0.14	0.15	1.05	0.6	0.7	1.13	0.15	0.18	1.20	0.4	0.4	1.05	0.07	0.08	1.14	0.1	0.1	1.11	0.02	0.03	1.39	
Methamphetamine	0.9	0.9	0.97	0.12	0.12	0.99	0.4	0.4	1.13	0.11	0.12	1.07	0.1	0.1	1.10	0.04	0.05	1.25	0.0	0.0	1.10	0.02	0.02	1.31	
Sedatives	0.3	0.2	0.75	0.09	0.09	1.06	0.2	0.1	0.78	0.10	0.09	0.86	0.1	0.1	1.13	0.03	0.03	1.18	0.0	0.0	1.07	0.02	0.02	1.14	
<b>ILLICIT DRUGS OTHER THAN MARIJUANA<sup>1</sup></b>	10.8	10.5	0.98	0.40	0.41	1.02	4.3	4.5	1.04	0.43	0.51	1.17	2.5	2.5	1.01	0.16	0.20	1.20	1.1	1.1	1.03	0.08	0.09	1.03	

\* Low precision; no estimate reported.

LSD = lysergic acid diethylamide; PCP = phencyclidine; PUF = public use file, which refers to estimates produced from a NSDUH public use file (a PUF is a NSDUH restricted-use file that has undergone statistical disclosure limitation treatment); RUF = restricted-use file, which refers to estimates produced from a NSDUH RUF and published in NSDUH's detailed tables available at <http://www.samhsa.gov/data/>; SE = standard error that is associated with the estimate.

NOTE: Ratio of estimates = PUF estimate ÷ RUF estimate; Ratio of SEs = PUF SE ÷ RUF SE.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>2</sup> Nonmedical use of prescription-type pain relievers, tranquilizers, stimulants, or sedatives; does not include over-the-counter drugs.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2004.