Table MH10.5 Received Mental Health Treatment/Counseling and/or Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Adults Aged 18 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse Status and Past Year Level of Mental Illness: Comparison of Percentages and Standard Errors from Restricted-Use File and Public Use File Estimates, 2010 NSDUH

Illicit Drug or Alcohol Dependence or Abuse Status/Level of Mental	MENTAL HEALTH TREATMENT OR SUBSTANCE USE TREATMENTAT A SPECIALTY FACILITY ^{1,2}							MENTAL HEALTH TREATMENT ONLY						SUBSTANCE USE TREATMENT AT A SPECIALTY FACILITY ONLY ²						MENTAL HEALTH TREATMENT AND SUBSTANCE USE TREATMENT AT A SPECIALTY FACILITY 1,2					
	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	
TOTAL	14.4	14.2	0.99	0.27	0.26	0.95	13.3	13.1	0.98	0.27	0.25	0.92	0.6	0.6	1.05	0.05	0.06	1.26	0.5	0.5	0.99	0.04	0.04	0.98	
DEPENDENCE OR ABUSE	28.4	28.2	0.99	0.97	1.14	1.17	21.0	20.7	0.99	0.88	0.95	1.08	3.4	3.6	1.04	0.38	0.43	1.16	3.9	3.9	0.98	0.42	0.42	1.00	
Any Mental Illness	48.6	47.4	0.97	1.74	2.36	1.36	36.8	35.6	0.97	1.68	1.95	1.16	2.9	3.1	1.08	0.54	0.65	1.21	8.9	8.6	0.97	1.02	1.02	1.00	
Serious Mental Illness	73.5	74.4	1.01	2.63	2.87	1.09	53.4	53.5	1.00	3.23	3.08	0.96	3.6	4.5	1.23	1.30	1.73	1.33	16.4	16.4	1.00	2.47	2.89	1.17	
Moderate Mental Illness	43.5	40.8	0.94	3.42	4.30	1.26	33.8	31.6	0.94	3.21	3.60	1.12	2.4	1.7	0.74	0.81	0.65	0.81	7.2	7.2	1.01	1.71	1.81	1.06	
Mild Mental Illness	34.3	33.1	0.96	2.49	2.71	1.09	26.9	25.9	0.96	2.47	2.72	1.10	2.6	2.9	1.11	0.69	0.86	1.25	4.7	4.3	0.90	1.08	0.59	0.55	
No Mental Illness	15.7	16.1	1.02	1.05	0.98	0.93	11.0	11.3	1.02	0.92	0.91	0.98	3.8	3.9	1.03	0.50	0.52	1.03	0.8	0.8	1.07	0.17	0.21	1.22	
NO DEPENDENCE OR ABUSE	13.0	12.8	0.98	0.28	0.28	1.00	12.5	12.3	0.98	0.28	0.28	1.00	0.3	0.3	1.06	0.04	0.05	1.17	0.2	0.2	0.98	0.02	0.03	1.11	
Any Mental Illness	42.1	41.8	0.99	1.00	0.97	0.97	40.9	40.6	0.99	1.00	0.99	0.99	0.6	0.6	1.02	0.15	0.16	1.05	0.7	0.7	0.97	0.13	0.15	1.08	
Serious Mental Illness	66.7	66.1	0.99	2.06	2.36	1.14	65.2	64.8	0.99	2.06	2.37	1.15	0.0	0.0	0.02	0.02	0.00	0.02	1.4	1.3	0.94	0.38	0.41	1.08	
Moderate Mental																									
Illness	49.7	50.0	1.01	1.99	2.40	1.21	48.1	48.3	1.00	1.97	2.34	1.19	0.8	1.0	1.19	0.32	0.38	1.21	0.7	0.7	0.97	0.33	0.43	1.32	
Mild Mental Illness	30.0	29.7	0.99	1.24	1.18	0.95	28.9	28.7	0.99	1.23	1.20	0.97	0.6	0.6	0.93	0.23	0.23	0.97	0.4	0.4	1.03	0.14	0.13	0.92	
No Mental Illness	7.4	7.2	0.97	0.25	0.23	0.91	7.1	6.9	0.97	0.25	0.23	0.93	0.2	0.3	1.07	0.04	0.04	1.07	0.1	0.1	0.98	0.01	0.02	1.10	

^{*} Low precision; no estimate reported.

PUF = public use file, which refers to estimates produced from a NSDUH public use file (a PUF is a NSDUH restricted-use file that has undergone statistical disclosure limitation treatment); RUF = restricted-use file, which refers to estimates produced from a NSDUH RUF and published in NSDUH's mental health detailed tables available at http://www.samhsa.gov/data/; SE = standard error that is associated with the estimate.

NOTE: Some 2010 RUF estimates and SEs may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings).

NOTE: Ratio of estimates = PUF estimate \div RUF estimate; Ratio of SEs = PUF SE \div RUF SE.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, and serious mental illness. Any mental illness includes individuals in any of the three categories. Estimates for mental illness presented in this table may differ from previously published mental illness estimates because of revised estimation procedures. For details on the methodology, see the NSDUH short report titled Revised Estimates of Mental Illness from the National Survey on Drug Use and Health at http://www.samhsa.gov/data/.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010.

¹ Mental Health Treatment/Counseling is defined as having received inpatient care or outpatient care or having used prescription medication for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

² Received Illicit Drug or Alcohol Treatment at a Specialty Facility refers to treatment received at a hospital (inpatient only), rehabilitation facility (inpatient or outpatient), or mental health center in order to reduce or stop drug or alcohol use, or for medical problems associated with drug or alcohol use.