

Table MH13.12 Substance Use in the Past Year and Past Month among Youths Aged 12 to 17, by Past Year Major Depressive Episode (MDE): Comparison of Percentages and Standard Errors from Restricted-Use File and Public Use File Estimates, 2013 NSDUH

Substance	TOTAL ¹						MDE						NO MDE					
	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs
PAST YEAR USE																		
Illicit Drugs ²	17.2	16.9	0.98	0.35	0.39	1.12	33.2	32.2	0.97	1.32	1.53	1.16	15.1	14.8	0.98	0.34	0.37	1.10
Marijuana and Hashish	13.4	13.2	0.98	0.31	0.37	1.16	25.7	25.3	0.98	1.22	1.45	1.19	11.8	11.5	0.98	0.31	0.34	1.09
Cocaine	0.5	0.5	1.02	0.06	0.08	1.30	1.1	1.2	1.13	0.26	0.39	1.47	0.4	0.4	0.99	0.05	0.07	1.26
Crack	0.0	0.0	0.69	0.02	0.01	0.81	0.1	0.0	0.00	0.08	0.00	0.00	0.0	0.0	0.94	0.01	0.01	1.02
Heroin	0.1	0.1	0.94	0.03	0.03	0.95	0.3	0.3	0.98	0.14	0.16	1.15	0.1	0.1	0.93	0.03	0.03	0.91
Hallucinogens	1.9	1.9	1.00	0.12	0.15	1.27	4.2	4.2	1.02	0.47	0.64	1.36	1.6	1.6	0.99	0.12	0.13	1.15
LSD	0.6	0.6	0.92	0.06	0.08	1.26	1.1	1.1	0.99	0.26	0.36	1.38	0.5	0.4	0.89	0.06	0.07	1.13
PCP	0.1	0.1	0.87	0.03	0.03	1.04	0.2	0.3	1.25	0.13	0.17	1.31	0.1	0.1	0.81	0.02	0.02	0.93
Ecstasy	0.9	1.0	1.02	0.08	0.09	1.14	2.0	2.1	1.07	0.29	0.45	1.57	0.8	0.8	1.01	0.08	0.08	1.09
Inhalants	1.9	1.9	0.96	0.12	0.12	1.04	4.8	4.2	0.88	0.55	0.48	0.88	1.5	1.5	0.99	0.11	0.11	1.01
Nonmedical Use of Psychotherapeutics ^{3,4}	5.8	5.6	0.97	0.20	0.27	1.33	12.8	11.9	0.93	0.87	0.96	1.10	4.8	4.8	0.99	0.20	0.25	1.24
Pain Relievers	4.6	4.5	0.96	0.18	0.22	1.20	10.3	9.5	0.92	0.80	0.87	1.08	3.9	3.8	0.98	0.18	0.21	1.15
OxyContin [®]	0.6	0.6	0.96	0.06	0.06	1.01	1.7	1.6	0.95	0.32	0.37	1.14	0.4	0.4	0.96	0.06	0.06	1.00
Tranquilizers	1.4	1.4	0.99	0.09	0.12	1.31	4.2	4.2	1.00	0.49	0.53	1.09	1.0	1.0	0.98	0.09	0.11	1.23
Stimulants ⁴	1.1	1.1	1.01	0.09	0.11	1.25	3.1	3.2	1.03	0.45	0.56	1.24	0.8	0.8	1.01	0.08	0.10	1.26
Methamphetamine ⁴	0.3	0.3	0.96	0.04	0.05	1.24	0.7	0.6	0.81	0.20	0.21	1.05	0.2	0.2	1.05	0.04	0.05	1.37
Sedatives	0.3	0.2	0.94	0.04	0.05	1.21	0.7	0.6	0.79	0.19	0.17	0.89	0.2	0.2	0.99	0.04	0.05	1.28
Illicit Drugs Other Than Marijuana ²	8.0	7.7	0.97	0.24	0.28	1.18	17.6	16.2	0.92	1.00	1.17	1.17	6.7	6.6	0.99	0.24	0.26	1.09
PAST MONTH USE																		
Daily Cigarette Use ⁵	1.1	1.1	0.98	0.08	0.10	1.24	1.8	1.8	1.03	0.27	0.41	1.49	1.0	0.9	0.96	0.09	0.10	1.14
Heavy Alcohol Use ⁶	1.2	1.1	0.94	0.09	0.11	1.17	1.2	1.0	0.88	0.26	0.18	0.68	1.2	1.1	0.94	0.10	0.12	1.21

* Low precision; no estimate reported.

PUF = public use file, which refers to estimates produced from a NSDUH public use file (a PUF is a NSDUH restricted-use file that has undergone statistical disclosure limitation treatment); RUF = restricted-use file, which refers to estimates produced from a NSDUH RUF and published in NSDUH's mental health detailed tables available at <http://www.samhsa.gov/data/>; SE = standard error that is associated with the estimate.

NOTE: $Ratio\ of\ estimates = PUF\ estimate \div RUF\ estimate$; $Ratio\ of\ SEs = PUF\ SE \div RUF\ SE$.

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. Respondents with unknown past year MDE data were excluded.

¹ Estimates in the Total columns represent all youths aged 12 to 17, including those with unknown past year MDE information.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 national findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

⁵ Daily Cigarette Use is defined as smoking on each of the past 30 days.

⁶ Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple hours of each other) on each of 5 or more days in the past 30 days.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013.