

Refid: 12, Skateboards: Are they really perilous? A retrospective study from a district hospital.

Rethnam U, Yesupalan RS, Sinha A.

BACKGROUND: Skateboarding has been a popular sport among teenagers even with its attendant associated risks. The literature is packed with articles regarding the perils of skateboards. Is the skateboard as dangerous as has been portrayed?

METHODS: This was a retrospective study conducted over a 5 year period. All skateboard related injuries seen in the Orthopaedic unit were identified and data collated on patient demographics, mechanism & location of injury, annual incidence, type of injury, treatment needed including hospitalisation.

RESULTS: We encountered 50 patients with skateboard related injuries. Most patients were males and under the age of 15. The annual incidence has remained low at about 10. The upper limb was predominantly involved with most injuries being fractures. Most injuries occurred during summer. The commonest treatment modality was plaster immobilisation. The distal radius was the commonest bone to be fractured. There were no head & neck injuries, open fractures or injuries requiring surgical intervention.

CONCLUSION: Despite its negative image among the medical fraternity, the skateboard does not appear to be a dangerous sport with a low incidence and injuries encountered being not severe. Skateboarding should be restricted to supervised skateboard parks and skateboarders should wear protective gear. These measures would reduce the number of skateboarders injured in motor vehicle collisions, reduce the personal injuries among skateboarders, and reduce the number of pedestrians injured in collisions with skateboarders.

 and go to or Skip to Next

Comparative Effectiveness and Safety of Pharmacologic Therapies for the Management of Crohn's Disease Abstract Review Form
WE SHOULD INCLUDE BOTH RCTS AND OBSERVATIONAL STUDIES FOR THE ABSTRACT REVIEW

Drugs of interest: ASA: sulfasalazine, mesalamine
 Anti-metabolite: azathioprine, methotrexate, 6-mercaptopurine
 Biologic: adalimumab, infliximab, certolizumab pegol, natalizumab, ustekinumab
 Corticosteroids: prednisone, prednisolone, 6-methylprednisolone, hydrocortisone, budesonide

[Click here to view KQs and exclusion/inclusion criteria](#)
1. Exclude article because... (check the first reason why study should be excluded)

- No original data (e.g., review article, commentary, or editorial)
- No subjects with Crohn's disease
- Does not evaluate a drug of interest
- Does not apply to any of the key questions (specify):
- Case report or case series
 unless hepatosplenic t-cell lymphoma or progressive multifocal leukoencephalopathy
- Other observational study or open label trial without a comparison group
 unless hepatosplenic t-cell lymphoma or progressive multifocal leukoencephalopathy
- No human data reported
- Not written in English
- Abstract only
- Other reason for exclusion

2. Unclear

- Unclear - pull article for review

3. Include article for review

- Include

4. Exclude from review, but pull for handsearching

- Pull for handsearching (e.g., systematic review that applies to the key question and published since 2005)

5. Comments (limit 250 characters)

 and go to or Skip to Next