

# ***Technical Report***

---

## **Screening for Cervical Cancer in Primary Care: A Decision Analysis for the U.S. Preventive Services Task Force**

### **Prepared for:**

Agency for Healthcare Research and Quality  
U.S. Department of Health and Human Services  
5600 Fishers Lane  
Rockville, MD 20857  
[www.ahrq.gov](http://www.ahrq.gov)

**Contract No. HHSA-290-2012-00015-I**

### **Prepared by:**

Harvard T.H. Chan School of Public Health  
Department of Health Policy and Management  
Center for Health Decision Science  
Boston, MA

### **Investigators:**

Jane J. Kim, PhD  
Emily A. Burger, PhD  
Catherine Regan, BA  
Stephen Sy, MS

# Table of Contents

|  |           |
|--|-----------|
| <b>Chapter 1. Introduction.....</b>                              | <b>1</b>  |
| <b>Chapter 2. Methods .....</b>                                  | <b>2</b>  |
| Model.....   | 2         |
| Model Calibration.....   | 3         |
| Model Validation.....  | 4         |
| Screening Strategies.....  | 4         |
| Screening Inputs .....   | 5         |
| Screening Test Characteristics.....                              | 5         |
| Colposcopy/Biopsy and Precancer Treatment .....                  | 6         |
| Outcomes .....   | 6         |
| Scenario Analysis .....  | 7         |
| Sensitivity Analysis .....                                       | 7         |
| <b>Chapter 3. Results .....</b>                                  | <b>8</b>  |
| Health Benefits and Harms.....                                   | 8         |
| Relative Efficiency Analysis .....                               | 9         |
| Summary by Efficiency Outcome .....                              | 9         |
| Summary by Screening Modality, Screening Interval, and Ages..... | 11        |
| Sensitivity Analysis .....                                       | 12        |
| Test Characteristics.....  | 12        |
| Followup of HPV-Positive Women .....                             | 14        |
| HPV-16/18 Vaccinated Women .....                                 | 15        |
| <b>Chapter 4. Discussion.....</b>                                | <b>16</b> |
| Strengths and Limitations of Modeling .....                      | 17        |
| Summary.....   | 18        |
| <b>References.....</b>   | <b>19</b> |

## Figures

- Figure 1. Model Schematic
- Figure 2. Prevalence of HPV by Age and Type
- Figure 3. Type Distribution of HPV in CIN2 and CIN3
- Figure 4. Type Distribution of HPV in Cancer
- Figure 5. Cervical Cancer Incidence per 100,000 Women, by Age and Model (Natural History)
- Figure 6. Cervical Cancer Incidence and Mortality by Age (With Screening)
- Figure 7. Outcomes From HPV-FOCAL Trial (With Screening)
- Figure 8. Flow Diagram for Alternative Triage Strategies for hrHPV-Positive Women
- Figure 9. Efficiency Frontiers: Colposcopies per Life-Year Gained, Varying HPV Testing Switch Age and Interval (Screening End Age 65 Years), Under Different Scenarios of Triage Options for HPV-Positive Women (Scenarios A-D)
- Figure 10. Efficiency Frontiers: Tests per Life-Year Gained, Varying HPV Testing Switch Age and Interval (Screening End Age 65 Years), Under Different Scenarios of Triage Options for HPV-Positive Women (Scenarios A-D)
- Figure 11. Efficiency Frontiers: Colposcopies per Cervical Cancer Case Averted, Varying HPV Testing Switch Age and Interval (Screening End Age 65 Years), Under Different Scenarios of Triage Options for HPV-Positive Women (Scenarios A-D)

Figure 12. Efficiency Frontiers: Colposcopies per Life-Year Gained When Varying Screening End Age (65, 70, and 75 Years), Under Different Scenarios of Triage Options for HPV-Positive Women (Scenarios A-D)

Figure 13. Efficiency Frontiers: Tests per Life-Year Gained When Varying Screening End Age (65, 70, and 75 Years), Under Different Scenarios of Triage Options for HPV-Positive Women (Scenarios A-D)

Figure 14. Efficiency Frontiers: Colposcopies per Cervical Cancer Case Averted When Varying Screening End Age (65, 70, and 75 Years), Under Different Scenarios of Triage Options for HPV-Positive Women (Scenarios A-D)

Figure 15. Efficiency Frontiers: Colposcopies per Life-Year Gained for All Strategies, Varying HPV Testing Switch Age, Screening Interval, and Screening End Age, Under Different Scenarios of Triage Options for HPV-Positive Women (Scenarios A-D)

Figure 16. Efficiency Frontiers: Tests per Life-Year Gained for All Strategies, Varying HPV Testing Switch Age, Screening Interval, and Screening End Age, Under Different Scenarios of Triage Options for HPV-Positive Women (Scenarios A-D)

Figure 17. Efficiency Frontiers: Colposcopies per Cervical Cancer Case Averted for All Strategies, Varying HPV Testing Switch Age, Screening Interval, and Screening End Age, Under Different Scenarios of Triage Options for HPV-Positive Women (Scenarios A-D)

## **Tables**

Table 1. Key Model Attributes

Table 2. Cervical Cancer Screening Strategies

Table 3. Screening Test Characteristics

Table 4. Distribution of Abnormal Cytology Results Conditioned on Histology Result

Table 5. Outcomes for Cervical Cancer Screening Strategies Over the Lifetime of Screening (Screening End Age 65 Years)

Table 6. Outcomes for Cervical Cancer Screening Strategies Over the Lifetime of Screening (Screening End Age 70 Years)

Table 7. Outcomes for Cervical Cancer Screening Strategies Over the Lifetime of Screening (Screening End Age 75 Years)

Table 8. Sensitivity Analysis Summary: Impact of Uncertainty on Colposcopies per Life-Year Gained

Table 9. Sensitivity Analysis Summary: Impact of Uncertainty on Tests per Life-Year Gained

Table 10. Sensitivity Analysis Summary: Impact of Uncertainty on Colposcopies per Cervical Cancer Case Averted

## **Appendix Tables**

Appendix Table 1. Natural History Model Parameters, Postcalibration

Appendix Table 2. Efficient Cervical Cancer Screening Strategies (Screening End Age 65 Years) in Terms of Colposcopies per Life-Year Gained

Appendix Table 3. Efficient Cervical Cancer Screening Strategies (Screening End Age 65 Years) in Terms of Tests per Life-Year Gained

Appendix Table 4. Efficient Cervical Cancer Screening Strategies (Screening End Age 65 Years) in Terms of Colposcopies per Cervical Cancer Case Averted

Appendix Table 5. Efficient Cervical Cancer Screening Strategies (Screening End Age 70 Years) in Terms of Colposcopies per Life-Year Gained

Appendix Table 6. Efficient Cervical Cancer Screening Strategies (Screening End Age 70 Years) in Terms of Tests per Life-Year Gained

Appendix Table 7. Efficient Cervical Cancer Screening Strategies (Screening End Age 70 Years) in Terms of Colposcopies per Cervical Cancer Case Averted

Appendix Table 8. Efficient Cervical Cancer Screening Strategies (Screening End Age 75 Years) in Terms of Colposcopies per Life-Year Gained

Appendix Table 9. Efficient Cervical Cancer Screening Strategies (Screening End Age 75 Years) in Terms of Tests per Life-Year Gained

Appendix Table 10. Efficient Cervical Cancer Screening Strategies (Screening End Age 75 Years) in Terms of Colposcopies per Cervical Cancer Case Averted

Appendix Table 11. Efficient Cervical Cancer Screening Strategies, Varying HPV Testing Switch Age, Screening Interval, and Screening End Age in Terms of Colposcopies per Life-Year Gained

Appendix Table 12. Efficient Cervical Cancer Screening Strategies, Varying HPV Testing Switch Age, Screening Interval, and Screening End Age in Terms of Tests per Life-Year Gained

Appendix Table 13. Efficient Cervical Cancer Screening Strategies, Varying HPV Testing Switch Age, Screening Interval, and Screening End Age in Terms of Colposcopies per Cervical Cancer Case Averted

Appendix Table 14. Range of Outcomes for Cervical Cancer Screening Strategies Over Lifetime of Screening (Screening End Age 65 Years) Across 50 Top-Fitting Parameter Sets

Appendix Table 15. Outcomes for Cervical Cancer Screening Strategies Over Lifetime of Screening (Screening End Age 65 Years), Assuming Cytology Sensitivity of 81.5% and Specificity of 88.0%

Appendix Table 16. Outcomes for Cervical Cancer Screening Strategies Over Lifetime of Screening (Screening End Age 65 Years), Assuming Cytology Sensitivity of 51.4% and Specificity of 93.6%

Appendix Table 17. Outcomes for Cervical Cancer Screening Strategies Over Lifetime of Screening (Screening End Age 65 Years), Assuming Lower HPV Sensitivity (Relative Sensitivity of 1.15)

Appendix Table 18. Outcomes for Cervical Cancer Screening Strategies Over Lifetime of Screening (Screening End Age 65 Years), Assuming Imperfect Colposcopy/Biopsy Performance

Appendix Table 19. Outcomes for Cervical Cancer Screening Strategies Over Lifetime of Screening (Screening End Age 65 Years), Assuming CIN2/3 Treatment Effectiveness of 82%

Appendix Table 20. Outcomes for Cervical Cancer Screening Strategies Over Lifetime of Screening (Screening End Age 65 Years), Assuming Referral to Colposcopy for LSIL or Worse Result on Cytology Triage

Appendix Table 21. Outcomes for Cervical Cancer Screening Strategies Over Lifetime of Screening (Screening End Age 65 Years), Assuming 6-Month Followup for HPV-Positive, Cytology-Negative Women

Appendix Table 22. Outcomes for Cervical Cancer Screening Strategies Over Lifetime of Screening (Screening End Age 65 Years), Assuming 24-Month Followup for HPV-Positive, Cytology-Negative Women

Appendix Table 23. Outcomes for Cervical Cancer Screening Strategies Over Lifetime of Screening (Screening End Age 65 Years), Assuming Colposcopy Referral for All HPV-Positive Women

Appendix Table 24. Outcomes for Cervical Cancer Screening Strategies Over Lifetime of Screening (Screening End Age 65 Years) in Women Vaccinated Against HPV-16/18

# Chapter 1. Introduction

Despite dramatic reductions in cervical cancer since the introduction of Papanicolaou (Pap) cytology testing in the United States, roughly 12,820 women are expected to develop and 4,210 women are expected to die from cervical cancer in 2017.<sup>1</sup> Information on the natural history of human papillomavirus (HPV) and its causal role in cervical disease, coupled with technologies to improve detection of precancerous lesions, continue to emerge and have prompted revisions to screening guidelines. In 2012, cervical cancer screening guidelines were harmonized across several major guidelines-making organizations, including the U.S. Preventive Services Task Force (USPSTF),<sup>2-4</sup> with recommendations for routine cytology screening every 3 years starting at age 21 years, with an option to switch to cytology and HPV “cotesting” every 5 years starting at age 30 years. Screening end age is recommended at age 65 years, provided a history of regular screening without abnormalities in the past 10 to 20 years.<sup>2-4</sup> Since 2012, new evidence on primary HPV testing has emerged, contributing to the U.S. Food and Drug Administration (FDA) approval of the first standalone HPV test for primary screening in women age 25 years and older. Interim clinical guidance on the use of primary HPV testing has been issued from several professional organizations.<sup>5</sup>

While empirical studies such as randomized clinical trials provide high-quality evidence on the effectiveness of screening, outcomes are usually based on intermediate endpoints after a limited number of rounds of screening. Mathematical disease simulation models can complement such evidence by extrapolating data beyond the trial period to project the benefits and harms of screening in the long term, over multiple rounds. Models can also explore the impact of alternative scenarios that have not been examined in empirical studies.

This decision analysis using a cervical cancer disease simulation model accompanies the systematic review that is being conducted by the Kaiser Permanente Evidence-based Practice Center (EPC) to update the evidence and address gaps in the expected benefits and harms of cervical cancer screening strategies in primary care.<sup>6</sup> The key questions for the decision analysis center around the long-term impact of primary HPV screening compared to current guidelines-based screening strategies in terms of health benefits and harms in the general population:

1. How does the effectiveness of primary HPV screening in reducing cervical cancer incidence and mortality vary by 1) age to start HPV screening, 2) rescreening interval (following an HPV-negative result), and 3) age to stop HPV screening?
2. How do the harms of primary HPV screening vary by 1) age to start HPV screening, 2) rescreening interval, and 3) age to stop HPV screening?
3. Which cervical cancer screening strategies are considered efficient?

Two variations of triage of HPV-positive women were included in the base case. While the key questions of the analysis focused on primary HPV testing alone, we also considered other variations of cotesting. In addition, sensitivity analysis was conducted to assess the impact of uncertainty in the data and alternative screening scenarios.

# Chapter 2. Methods

## Model

An overview of the decision model in terms of model attributes, natural history, and screening strategies is provided below and summarized in **Table 1**. The model is a microsimulation (i.e., individual-based) model of HPV-induced cervical carcinogenesis, in which individual women enter the model at an early age (i.e., age 9 years) and are followed over their lifetimes.<sup>7,8</sup> The main health states of the model comprise HPV infection (by genotype), precancer (i.e., cervical intraepithelial neoplasia [CIN], grades 2 and 3) and invasive cancer (by stage) (**Figure 1**). The model focuses on squamous cell carcinoma (SCC), the most common histologic subtype of cervical cancer.

The current analysis was conducted using a single hypothetical birth cohort assumed to be born in year 1996 that would begin screening at age 21 years in year 2017. Screening is used to detect the presence of high-grade precancer, which may resolve spontaneously or can be treated and removed before progressing to cancer; therefore, reductions in cervical cancer morbidity and mortality due to screening result from both the prevention and the earlier detection of invasive cancer. The effectiveness of screening strategies depends on screening ages, screening interval, test characteristics, management of abnormal/positive results, and treatment efficacy. The model was used to project estimates of both benefits and harms, including life-years gained, cervical cancer cases and deaths, screening tests, diagnostic procedures, and false-positive results under various scenarios of primary screening tests, screening ages, screening intervals, and triage algorithms for screen-positive women. These measures are calculated as the cumulative number of events or time spent in the different health states, which are then modified by the interventions, over the selected time horizon (e.g., lifetime). These measures in totality capture the benefits and harms for the strategies being considered. To examine the relative tradeoff of harms versus benefits among the strategies, we calculated three efficiency outcomes in terms of the incremental number colposcopies per life-year gained, incremental number of screening tests per life-year gained, and incremental number of colposcopies per cervical cancer case averted.

## Natural History

Upon entry into the model, each woman faces monthly transitions between health states that describe underlying true health, including HPV infection, precancer (i.e., CIN2 and CIN3), and invasive cancer (i.e., local, regional, and distant). CIN2 and CIN3 are modeled as nonsequential precancerous health states with distinct probabilities of regression to normal or progression to cancer, whereas CIN1 is interpreted as a microscopic manifestation of acute HPV infection and is therefore incorporated into the HPV-infected state. States are further stratified into oncogenic HPV types 16, 18, 31, 33, 45, 52, and 58, each considered separately; pooled other high-risk types; and pooled low-risk HPV types. Transition probabilities can vary by age, HPV type, duration of infection or lesion, and history of prior HPV infection and CIN treatment. Cancer detection can occur through symptoms or screening. Each month, all women are subjected to rates of hysterectomy and all-cause mortality<sup>9,10</sup> and excess mortality from cervical cancer.<sup>10</sup>

## Screening, Diagnosis, and Precancer Treatment

Screening assumptions in the model can vary by screening start age, stop age, interval between negative screens, coverage, triage testing, and compliance to recommended followup. Tests for primary screening and triage include cytology (conventional or liquid-based), and high-risk HPV DNA testing (pooled or genotyping), as well as cytology and HPV cotesting. Management of screen-positive women can vary by age, followup test, time to followup test(s), and number of negative followup tests required to return to routine screening.

## Cancer Treatment and Survival

Cancer staging (i.e., local to regional to distant) and progression is modeled, accounting for symptomatic detection and the possibility of downstaging at diagnosis due to screening. In addition to all-cause mortality, women with cervical cancer are subject to excess mortality, based on 5-year survival estimates depending on cancer stage, age, and time since diagnosis according to Surveillance, Epidemiology, and End Results (SEER) data.<sup>11</sup>

## Model Calibration

A process of model calibration and validation was undertaken to ensure fit to observed data. Calibrated parameters included HPV incidence (by age and genotype), CIN progression and regression, and HPV natural immunity following type-specific HPV infection and clearance. Baseline values for each of the uncertain parameters were randomly selected from a predetermined plausible range, creating a unique natural history parameter set. Goodness of fit was ascertained by calculating the likelihood of model-projected outcomes from each parameter set against corresponding calibration targets. To capture uncertainty in the natural history parameters, the 50 best-fitting sets were used in all analyses; the results are reported as the mean value across the 50 sets (the minimum and maximum values are also reported in a sensitivity analysis). The calibrated model parameter values used in this analysis are summarized in **Appendix Table 1**.<sup>12-22</sup>

For calibration target data, the data sources were selected on the basis of representativeness of the general population, sampling methods, and sample size. All data were from populations prior to HPV vaccination. Age-specific prevalence of HPV infections was based on data from the New Mexico HPV Pap Registry (NMHPVPR), the only statewide screening registry in the United States.<sup>23</sup> The model was fitted to prevalence of HPV types 16, 18, 31, 33, 45, 52, and 58 separately, as well as other pooled high-risk HPV types. HPV type distribution in cases of CIN and cancer were also included as calibration target data. For CIN2 and CIN3, HPV type distribution was based on data from the NMHPVPR;<sup>24</sup> for cancer, HPV type distribution in SCC was based on a recent study by the Centers for Disease Control and Prevention using tissue samples from U.S. population-based cancer registries.<sup>25</sup> Model fit to calibration targets are displayed in **Figures 2–4**.



## Model Validation

Age-specific cervical cancer incidence rates under an assumption of no intervention (i.e., natural history) were projected by the model and compared against cancer registry data from the 1950s and early 1960s, before Pap smear screening was widely performed (**Figure 5**).<sup>26,27</sup> Given the limited data from only a few states (Connecticut, New York, and Hawaii), and the potential changes in sexual behavior and other risk factors since the prescreening era, these data were not used directly to calibrate the model but instead were used to assess predictive validity for overall underlying risk.

In addition, model-projected outcomes of cervical cancer incidence and mortality rates were compared against those reported in SEER cancer registries in recent years (i.e., 2000–2012), under assumptions of screening practice patterns reported in the NMHPVPR (**Figure 6**).<sup>8,28-30</sup> Screening practice patterns included estimated proportions of women never screened and screened at different intervals (e.g., annual or biennial) and proportions of women who do not comply to followup diagnostic testing, precancer treatments, or both.

Additional model validation exercises included simulating the protocol from the HPV-FOCAL trial and comparing model projections against reported outcomes.<sup>31</sup> We simulated three screening scenarios involving switching to primary HPV testing at ages 27, 34, and 52 years to reflect women who switch from cytology-based routine screening to primary HPV testing under the trial screening protocol. We projected the baseline and cumulative 12-month CIN2+, CIN3+, and colposcopy rates per 1,000 women in the first round of the HPV and control arms of the HPV-FOCAL trial for ages 25–29, 30–34, and 35–65 years (**Figure 7**). Comparisons of model projections against observational data have been previously reported.<sup>8</sup>

## Screening Strategies

The analysis focused on the comparative effectiveness and harms of primary HPV testing, compared to currently recommended screening strategies, as well as other variations of cotesting. **Table 2** summarizes the 19 main strategies evaluated. Guideline-based screening strategies comprised cytology alone every 3 years from ages 21 to 65 years (strategy 1) and cytology alone every 3 years from ages 21 to 29 years, with a switch to cytology and HPV cotesting every 5 years from ages 30 to 65 years (strategy 2).<sup>2-4</sup> Management of women with equivocal or abnormal tests was assumed to follow established guidelines.<sup>3,32</sup> For cotesting, HPV-positive/cytology-negative women were managed by repeat cotesting at 12 months, with referral to colposcopy for any positive result.

The primary HPV testing strategies (strategies 3–14) were varied by: 1) age to switch from cytology to HPV screening, 2) rescreening interval following an HPV-negative result, and 3) triage options for an HPV-positive result. For the base case, *age to switch to HPV screening* was evaluated at ages 25, 27, and 30 years, following cytology-only screening starting at age 21 years. Age 25 years was selected to reflect the FDA-approved age threshold for primary HPV testing; age 27 years was selected to coincide with timing of 3-year cytology testing (at ages 21 and 24 years); and age 30 years was selected to be consistent with the age threshold to begin

cotesting. The *rescreening interval* for primary HPV testing was evaluated at every 3 years and every 5 years, consistent with current U.S. guidelines for cytology-only screening and cotesting. Two triage strategies for HPV-positive screening results were examined (**Figure 8**): a) assuming HPV-16/18 genotype information is available, HPV 16/18-positive women are referred to colposcopy, whereas women positive for other high-risk HPV types receive cytology triage (those with a cytology result of atypical squamous cells of undetermined significance [ASC-US] or worse are referred to colposcopy; those with a cytology-negative result receive a followup test in 12 months); and b) all women with high-risk HPV receive cytology triage. A referral threshold of a cytology result of low-grade squamous intraepithelial lesion (LSIL) or worse was also evaluated, and the interval for followup testing was varied (e.g., 6 or 24 months) in sensitivity analysis. An additional triage strategy in which all women with high-risk HPV are referred for immediate colposcopy was also included in sensitivity analysis. Additional cotesting strategies (strategies 15–19) were included, varying the age to switch and rescreening interval.

In the base-case analysis, age to stop screening was 65 years, assuming no recent history of abnormal results, consistent with current guidelines; we evaluated the impact of extending the age threshold at which to terminate screening to ages 70 and 75 years. We assumed full compliance to screening initiation, rescreening interval, and followup for both diagnostic and precancer treatment referrals. Furthermore, the base-case analysis focused on women who did not receive HPV vaccination.

## Screening Inputs

### Screening Test Characteristics

Test sensitivity and specificity values, defined at a disease threshold of CIN2 or worse (CIN2+), were required as model inputs and were informed from the studies reviewed by the EPC,<sup>6</sup> as well as from the published literature (**Table 3**).<sup>33-37</sup> For strategies involving cytology testing, we applied estimates of test sensitivity and specificity assuming a positivity threshold of ASC-US, with base-case values obtained from a meta-analysis conducted by Koliopoulos et al.<sup>33</sup> These pooled estimates of sensitivity and specificity of cytology for detection of CIN2+ (72.7% and 91.9%, respectively) were based on 18 studies identified in a systematic review. The ranges of test performance values for cytology were informed by the Koliopoulos study, as well as estimates reported in the recent U.S.-based Addressing the Need for Advanced HPV Diagnostics (ATHENA) study.<sup>34</sup>

The model also required data on the distribution of abnormal cytology results (e.g., proportion of women with ASC-US, LSIL, and high-grade squamous intraepithelial lesion result) conditional on histologic diagnosis (**Table 4**). These estimates were based on data from the ATHENA study, which reported the baseline cytology results by central pathology review diagnosis of 8,000 women age 25 years and older.<sup>38</sup>

The main inputs of test performance for primary HPV testing, alone and as part of cotesting, were based on ATHENA and studies that were included in EPC review.<sup>34-37</sup> Given the wide variation in absolute test characteristics across studies due to differences in protocols and

populations, we elected to utilize relative sensitivity and specificity values, compared with cytology testing (positivity threshold of ASC-US+) (**Table 3**). Our base-case estimates were anchored on the ATHENA study,<sup>34</sup> which provided verification bias-adjusted estimates and included both HPV (cobas HPV test [Roche]) and cotesting strategies with similar followup algorithms as what was evaluated in this analysis. The worst- and best-case values for sensitivity and specificity for all screening test modalities were informed by data reported in ATHENA,<sup>34</sup> a meta-analysis by Arbyn et al,<sup>35</sup> and the New Technology in Cervical Cancer (NTCC) trial,<sup>36,37</sup> which reflect variations across testing modalities that include the cobas HPV test, Hybrid Capture 2 (Qiagen), and polymerase chain reaction-based tests.

## Colposcopy/Biopsy and Precancer Treatment

The sensitivity and specificity of colposcopy and biopsy were assumed to be perfect (100%) in the base case, although we explored the impact of error in histologic diagnosis using data from the NMHPVPR in sensitivity analysis.<sup>39</sup> We assumed that, with active surveillance of women who receive precancer treatment, the effectiveness of treatment in removing a CIN2 or CIN3 lesion (e.g., via loop electrosurgical excisional procedure [LEEP]) is ultimately 100%, but also explored a lower treatment effectiveness of 82%.<sup>40</sup> The model assumed that HPV infections are also removed with precancer treatment, and therefore, treated women return to an uninfected state.

## Outcomes

The model generated a number of outcomes associated with each screening strategy, reflecting both health effects and harms over the lifetime of the screening cohort (i.e., ages 21 to 100 years): total number of cytology and HPV tests (including screening, triage, and surveillance), colposcopies, CIN2 and CIN3 detected, CIN3+ detected (including CIN3 and cervical cancers detected through screening), false-positive screening results (defined as total number of colposcopies without underlying CIN2, CIN3, or cancer), cervical cancer cases, cervical cancer deaths, and life-years.

The relative efficiency of each screening strategy was evaluated and expressed as the incremental number of colposcopies per life-year gained, defined as the additional number of colposcopies divided by the additional life-years of a specific strategy (strategy x) compared to the strategy with the next fewer colposcopies (strategy y):

$$\frac{Colposcopies_{Strat\ x} - Colposcopies_{Strat\ y}}{Life-years_{Strat\ x} - Life-years_{Strat\ y}}$$

Strategies with a higher number of colposcopies and lower life-years than an alternative strategy were considered “inefficient” and eliminated from the calculation; all other strategies were considered “efficient.” Because there is no consensus on the appropriate metric to assess efficiency, we also presented results in terms of the incremental number of total screening tests per life-year gained and the incremental number of colposcopies per cervical cancer case averted.

## Scenario Analysis

For primary HPV testing, the different options for triaging HPV-positive women are not equally available or preferred; therefore, analyses to determine the relative efficiency of the screening strategies were conducted under four different scenarios of triage availability:

- **Scenario A:** Included only 16/18 genotype triage option for HPV-positive women (strategies 3–8)
- **Scenario B:** Included only cytology triage option for HPV-positive women (strategies 9–14)
- **Scenario C:** Included both 16/18 genotyping and cytology triage options for HPV-positive women (strategies 3-14)
- **Scenario D:** Included both 16/18 genotyping and cytology triage options for HPV-positive women, plus additional cotesting strategies (strategies 3–19)

Each scenario analysis above also included the current guidelines-based strategies of cytology testing alone every 3 years, with or without a switch to cotesting every 5 years at age 30 years (strategies 1 and 2).

## Sensitivity Analysis

We assessed the impact of uncertainty in the data, alternative screening management protocols, and screening in HPV-vaccinated women. Data uncertainty focused on the underlying natural history of disease (i.e., transition probabilities), screening test characteristics, colposcopy/biopsy performance, and precancer treatment effectiveness. Alternative screening scenarios included variations in management of HPV-positive women, including cytology triage with a colposcopy referral threshold of LSIL (base case assumed ASC-US), varying intervals for followup testing from 6 to 24 months (base case assumed 12 months), and immediate colposcopy for all HPV-positive women. To reflect a low-risk population, we evaluated screening in HPV-vaccinated women. We assumed women were fully vaccinated with the three-dose HPV-16/18 vaccine in preadolescence and that vaccination conferred 100% protection against HPV-16 and HPV-18 infections over the lifetime.

# Chapter 3. Results

## Health Benefits and Harms

In the absence of screening, the lifetime risk of cervical cancer was 1.9% (range across 50 best-fitting sets, 1.3% to 2.4%) and lifetime mortality from cervical cancer was 0.83% (range, 0.58% to 1.08%), resulting in a life expectancy of 63.921 years (range, 63.845 to 64.006 years) for 20-year-old women. Under scenarios of screening, model outcomes of screening tests (cytology, HPV, and total), colposcopies, CIN2/CIN3 detected, CIN3+ detected, false positives, cervical cancer cases, cervical cancer deaths, and life-years per 1,000 women were projected, separately for screening end ages of 65, 70, and 75 years (**Tables 5–7**). Compared to no screening, all cervical cancer screening strategies led to substantial reductions in cancer cases and deaths and gains in life-years. Compared to the current guidelines-based strategies of cytology alone every 3 years starting at age 21 years, with or without a switch to cotesting every 5 years starting at age 30 years, all new alternative strategies were more effective. For example, when screening ended at age 65 years (**Table 5**), cervical cancer deaths associated with the guidelines-based strategies (strategies 1 and 2) ranged from 0.30 to 0.76 deaths per 1,000 women, whereas the new alternative strategies involving primary HPV testing or cotesting (strategies 3–19) had fewer cervical cancer deaths, ranging from 0.23 to 0.29 deaths per 1,000 women.

Earlier switch age, more frequent intervals, and cotesting strategies generally led to a greater number of lifetime total tests. The proportion of tests that were cytology versus HPV test depended on the particular strategy. Screening strategies that involved switching to primary HPV testing alone (strategies 3–14) were associated with a substantially lower total number of cytology tests—3 to 8 times fewer cytology tests than with cytology alone or cotesting. In contrast, strategies involving HPV testing alone with 16/18 genotype triage (strategies 3–8) had the highest number of HPV tests, followed closely by HPV testing alone with cytology triage (strategies 9–14) and cotesting (strategies 2 and 15–19); cytology only, which utilizes HPV testing only for triage of ASC-US, had the lowest number of HPV tests. In total, the cotesting strategies had nearly double the total tests of the primary HPV testing strategies. The HPV testing strategy with cytology triage (strategies 9–14) led to a slightly higher number of tests than HPV testing with 16/18 genotyping (strategies 3–8). Continually screening with cytology alone every 3 years (strategy 1) had the fewest number of total tests than other HPV testing and cotesting strategies involving 3-year screening.

Cytology testing alone every 3 years also yielded the lowest number of lifetime colposcopies (i.e., 645 per 1,000 women). All other strategies increased colposcopies substantially, ranging from 1,452 to 2,535 per 1,000 women—up to 4-fold higher when screening with primary HPV testing or cotesting every 3 years starting at age 25 years. All else equal, colposcopies were generally highest for cotesting (strategies 2 and 15–19), followed closely by primary HPV testing with 16/18 genotype triage (strategies 3–8). HPV testing with 16/18 genotype triage had 12% to 14% greater colposcopies than HPV testing with cytology triage.

Likewise, the lowest numbers of CIN2/CIN3 and CIN3+ detected were from cytology testing alone every 3 years; for example, 160 CIN2/CIN3 cases detected per 1,000 women (strategy 1)

versus 198 to 223 CIN2/CIN3 cases detected per 1,000 women for the strategies involving primary HPV testing and cotesting (strategies 2–19). Cotesting strategies (strategies 2 and 15–19) yielded the highest numbers of CIN2/CIN3 and CIN3+ detected, but only marginally higher than primary HPV testing with 16/18 genotype triage (strategies 3–8), followed by primary HPV testing with cytology triage (strategies 9–14). Mimicking the trend of colposcopies, the number of false positives increased dramatically from cytology testing every 3 years to HPV testing or cotesting, irrespective of screening switch age or interval, up to 5-fold greater with cotesting or HPV testing every 3 years starting at age 25 years (strategies 3, 9, and 15).

The current guidelines-based strategy of cytology alone every 3 years (strategy 1) was the least effective in terms of cervical cancer cases, deaths, and life-years. Across the testing modalities, the most effective strategy was primary HPV testing with 16/18 genotype triage (strategies 3–8); the lowest number of cancer cases and deaths, and highest number of life years, occurred with 3-year screening with a switch age to primary HPV testing of 25 years (strategy 3). However, the difference in benefit between strategies involving primary HPV testing with 16/18 genotype triage compared to HPV testing with cytology triage and cotesting were quite small, especially at earlier switch ages and more frequent intervals.

As the screening end age extended to age 70 and 75 years (**Tables 6 and 7**), the absolute number of screening tests, colposcopies, CIN2/CIN3 detected, CIN3+ detected, false positives, and life-years increased, while the cervical cancer cases and deaths decreased. The trends in outcomes between the strategies (i.e., by screening modality, switch age, and interval) remained consistent irrespective of screening end age.

## Relative Efficiency Analysis

Three different metrics of colposcopies per life-year gained, screening tests per life-year gained, and colposcopies per cancer case averted were calculated to reflect different tradeoffs in harms and benefits. Each scenario analysis (A–D), representing different assumptions of the availability of triage strategies for primary HPV testing, included the current guidelines-based strategies of cytology testing alone every 3 years, with or without a switch to cotesting every 5 years at age 30 years (**Figures 9–11 and Appendix Tables 2–4**). These analyses were repeated assuming extension of the screening end age to 70 and 75 years (**Figures 12–14 and Appendix Tables 5–10**), and a comprehensive analysis included all screening strategies, varying screening test modality, triage approach, age to switch to HPV primary testing, interval, and age to end screening (**Figures 15–17 and Appendix Tables 11–13**).

## Summary by Efficiency Outcome

### Colposcopies per Life-Year Gained

In all four scenarios of HPV triage and cotesting availability (**Figure 9 and Appendix Table 2**), the strategy with the lowest number of colposcopies per life-year gained was the current guidelines-based strategy of cytology testing alone every 3 years (strategy 1), with 3 colposcopies per life-year gained compared to no screening. By comparison, primary HPV and

cotesting strategies increased both life-years and number of colposcopies. Efficient strategies included primary HPV testing, either with 16/18 genotyping or cytology triage, every 5 years with a switch age of 25, 27, and 30 years (strategies 6–8 and 12–14). For example, with 16/18 genotype triage availability (Scenario A), number of colposcopies per life-year gained increased from 86 to 297, as the switch age from cytology to HPV primary testing decreased from 30 to 25 years. Switching to 3-year primary HPV testing at age 25 years (strategy 3), the most effective strategy, had a substantially higher ratio of 2,082 colposcopies per life-year gained, compared to 5-year screening at the same switch age of 25 years (strategy 6). When assuming cytology triage for HPV-positive women (Scenario B), the corresponding ratios were lower for every 5-year screening with HPV primary testing, but slightly higher for every 3-year screening. In both Scenarios C and D, which included both triage strategies for HPV-positive women, followup with 16/18 genotype triage was not efficient when compared against cytology triage when HPV primary testing was conducted every 5 years. Screening every 3 years with either triage strategy required a much greater number of colposcopies per life-year gained, ranging from 2,188 (cytology triage, strategy 9) to 3,822 (16/18 genotype triage, strategy 3). The guidelines-based cotesting strategy (strategy 2) and additional cotesting strategies with earlier switch ages (25 and 27 years) and greater frequency (3-year interval) were not efficient in any scenario (strategies 15–19).

### **Tests per Life-Year Gained**

When the analysis was expressed in terms of tests (both cytology and HPV) per life-year gained, the only efficient strategies were primary HPV testing at a switch age of 25 years, with either 5- or 3-year screening (strategies 3, 6, 9, and 12) (**Figure 10** and **Appendix Table 3**). In Scenario A, HPV primary testing with 16/18 genotype triage every 5 years (strategy 6) was associated with 43 tests per life-year gained; the ratio for this same strategy increased substantially to 22,335 tests per life-year gained with every 3-year screening (strategy 3). In Scenario B, with cytology triage, the corresponding ratios increased slightly to 44 tests per life-year gained with 5-year screening (strategy 12), and 28,636 tests per life-year gained with 3-year screening (strategy 9). When both triage options were available (Scenarios C and D), 16/18 genotyping was more efficient than cytology triage. Cytology only and cotesting strategies (strategies 1, 2, and 15–19) were not efficient in any of the scenarios compared to primary HPV testing.

### **Colposcopies per Cervical Cancer Case Averted**

Efficient strategies were consistent with those identified in the analysis of colposcopies per life-year gained (**Figure 11** and **Appendix Table 4**). Across all scenarios, cytology-only screening every 3 years (strategy 1) had the lowest ratio of 39 colposcopies per case averted. In Scenario A, primary HPV testing every 5 years with a switch age of 30 years (strategy 8) was associated with a ratio of 766 colposcopies per case averted; shifting the age of switching from cytology to primary HPV testing required a greater number of colposcopies per case averted (1,432 for switch age 27 years and 2,120 for switch age 25 years). The most effective strategy, HPV testing every 3 years with a switch age of 25 years (strategy 3), increased the ratio to 8,580 colposcopies per case averted. By comparison, when cytology was the only triage option for HPV-positive women (Scenario B), the corresponding ratios were uniformly lower, ranging from 640 to 1,735 colposcopies per case averted for HPV testing every 5 years at switch ages of 30 to 25 years,

respectively; 3-year primary HPV testing with a switch age of 25 years (strategy 9) was associated with 7,018 colposcopies per case averted. When both triage options were equally available, we found that the strategies involving cytology triage for HPV-positive women were more efficient than 16/18 genotyping; 3-year HPV testing at age 25 years with 16/18 genotyping (strategy 3), the most effective strategy, had a much higher ratio of 23,974 colposcopies per case averted. In all scenarios, cotesting strategies (strategies 2 and 15–19) were not efficient.

## **Summary by Screening Modality, Screening Interval, and Ages**

### **Screening Modality and Triage**

Across all three efficiency outcomes, strategies involving primary HPV testing, with either 16/18 genotype or cytology triage (depending on the scenario), consistently remained on the efficiency frontier. When the two triage strategies were compared in the same analysis (Scenarios C and D), cytology triage was more efficient than 16/18 genotype triage for the two efficiency metrics that used colposcopy as a measure of harm (per life-year gained and per case averted); however, 16/18 genotype testing was the preferred triage option when using screening tests as the measure of harm (per life-year gained).

Cytology only, reflecting a currently recommended strategy, had the lowest benefit in terms of life-years and cancer cases, as well as the lowest number of colposcopies, and therefore yielded the lowest ratios when considering colposcopies as the measure of harm. When instead considering screening tests, cytology only was no longer on the efficiency frontier. Strategies involving cotesting, including one that is current recommended in the United States, were universally not efficient across any of the measures.

### **Screening Interval**

Strategies involving 5-year screening were much more efficient than strategies with 3-year screening, which either were not on the efficiency frontier or had exceedingly high (i.e., unattractive) harm-to-benefit ratios. For all three efficiency measures, strategies involving switching to primary HPV testing every 5 years remained on the efficiency frontier.

### **Age to Switch From Cytology-Only Screening**

For efficiency outcomes using colposcopies as a measure of harm, switching from cytology only to primary HPV testing at ages 25, 27, and 30 years were found to be efficient, and the harm-to-benefit ratios decreased (i.e., became more attractive) as the switch age extended from age 25 to 30 years. When using screening tests as the measure of harm, only two strategies were efficient, both involving switching to primary HPV testing at age 25 years (5- and 3-year intervals).

### **Age to End Screening**

When the analyses were repeated with the age to end screening extended to 70 and 75 years (base case, 65 years), we found that our findings were very robust and that all of the same strategies were on the efficiency frontier, as in the base-case analysis for each of the three



efficiency outcomes (**Figures 12–14** and **Appendix Tables 5–10**). The corresponding ratios increased (i.e., became less attractive) as the end age increased, indicating that although screening is more effective when continued to later ages, it also becomes less efficient.

When we conducted a comprehensive analysis with all possible strategies, including varying screening end age, we found that most of the strategies on the efficiency frontier involved extending screening end age to 70 and 75 years (**Figures 15–17** and **Appendix Tables 11–13**). When using colposcopies as a measure for harms under Scenario D, screening with cytology alone every 3 years (strategy 1) was efficient when screening ended at 65, 70, and 75 years; ratios for colposcopies per life-year gained were 3, 17, and 26, respectively. Next efficient strategies included switch to 5-year primary HPV testing with cytology triage at age 30 years (strategy 14), with screening end ages of 70 and 75 years, which increased colposcopies per life-year gained to 95 and 99 colposcopies per life-year gained, respectively. Those strategies with earlier switch ages (27 and 25 years, strategies 13 and 12) increased the number of colposcopies required per life-year gained to 135 and 225, respectively. Three-year HPV testing with 16/18 genotype triage at a switch age of 25 years (strategy 3) until age 75 years was the most effective strategy, with the highest ratio of 6,239 colposcopies per life-year gained. When considering colposcopies per case averted, nearly all of the same strategies were efficient, except that the screening end age for primary HPV testing was 75 years. Strategies with a switch to HPV testing before age 30 years required greater than 1,000 colposcopies per case averted; strategies with a switch age of 25 years had ratios ranging from 2,064 to 25,112 colposcopies per case averted, depending on triage strategy and interval.

When using screening tests as a measure of harms, we again found that the only efficient strategies involved primary HPV testing with 16/18 genotype triage at a switch age of 25 years (strategies 3 and 6). Extending the end age of screening increased efficiency ratios dramatically, from 43 to 707 to 1,497 screening tests per life-year gained when screening occurred every 5 years up to ages 65 to 70 to 75 years, respectively. This same strategy occurring every 3 years up to age 75 years required 69,064 screening tests per life-year gained.

## Sensitivity Analysis

Because of the inherent uncertainty in key model parameters, we undertook several sensitivity analyses to explore the robustness of results. The range (minimum and maximum) of base-case results across the 50 best-fitting parameter sets are presented in **Appendix Table 14** to show the variation in outcomes when taking into account the uncertainty in the natural history parameters. Despite these variations, the rank order of the strategies according to each outcome was stable over the multiple sets.

## Test Characteristics

We undertook several analyses to assess the impact of test performance characteristics on base-case results (Scenario D) (**Tables 8–10** and **Appendix Tables 15–19**). When test sensitivity for cytology was increased to the upper-bound, best-case value (81.5%), with a corresponding decrease in specificity (88.0%), we found that both cervical cancer cases and deaths decreased

for all strategies, with the biggest decrease in the cytology-only strategy (decrease of 15% in cervical cancer cases and 13% in cervical cancer deaths, strategy 1) (**Appendix Table 15**). Nonetheless, because of the lower specificity, both numbers of colposcopies and false positives increased considerably, whereas the number of tests only increased marginally. For all three efficiency metrics, the efficient strategies remained the same as in the base case, but the ratios generally increased (i.e., became less attractive) given the increase in resource use. For example, ratios associated with a switch to 5-year primary HPV testing at ages 30, 27, and 25 years increased to 80, 167, and 323 colposcopies per life-year gained (strategies 12–14) (**Table 8**); switching to 3-year primary HPV testing at age 25 years (strategies 3 and 9) was still associated with ratios greater than 2,000 colposcopies per life-year gained. The increase in ratios in terms of tests per life-year gained (**Table 9**) and colposcopies per cancer case averted (**Table 10**) was not as pronounced.

When cytology specificity was increased (93.6%), with a decrease in sensitivity (51.4%) (**Appendix Table 16**), the effectiveness of all strategies decreased—especially for screening with cytology alone—but given the corresponding decrease in colposcopies, the ratios using this measure decreased (became more attractive) for all strategies. Efficiency ratios that used colposcopies as a measure of harm decreased by up to 60% for the HPV testing strategies (**Tables 8 and 10**). While switching to 5-year primary HPV testing with 16/18 genotype triage at age 25 years (strategy 6) became an efficient strategy in terms of colposcopies per life-year gained (873 colposcopies per life-year gained) (**Table 8**); the same strategy with cytology triage was associated with a far lower ratio (104 colposcopies per life-year gained, strategy 12). Since screening tests changed only marginally, the ratios using tests per life-year gained were stable (**Table 9**), switching to 5-year primary HPV testing starting at age 25 years at 43 tests per life-year gained; this same strategy at 3-year intervals remained exceedingly high.

We explored the lower-bound (worst-case) relative sensitivity of HPV testing, which impacted both HPV testing alone and cotesting, and found that despite a decrease in the effectiveness of the primary HPV testing strategies, they still provided greater benefits than the current guidelines-based strategies (**Appendix Table 17**). Since the decrease in effectiveness was also accompanied by a decrease in colposcopies, the ratios among efficient strategies improved and more strategies involving 3-year screening with HPV testing alone (strategies 3, 9, and 10) (**Tables 8 and 10**) became efficient, likely to offset the lower sensitivity value.

When we introduced error in the performance of colposcopy/biopsy in classifying a woman's true histologic status, we found that the number of lifetime colposcopies decreased by up to 16% for the HPV and cotesting strategies; this decrease in colposcopies was also accompanied by a decrease in false-positive results, but also an increase in cancer cases and deaths (**Appendix Table 18**). When the effectiveness of precancer treatment (i.e., LEEP) was decreased to 82%, we found only small changes in the number of tests and colposcopies, but 12% to 17% increase in cervical cancer cases and 4% to 7% increase in cervical cancer deaths across all strategies (**Appendix Table 19**). Despite these variations in outcomes, the base-case results of the efficiency analyses remained stable under both sensitivity analyses, with slight decreases in the ratios due to the relatively greater reductions in harms (i.e., colposcopies and tests) than benefits (i.e., life-years and cases averted).

## Followup of HPV-Positive Women

Given that the effectiveness and efficiency of HPV testing depends heavily on the management of screen-positive women, we examined alternative followup algorithms based on protocols from empirical studies (**Tables 8—10** and **Appendix Tables 20—23**). In either primary HPV testing strategy, for women who receive cytology triage (i.e., non-16/18, high-risk positive women in 16/18 triage option; all high-risk positive women in cytology triage option), we explored a more stringent cutoff of LSIL or worse as the threshold to refer women directly to colposcopy (vs. ASC-US or worse in the base case). Overall, all HPV testing strategies had lower effectiveness, as well as decreases in number of tests, colposcopies, and false-positive results (**Appendix Table 20**). Since the change in the measures of harms was slightly larger than the change in measures of benefit, the ratios for all strategies across the three efficiency outcomes marginally decreased (i.e., became more attractive).

We also varied the time to repeat testing for women who receive a normal results upon cytology triage (in both 16/18 genotype and cytology triage options for HPV testing), as well as those who receive HPV-positive and cytology-negative results on cotesting (12 months in the base case). In one scenario, we decreased the time to followup to 6 months, and in another scenario, we increased the time to followup to 24 months. We found that varying the followup interval had a large impact on number of colposcopies and false positives and a smaller impact on effectiveness, especially for HPV testing with cytology triage and cotesting, which send greater proportions of women to repeat testing. When followup was 6 months (**Appendix Table 21**), both colposcopies and effectiveness increased; this result led to similar efficient strategies as in the base-case analysis, but overall higher ratios indicating lower efficiency. Additionally, switching to 3-year cotesting at age 25 years (strategy 15) became the most effective strategy, although with relative high ratios for all three efficiency outcomes. In contrast, when time to followup was extended to 24 months (**Appendix Table 22**), HPV testing and cotesting strategies were less effective, but were more efficient in terms of colposcopies per life-year gained and per cancer case averted. For example, colposcopies per life-year gained associated with switching to 5-year HPV testing at ages 30, 27, and 25 years decreased to 48, 109, and 114 (73, 143, and 195 in the base case; strategies 14, 13, and 12), respectively (**Table 8**).

We evaluated a third alternative triage option in which all HPV-positive women are referred directly to colposcopy. Not surprisingly, the number of colposcopies and false positives was much greater (25% to 29% higher than with 16/18 genotype triage; 40% to 48% higher than with cytology triage), with only a nominal increase in effectiveness (**Appendix Table 23**). For ratios that used colposcopies as a measure of harm, all strategies that referred HPV-positive women to colposcopy without further testing were not efficient, and the base-case strategies and ratios remained the same. Because the number of tests decreased given the removal of repeat testing for HPV-positive women, when defining efficiency in terms of tests per life-year gained, the efficient strategies of 16/18 genotype triage in the base case were replaced by the same strategies of referring all HPV-positive women to colposcopy (i.e., switching to primary HPV testing at age 25 years); as in the base-case analysis, ratios associated with 3-year HPV testing (>29,000 tests per life-year gained) were much higher than with 5-year testing (42 tests per life-year gained).

## HPV-16/18 Vaccinated Women

We evaluated the screening strategies in women assumed to be completely protected from HPV-16/18 infections over the lifetime due to vaccination, to reflect a low-risk population (**Tables 8–10** and **Appendix Table 24**). When protection against HPV-16/18 is complete, the 16/18 genotype and cytology triage options become equivalent, and therefore, our analysis set reduced to cytology alone, HPV primary testing with cytology triage, and cotesting. Women in this low-risk group faced significant reductions in cervical cancer cases and deaths, as well as high reductions in colposcopies and false-positive results. The same strategies were identified as efficient as in the base case; however, the ratios for all efficiency outcomes increased considerably, likely because these strategies (targeted to unvaccinated women in our base-case analysis) remain too intensive in women with considerably lower cervical cancer risk. For example, in terms of colposcopies per life-year gained, the ratio for switching to 5-year HPV testing at age 30 years (strategy 14) increased from 73 to 113; ratios for switching to 5-year HPV testing at younger ages, 27 and 25 years, more than doubled, to 402 and 463, respectively (143 and 195 in the base case; strategies 13 and 12). A similar trend was observed in terms of colposcopies per cancer case averted. When testing was used as a measure of harm, the ratios associated with 5-year HPV testing at age 25 years also doubled; ratios associated with 3-year screening ranged from 159,953 per life-year gained (primary HPV testing, strategy 9) to 429,590 per life-year gained (cotesting, strategy 15).

## Chapter 4. Discussion

This report summarizes the findings from a model-based decision analysis on the long-term health effects, harms, and efficiency of primary HPV testing strategies to inform the updated USPSTF recommendations for cervical cancer screening in the United States. This analysis extends the 2012 decision analysis, which primarily evaluated cytology-based strategies,<sup>41</sup> by focusing specifically on HPV testing for primary screening and including variations in age to switch from cytology-only screening to HPV testing, the rescreening interval, triage options for HPV-positive women, and screening end age. For strategies that overlapped in both reports (e.g., 3-year cytology alone, strategy 1), our results were quite similar to the findings from the previous report.

Consistent with short-term evidence from clinical studies, the model projected that strategies employing primary HPV testing or cotesting were associated with greater health benefits compared to current guidelines-based cytology testing alone but come at a harm of greater testing, colposcopies, and false positives. In all analyses, across three different efficiency measures, primary HPV testing strategies occurring at 5-year intervals were efficient, with the harm-to-benefit ratio decreasing (i.e., becoming more attractive) as the switch age extended from 25 to 30 years. By comparison, the more effective strategies involving 3-year HPV testing generally had exceedingly high ratios. Which HPV testing triage option was efficient depended on the measure of harm used: in terms of colposcopies per life-year gained or per cancer case averted, cytology triage for HPV-positive women was uniformly more efficient than 16/18 genotyping triage; most of the efficient strategies when using colposcopies as a measure of harm consistently involved 5-year HPV testing (switching at ages 25, 27, and 30 years). In contrast, in terms of screening tests per life-year gained, 16/18 genotyping triage was more efficient, mostly involving switching to primary HPV testing at age 25 years (5- and 3-year intervals).

Cotesting strategies were predominantly inefficient and appeared on the efficiency frontier only under two scenarios (repeat followup HPV testing at 6 months and in low-risk, HPV-vaccinated women), but in both of those cases were associated with much higher ratios compared to strategies involving HPV testing alone. When colposcopies were used as the measure of harm, cytology testing alone every 3 years was associated with very low (i.e., attractive) ratios; however, when using total tests as the measure of harm, cytology testing was inefficient across all analyses.

These findings were robust when varying age to end screening from age 65 to 70 and 75 years (assuming no recent abnormal results) with only slight increases in each of the ratios due to decreased efficiency of screening in older ages. When competing all strategies, including end ages for screening, we found that most of the efficient strategies across the three outcomes involved extending the screening age to 70 or 75 years. However, given the vast uncertainties regarding the natural history of HPV infection and screening effectiveness in older women, which were not extensively explored in the current analysis, our findings of screening end age should be interpreted with caution.

When multiple strategies are identified as efficient, selecting the “optimal” strategy depends on a

threshold ratio that would be considered a reasonable balance of harms and benefits. The desired thresholds for each of the three efficiency measures is not clear when using intermediate metrics such as colposcopies or screening tests as a proxy for harm, as it is difficult to compare head-to-head against other (noncervical cancer) health interventions. Although costs were not considered in this analysis, the relative efficiency with respect to costs per life-year gained, a standard metric in traditional cost-effectiveness analysis, can provide another dimension of the tradeoff of harms and benefits that can more easily be benchmarked against other health interventions. In the absence of such a standard metric in this report, we elected to express results using three different metrics of efficiency to help interpret results in the context of different potential tradeoffs that might be considered important by decisionmakers.

## Strengths and Limitations of Modeling

Disease simulation models, when paired with robust data on disease burden and intervention effects, can be powerful tools in projecting long-term outcomes to inform decisionmaking in a timely manner.<sup>42</sup> While most empirical studies on screening effectiveness report findings after only one or two rounds of screening, we can use the model to evaluate the implications of multiple screenings over an extended period and under different combinations of ages to switch, screening intervals, and management algorithms of screen-positive women. As with all model-based analyses, however, this analysis is subject to important limitations.

First, our analysis is based on assumptions of perfect compliance to screening intervals and management of screen-positive women; however, it is well documented that screening practice is not perfect and quite variable across the United States. How loss to followup might differ across testing modalities, age, and interval is uncertain but could impact the overall effectiveness and relative efficiency of the screening strategy. Second, although we examined a number of unique strategies, there may be other strategies that could lead to a more attractive balance of harms and benefits; for example, we restricted our rescreening interval to be no less frequent than every 5 years, but extending intervals to be even longer (e.g., 7 or 10 years) may be more efficient without compromising on effectiveness. Third, as mentioned previously, we did not explore different assumptions regarding the natural history of HPV infection in older women, nor did we examine other strategies or criteria to determine when to stop screening. There is much uncertainty regarding the prevalence and clinical importance of a newly-acquired HPV infection versus reactivation of a previously-acquired infection in older ages, which may impact the optimal age at which to stop screening. Furthermore, recent studies indicate that the incidence and mortality rates from cervical cancer are grossly underestimated by SEER given high rates of hysterectomies in U.S. women, and suggest that the current recommendation for terminating screening may not be optimal.<sup>43,44</sup> The findings from our model, which do correct for hysterectomy rates by age in the population, indicate efficiency and greater effectiveness by extending the screening end age to 70 or 75 years; however, other screening exit criteria and strategies should be further explored in future analyses under various assumptions of disease risk and screening effect at older ages.

Fourth, issues regarding HPV-negative cancers and the implications for the relative effectiveness of HPV testing alone versus cytology alone or cotesting were not fully addressed.<sup>45</sup> The

sensitivity analysis in which HPV relative test sensitivity (compared to cytology) was decreased to a lower bound estimate mimics a scenario of greater missed disease due to HPV negativity; this scenario was the only one in which strategies involving 3-year HPV screening became more efficient with ratios comparable to 5-year HPV screening in the base-case analysis. In our assessment of screening in a low-risk population, we only represented one very specific subset of low-risk women, those who receive protection against HPV-16/18 infection and disease from vaccination. While there are other low-risk segments of the population, this question will become increasingly more pertinent as vaccinated women enter screening age. Finally, it is important to underscore that the results from the model represent average outcome across the whole population and is intended to inform guidelines at the population level, not at an individual level.

## Summary

In summary, the results from the model indicate that primary HPV screening has the potential to increase the effectiveness of screening compared to current U.S. guidelines-based strategies and may represent a reasonable balance of harms and benefits when administered every 5 years. The optimal age at which to switch from cytology to HPV testing and the optimal management of HPV-positive women depend on which outcome (colposcopies or tests) is used as the proxy for harms.

## References


1. Cancer Facts and Figures 2017. <https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-statistics/annual-cancer-facts-and-figures/2017/cancer-facts-and-figures-2017.pdf>. (Last accessed January 31, 2017).
2. U.S. Preventive Services Task Force. Screening for cervical cancer: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2012;156:880-91.
3. Saslow D, Solomon D, Lawson HW, et al. American Cancer Society, American Society for Colposcopy and Cervical Pathology, and American Society for Clinical Pathology screening guidelines for the prevention and early detection of cervical cancer. *CA Cancer J Clin*. 2012;62:147-72.
4. ACOG Practice Bulletin Number 131: screening for cervical cancer. *Obstet Gynecol*. 2012;120:1222-38.
5. Huh WK, Ault KA, Chelmow D, et al. Use of primary high-risk human papillomavirus testing for cervical cancer screening: Interim clinical guidance. *Gynecol Oncol*. 2015; 136(2):178-82.
6. Melnikow J, Henderson JT, Burda BU, et al. Screening for Cervical Cancer With High-Risk Human Papillomavirus Testing: A Systematic Evidence Review for the U.S. Preventive Services Task Force. Evidence Synthesis No. 158. AHRQ Publication No. 17-05231-EF-1. Rockville, MD: Agency for Healthcare Research and Quality; 2017.
7. Campos NG, Burger EA, Sy S, et al. An updated natural history model of cervical cancer: derivation of model parameters. *Am J Epidemiol*. 2014;180:545-55.
8. Kim JJ, Campos NG, Sy S, et al. Inefficiencies and high-value improvements in U.S. cervical cancer screening practice: a cost-effectiveness analysis. *Ann Intern Med*. 2015;163:589-97.
9. National Center for Health Statistics. 2010 National Hospital Discharge Survey (NHDS) Public Use Micro-Data File and Documentation. Available at [https://www.cdc.gov/nchs/nhds/nhds\\_questionnaires.htm](https://www.cdc.gov/nchs/nhds/nhds_questionnaires.htm). (Last accessed January 31, 2017).
10. University of California Berkeley. Berkeley Mortality Database. Available at <http://demog.berkeley.edu/~bmd>. (Data downloaded on August 2, 2016).
11. U.S. National Cancer Institute. Surveillance, Epidemiology, End Results (SEER) Cancer Statistics Review, 1975-2013. Available at [https://seer.cancer.gov/csr/1975\\_2013/](https://seer.cancer.gov/csr/1975_2013/). (Last accessed January 31, 2017).
12. Campos NG, Burger EA, Sy S, et al. An updated natural history model of cervical cancer: derivation of model parameters. *Am J Epidemiol*. 2014;180(5):545-555.
13. Munoz N, Mendez F, Posso H, et al. Incidence, duration, and determinants of cervical human papillomavirus infection in a cohort of Colombian women with normal cytological results. *J Infect Dis*. 2004;190(12):2077-2087.
14. Herrero R, Hildesheim A, Rodriguez AC, et al. Rationale and design of a community-based double-blind randomized clinical trial of an HPV 16 and 18 vaccine in Guanacaste, Costa Rica. *Vaccine*. 2008;26(37):4795-4808.
15. Bratti MC, Rodriguez AC, Schiffman M, et al. Description of a seven-year prospective study of human papillomavirus infection and cervical neoplasia among 10000 women in Guanacaste, Costa Rica. *Rev Panam Salud Publica*. 2004;15(2):75-89.



16. Keefe KA, Schell MJ, Brewer C, et al. A randomized, double blind, Phase III trial using oral beta-carotene supplementation for women with high-grade cervical intraepithelial neoplasia. *Cancer Epidemiol Biomarkers Prev.* 2001;10(10):1029-1035.
17. Meyskens FL, Jr., Surwit E, Moon TE, et al. Enhancement of regression of cervical intraepithelial neoplasia II (moderate dysplasia) with topically applied all-trans-retinoic acid: a randomized trial. *J Natl Cancer Inst.* 1994;86(7):539-543.
18. Castle PE, Schiffman M, Wheeler CM, Solomon D. Evidence for frequent regression of cervical intraepithelial neoplasia-grade 2. *Obstet Gynecol.* 2009;113(1):18-25.
19. Moscicki AB, Ma Y, Wibbelsman C, et al. Rate of and risks for regression of cervical intraepithelial neoplasia 2 in adolescents and young women. *Obstet Gynecol.* 2010;116(6):1373-1380.
20. Wang SM, Colombara D, Shi JF, et al. Six-year regression and progression of cervical lesions of different human papillomavirus viral loads in varied histological diagnoses. *Int J Gynecol Cancer.* 2013;23(4):716-723.
21. McCredie MR, Sharples KJ, Paul C, et al. Natural history of cervical neoplasia and risk of invasive cancer in women with cervical intraepithelial neoplasia 3: a retrospective cohort study. *Lancet Oncol.* 2008;9(5):425-434.
22. McCrory D, Mather D, Bastian L, et al. Evaluation of cervical cytology. Evidence report/technology assessment No. 5. AHCPR publication no. 99-E010. Rockville: Agency for Health Care Policy and Research; 1999.
23. Wheeler CM, Hunt WC, Cuzick J, et al. A population-based study of human papillomavirus genotype prevalence in the United States: baseline measures prior to mass human papillomavirus vaccination. *Int J Cancer.* 2013;132:198-207.
24. Joste NE, Ronnett BM, Hunt WC, et al. Human papillomavirus genotype-specific prevalence across the continuum of cervical neoplasia and cancer. *Cancer Epidemiol Biomarkers Prev.* 2015;24:230-40.
25. Saraiya M, Unger ER, Thompson TD, et al. US assessment of HPV types in cancers: implications for current and 9-valent HPV vaccines. *J Natl Cancer Inst.* 2015;107:djv086.
26. Laskey PW, Meigs JW, Flannery JT. Uterine cervical carcinoma in Connecticut, 1935-1973: evidence for two classes of invasive disease. *J Natl Cancer Inst.* 1976;57:1037-43.
27. International Agency for Research on Cancer. Cancer in Five Continents, Volume 1. Available at <http://ci5.iarc.fr/Default.aspx>. (Last accessed January 31, 2017).
28. Cuzick J, Myers O, Hunt WC, et al. A population-based evaluation of cervical screening in the United States: 2008-2011. *Cancer Epidemiol Biomarkers Prev.* 2014;23:765-73.
29. Cuzick J, Myers O, Hunt WC, et al. Human papillomavirus testing 2007-2012: co-testing and triage utilization and impact on subsequent clinical management. *Int J Cancer.* 2015;136:2854-63.
30. Kinney W, Hunt WC, Dinkelspiel H, et al. Cervical excisional treatment of young women: a population-based study. *Gynecol Oncol.* 2014;132:628-35.
31. Ogilvie GS, Kraiden M, van Niekerk D, et al. HPV for cervical cancer screening (HPV FOCAL): complete round 1 results of a randomized trial comparing HPV-based primary screening to liquid-based cytology for cervical cancer. *Int J Cancer.* 2017;140:440-8.
32. Massad LS, Einstein MH, Huh WK, et al. 2012 updated consensus guidelines for the management of abnormal cervical cancer screening tests and cancer precursors. *J Low Genit Tract Dis.* 2013;17:S1-S27.

33. Koliopoulos G, Arbyn M, Martin-Hirsch P, Kyrgiou M, Prendiville W, Paraskevaidis E. Diagnostic accuracy of human papillomavirus testing in primary cervical screening: a systematic review and meta-analysis of non-randomized studies. *Gynecol Oncol*. 2007;104:232-46.
34. Cox JT, Castle PE, Behrens CM, Sharma A, Wright TC Jr, Cuzick J. Comparison of cervical cancer screening strategies incorporating different combinations of cytology, HPV testing, and genotyping for HPV 16/18: results from the ATHENA HPV study. *Am J Obstet Gynecol*. 2013;208:184 e1- e11.
35. Arbyn M, Ronco G, Anttila A, et al. Evidence regarding human papillomavirus testing in secondary prevention of cervical cancer. *Vaccine*. 2012;30(Suppl 5):F88-99.
36. Ronco G, Giorgi-Rossi P, Carozzi F, et al. Human papillomavirus testing and liquid-based cytology in primary screening of women younger than 35 years: results at recruitment for a randomised controlled trial. *Lancet Oncol*. 2006;7:547-55.
37. Ronco G, Segnan N, Giorgi-Rossi P, et al. Human papillomavirus testing and liquid-based cytology: results at recruitment from the new technologies for cervical cancer randomized controlled trial. *J Natl Cancer Inst*. 2006;98:765-74.
38. Wright TC, Stoler MH, Behrens CM, Sharma A, Zhang G, Wright TL. Primary cervical cancer screening with human papillomavirus: end of study results from the ATHENA study using HPV as the first-line screening test. *Gynecol Oncol*. 2015;136:189-97.
39. Stoler MH, Ronnett BM, Joste NE, et al. The interpretive variability of cervical biopsies and its relationship to HPV status. *Am J Surg Pathol*. 2015;39:729-36.
40. Ghaem-Maghani S, Sagi S, Majeed G, Soutter WP. Incomplete excision of cervical intraepithelial neoplasia and risk of treatment failure: a meta-analysis. *Lancet Oncol*. 2007;8:985-93.
41. Kulasingam SL, Havrilesky LJ, Ghebre R, Myers ER. Screening for cervical cancer: a modeling study for the US Preventive Services Task Force. *J Low Genit Tract Dis*. 2013;17:193-202.
42. Kim JJ, Tosteson AN, Zauber AG, et al. Cancer models and real-world data: better together. *J Natl Cancer Inst*. 2016;108.
43. Beavis AL, Gravitt PE, Rositch AF. Hysterectomy-corrected cervical cancer mortality rates reveal a larger racial disparity in the United States. *Cancer*. 2017 Jan 23.
44. Rositch AF, Nowak RG, Gravitt PE. Increased age and race-specific incidence of cervical cancer after correction for hysterectomy prevalence in the United States from 2000 to 2009. *Cancer*. 2014;120:2032-8.
45. Blatt AJ, Kennedy R, Luff RD, Austin RM, Rabin DS. Comparison of cervical cancer screening results among 256,648 women in multiple clinical practices. *Cancer Cytopathol*. 2015;123:282-8.


**Figure 1. Model Schematic**



The main health states of the model comprise HPV infection (by genotype), precancer (e.g., cervical intraepithelial neoplasia [CIN], grades 2 and 3), and invasive cancer (by stage). The model focuses on squamous cell carcinoma, the most common histologic subtype of cervical cancer.


Screening is used to detect the presence of high-grade precancer, which may resolve spontaneously or can be treated and removed before progressing to cancer, as well as for early detection of invasive cancer. The effectiveness of screening strategies depends on coverage by age, interval, test characteristics, treatment efficacy, and compliance to followup visits.

**Figure 2. Prevalence of HPV by Age and Type**




These graphs show postcalibration model fit to age- and type-specific HPV prevalence from the New Mexico HPV Pap Registry.<sup>23</sup> The model range shows the variation in model fit across the 50 best-fitting parameter sets; the red line shows the mean fit across the 50 sets.

**Figure 3. Type Distribution of HPV in CIN2 and CIN3**




These graphs show postcalibration model fit to HPV type distribution in CIN2 and CIN3 from the New Mexico HPV Pap Registry.<sup>24</sup> The model range shows the variation in model fit across the 50 best-fitting parameter sets; the red line shows the mean fit across the 50 sets.

Figure 4. Type Distribution of HPV in Cancer




This graph shows postcalibration model fit to HPV type distribution in cancer from U.S. population-based cancer registries.<sup>25</sup> The model range shows the variation in model fit across the 50 best-fitting parameter sets; the red line shows the mean fit across the 50 sets.

Figure 5. Cervical Cancer Incidence per 100,000 Women, by Age and Model (Natural History)



This graph shows model-projected cervical cancer incidence rates under a scenario of no intervention (i.e., natural history) compared against cancer registry data from the 1950s and early 1960s, before Pap smear screening was widely available in the United States. Data are from the Connecticut Tumor Registry (CTR) and IARC Cancer in Five Continents (volume 1), which included data from Connecticut, New York, and Hawaii.<sup>26,27</sup> Given the limited data from only a few states, and the potential changes in sexual behavior and other risk factors since the prescreening era, these data were not used directly to calibrate either model but instead were used to assess predictive validity for overall underlying risk. The model range shows the variation in model projections across the 50 best-fitting parameter sets; the red line shows the mean projection across the 50 sets. [Note: Both incidence and mortality rates from the model were calculated using the number of women alive as the denominator, not adjusting for women with hysterectomy, to match the estimates from the cancer registries.]


Figure 6. Cervical Cancer Incidence and Mortality by Age (With Screening)



This graph shows model-projected cervical cancer incidence and mortality rates under assumptions of screening practice patterns reported in the New Mexico HPV Pap Registry,<sup>8,28-30</sup> compared against those reported in SEER cancer registries in recent years (i.e., 2000–2013).<sup>11</sup> The model range shows the variation in model projections under different assumptions of noncompliance to followup diagnostic testing and/or precancer treatments. [Note: Both incidence and mortality rates from the model were calculated using the number of women alive as the denominator, not adjusting for women with hysterectomy, to match the estimates from SEER.]




**Figure 7. Outcomes From HPV-FOCAL Trial (With Screening)**



This graph shows model validation against baseline and 12-month outcomes reported in the HPV-FOCAL trial.<sup>31</sup> We simulated the trial protocol, including three screening scenarios involving switching to primary HPV testing at ages 27, 34, and 52 years. The model range shows the variation in projections across the 50 best-fitting parameter sets.


**Figure 8. Flow Diagram for Alternative Triage Strategies for (hr)HPV-Positive Women**




\* These variables were varied in sensitivity analysis.

Two triage strategies for HPV-positive screening results were examined: a) assuming HPV-16/18 genotype information is available, 16/18-positive women are referred to colposcopy, whereas women positive for other high-risk HPV types receive cytology triage (cytology of ASC-US or worse is referred to colposcopy; cytology-negative receives a followup test in 12 months); b) all women with high-risk HPV receive cytology triage (cytology of ASC-US or worse is referred to colposcopy; cytology-negative receives a followup test in 12 months). A referral threshold of cytology of LSIL or worse was also evaluated, and the interval for followup testing was varied (e.g., 6 or 24 months) in sensitivity analysis.


**Figure 9. Efficiency Frontiers: Colposcopies per Life-Year Gained, Varying HPV Testing Switch Age and Interval (Screening End Age 65 Years), Under Different Scenarios of Triage Options for HPV-Positive Women (Scenarios A–D)**




**Figure 10. Efficiency Frontiers: Tests per Life-Year Gained, Varying HPV Testing Switch Age and Interval (Screening End Age 65 Years), Under Different Scenarios of Triage Options for HPV-Positive Women (Scenarios A–D)**




**Figure 11. Efficiency Frontiers: Colposcopies per Cervical Cancer Case Averted Varying HPV Testing Switch Age and Interval (Screening End Age 65 Years), Under Different Scenarios of Triage Options for HPV-Positive Women (Scenarios A–D)**




**Figure 12. Efficiency Frontiers: Colposcopies per Life-Year Gained When Varying Screening End Age (65, 70, or 75 Years), Under Different Scenarios of Triage Options for HPV-Positive Women (Scenarios A–D)**



**Figure 13. Efficiency Frontiers: Tests per Life-Year Gained When Varying Screening End Age (65, 70, or 75 Years), Under Different Scenarios of Triage Options for HPV-Positive Women (Scenarios A–D)**




**Figure 14. Efficiency Frontiers: Colposcopies per Cervical Cancer Case Averted When Varying Screening End Age (65, 70, or 75 Years), Under Different Scenarios of Triage Options for HPV-Positive Women (Scenarios A–D)**







**Figure 15. Efficiency Frontiers: Colposcopies per Life-Year Gained for All Strategies, Varying HPV Testing Switch Age, Interval, and Screening End Age, Under Different Scenarios of Triage Options for HPV-Positive Women (Scenarios A–D)**



**Figure 16. Efficiency Frontiers: Tests per Life-Year Gained for All Strategies, Varying HPV Testing Switch Age, Interval, and Screening End Age, Under Different Scenarios of Triage Options for HPV-Positive Women (Scenarios A–D)**



**Figure 17. Efficiency Frontiers: Colposcopies per Cervical Cancer Case Averted for All Strategies, Varying HPV Testing Switch Age, Interval, and Screening End Age, Under Different Scenarios of Triage Options for HPV-Positive Women (Scenarios A–D)**



**Table 1. Key Model Attributes**

| <b>Attribute</b>                            | <b>Model</b>  |
|---|---|
| Mode of analysis, simulating life histories | Individual-based  |
| Cycle length                                | Monthly   |
| HPV types included                          | HPV-16; HPV-18; HPV-31; HPV-33; HPV-45; HPV-52; HPV-58; pooled HPV other high-risk; pooled HPV low-risk |
| Natural immunity                            | Reduced probability of future type-specific infection   |
| Health states included                      | Healthy, HPV, CIN2, CIN3, Cancer*   |
| Progression and regression probabilities    | Age-specific, function of HPV persistence   |
| Cancer staging                              | Local, regional, distant  |
| Screening                                   | Yes   |
| Diagnosis                                   | Yes   |
| Precancer treatment                         | Yes   |
| Vaccination                                 | Yes   |

\* The model focuses on squamous cell carcinoma, the most common histologic subtype of cervical cancer.

Abbreviations: CIN: cervical intraepithelial neoplasia; HPV: human papillomavirus.

**Table 2. Cervical Cancer Screening Strategies\***

| #  | Strategy Name                  | Screen (1) test, interval | Screen (1) start age | Screen (2) test, interval | Screen (2) start age | Triage strategies for HPV+ results |
|----|--------------------------------|---------------------------|----------------------|---------------------------|----------------------|------------------------------------|
| 1  | CYTO-3Y, 21                    | Cytology, 3y              | 21                   | --                        | --                   | HPV for ASC-US                     |
| 2  | CYTO-3Y, 21/COTEST-5Y, 30      | Cytology, 3y              | 21                   | Cotest, 5y                | 30                   | Repeat cotest, 12 mos              |
| 3  | CYTO-4Y, 21/HPV-3Y (16/18), 25 | Cytology, 4y              | 21                   | HPV, 3y                   | 25                   | HPV-16/18 genotype                 |
| 4  | CYTO-3Y, 21/HPV-3Y (16/18), 27 | Cytology, 3y              | 21                   | HPV, 3y                   | 27                   | HPV-16/18 genotype                 |
| 5  | CYTO-3Y, 21/HPV-3Y (16/18), 30 | Cytology, 3y              | 21                   | HPV, 3y                   | 30                   | HPV-16/18 genotype                 |
| 6  | CYTO-4Y, 21/HPV-5Y (16/18), 25 | Cytology, 4y              | 21                   | HPV, 5y                   | 25                   | HPV-16/18 genotype                 |
| 7  | CYTO-3Y, 21/HPV-5Y (16/18), 27 | Cytology, 3y              | 21                   | HPV, 5y                   | 27                   | HPV-16/18 genotype                 |
| 8  | CYTO-3Y, 21/HPV-5Y (16/18), 30 | Cytology, 3y              | 21                   | HPV, 5y                   | 30                   | HPV-16/18 genotype                 |
| 9  | CYTO-4Y, 21/HPV-3Y (cyto), 25  | Cytology, 4y              | 21                   | HPV, 3y                   | 25                   | Cytology triage                    |
| 10 | CYTO-3Y, 21/HPV-3Y (cyto), 27  | Cytology, 3y              | 21                   | HPV, 3y                   | 27                   | Cytology triage                    |
| 11 | CYTO-3Y, 21/HPV-3Y (cyto), 30  | Cytology, 3y              | 21                   | HPV, 3y                   | 30                   | Cytology triage                    |
| 12 | CYTO-4Y, 21/HPV-5Y (cyto), 25  | Cytology, 4y              | 21                   | HPV, 5y                   | 25                   | Cytology triage                    |
| 13 | CYTO-3Y, 21/HPV-5Y (cyto), 27  | Cytology, 3y              | 21                   | HPV, 5y                   | 27                   | Cytology triage                    |
| 14 | CYTO-3Y, 21/HPV-5Y (cyto), 30  | Cytology, 3y              | 21                   | HPV, 5y                   | 30                   | Cytology triage                    |
| 15 | CYTO-4Y, 21/COTEST-3Y, 25      | Cytology, 4y              | 21                   | Cotest, 3y                | 25                   | Repeat cotest, 12 mos              |
| 16 | CYTO-3Y, 21/COTEST-3Y, 27      | Cytology, 3y              | 21                   | Cotest, 3y                | 27                   | Repeat cotest, 12 mos              |
| 17 | CYTO-3Y, 21/COTEST-3Y, 30      | Cytology, 3y              | 21                   | Cotest, 3y                | 30                   | Repeat cotest, 12 mos              |
| 18 | CYTO-4Y, 21/COTEST-5Y, 25      | Cytology, 4y              | 21                   | Cotest, 5y                | 25                   | Repeat cotest, 12 mos              |
| 19 | CYTO-3Y, 21/COTEST-5Y, 27      | Cytology, 3y              | 21                   | Cotest, 5y                | 27                   | Repeat cotest, 12 mos              |

\* Management of women with abnormal screening results was assumed to follow clinical guidelines<sup>3,32</sup> and includes: for cytology testing, reflex HPV testing for women with atypical squamous cells of undetermined significance (ASC-US) and referral to colposcopy for women with more severe abnormal results; for cotesting, repeat cotesting in 12 months for women with cytology-negative, HPV-positive results; for HPV testing, two triage options were evaluated: “HPV (16/18)” strategies involved referral to colposcopy for women positive on HPV-16/18 genotype testing and cytology triage for women positive for other (non-16/18) high-risk HPV, and “HPV (cyto)” strategies involved cytology triage for all high-risk HPV-positive women. Strategies were evaluated in context of screening end age of 65, 70, and 75 years in separate analyses.

Abbreviations: Cyto: cytology; HPV: human papillomavirus.

**Table 3. Screening Test Characteristics**

| <b>Test Characteristic*</b>      | <b>Base-Case Value</b> | <b>Source</b> | <b>Worst-Case Value</b> | <b>Best-Case Value</b> | <b>Source</b> |
|----------------------------------|------------------------|---------------|-------------------------|------------------------|---------------|
| Cytology †<br>Sensitivity        | 0.727                  | 33            | 0.514                   | 0.815                  | 33,34         |
| Specificity                      | 0.919                  |               | 0.880                   | 0.936                  |               |
| HPV ‡<br>Relative sensitivity    | 1.24                   | 34            | 1.15                    | 1.37                   | 34-37         |
| Relative specificity             | 0.97                   |               | 0.96                    | 0.98                   |               |
| Cotest ‡<br>Relative sensitivity | 1.31                   | 34            | 1.20                    | 1.42                   | 34-37         |
| Relative specificity             | 0.93                   |               | 0.93                    | 0.94                   |               |

\* Sensitivity (specificity) for all tests defined as probability to detect presence (absence) of CIN2+.

† For cytology testing, positivity threshold is ASC-US.

‡ For HPV testing and cotesting, sensitivity and specificity are relative to cytology test characteristics.

**Table 4. Distribution of Abnormal Cytology Results Conditioned on Histology Result<sup>38</sup>**

| <b>Cytology result</b> | <b>&lt;CIN 2</b> | <b>CIN 2</b> | <b>CIN 3+</b> |
|------------------------|------------------|--------------|---------------|
| ASC-US                 | 0.6674           | 0.3026       | 0.2518        |
| LSIL                   | 0.2994           | 0.5395       | 0.2734        |
| ASC-H                  | 0.0130           | 0.0395       | 0.1007        |
| HSIL                   | 0.0202           | 0.1184       | 0.3741        |

Abbreviations: ASC-H: atypical squamous cells, cannot exclude HSIL; ASC-US: atypical squamous cells of undetermined significance; CIN: cervical intraepithelial neoplasia; HSIL: high-grade squamous intraepithelial lesion; LSIL: low-grade squamous intraepithelial lesion.

**Table 5. Outcomes for Cervical Cancer Screening Strategies Over the Lifetime of Screening (Screening End Age 65 Years)\***

| #  | Strategy                            | Per 1,000 women |           |              |        |                 |                 |                  |          |           |            |
|----|-------------------------------------|-----------------|-----------|--------------|--------|-----------------|-----------------|------------------|----------|-----------|------------|
|    |                                     | Cyto tests      | HPV tests | Total tests† | Colpos | CIN2,3 detected | CIN3+ detected‡ | False positives§ | CC cases | CC deaths | Life-years |
| 0  | No screening                        | 0               | 0         | 0            | 0      | 0               | 0               | 0                | 18.86    | 8.34      | 63921.34   |
| 1  | CYTO-3Y, 21-65                      | 13877           | 786       | 14662        | 645    | 160             | 46              | 484              | 2.34     | 0.76      | 64181.89   |
| 2  | CYTO-3Y, 21 / COTEST-5Y, 30-65      | 11425           | 8380      | 19806        | 1630   | 201             | 54              | 1429             | 1.08     | 0.30      | 64192.97   |
| 3  | CYTO-4Y, 21 / HPV-3Y (16/18), 25-65 | 1905            | 14807     | 16712        | 2530   | 218             | 57              | 2312             | 0.74     | 0.23      | 64195.61   |
| 4  | CYTO-3Y, 21 / HPV-3Y (16/18), 27-65 | 2876            | 13772     | 16648        | 2278   | 214             | 56              | 2063             | 0.83     | 0.25      | 64195.08   |
| 5  | CYTO-3Y, 21 / HPV-3Y (16/18), 30-65 | 3824            | 12428     | 16252        | 1978   | 205             | 54              | 1773             | 1.01     | 0.27      | 64193.51   |
| 6  | CYTO-4Y, 21 / HPV-5Y (16/18), 25-65 | 1706            | 10065     | 11771        | 2068   | 211             | 55              | 1857             | 0.79     | 0.25      | 64195.39   |
| 7  | CYTO-3Y, 21 / HPV-5Y (16/18), 27-65 | 2697            | 9290      | 11987        | 1861   | 207             | 55              | 1655             | 0.89     | 0.28      | 64194.69   |
| 8  | CYTO-3Y, 21 / HPV-5Y (16/18), 30-65 | 3675            | 8476      | 12151        | 1635   | 199             | 53              | 1435             | 1.05     | 0.29      | 64193.38   |
| 9  | CYTO-4Y, 21 / HPV-3Y (cyto), 25-65  | 2277            | 14790     | 17067        | 2209   | 217             | 56              | 1992             | 0.75     | 0.23      | 64195.53   |
| 10 | CYTO-3Y, 21 / HPV-3Y (cyto), 27-65  | 3205            | 13738     | 16943        | 1992   | 213             | 56              | 1779             | 0.85     | 0.25      | 64194.82   |
| 11 | CYTO-3Y, 21 / HPV-3Y (cyto), 30-65  | 4102            | 12397     | 16499        | 1734   | 203             | 54              | 1530             | 1.04     | 0.28      | 64193.19   |
| 12 | CYTO-4Y, 21 / HPV-5Y (cyto), 25-65  | 1993            | 10049     | 12042        | 1826   | 209             | 55              | 1617             | 0.81     | 0.25      | 64195.35   |
| 13 | CYTO-3Y, 21 / HPV-5Y (cyto), 27-65  | 2950            | 9273      | 12223        | 1648   | 205             | 54              | 1443             | 0.91     | 0.28      | 64194.44   |
| 14 | CYTO-3Y, 21 / HPV-5Y (cyto), 30-65  | 3888            | 8459      | 12348        | 1452   | 198             | 53              | 1254             | 1.08     | 0.29      | 64193.07   |
| 15 | CYTO-4Y, 21 / COTEST-3Y, 25-65      | 15723           | 14693     | 30416        | 2535   | 223             | 57              | 2312             | 0.76     | 0.23      | 64195.50   |
| 16 | CYTO-3Y, 21 / COTEST-3Y, 27-65      | 15765           | 13723     | 29488        | 2303   | 218             | 57              | 2084             | 0.83     | 0.25      | 64194.75   |
| 17 | CYTO-3Y, 21 / COTEST-3Y, 30-65      | 15456           | 12411     | 27867        | 2021   | 209             | 55              | 1812             | 1.03     | 0.27      | 64193.17   |
| 18 | CYTO-4Y, 21 / COTEST-5Y, 25-65      | 10944           | 9914      | 20859        | 2029   | 213             | 55              | 1816             | 0.82     | 0.26      | 64195.26   |
| 19 | CYTO-3Y, 21 / COTEST-5Y, 27-65      | 11275           | 9233      | 20508        | 1846   | 209             | 55              | 1637             | 0.89     | 0.27      | 64194.40   |

\* Outcomes calculated from age 20 to 100 years.

† Total number of tests, irrespective of primary, triage, or surveillance context.

‡ CIN3+ includes CIN3s and cervical cancers detected through screening (excludes clinically detected cancers).

§ Total number of colposcopies that did not result in CIN2, CIN3, or cancer detection.

Abbreviations: CC: cervical cancer; CIN: cervical intraepithelial neoplasia; Colpos: colposcopies; Cyto: cytology; HPV: Human papillomavirus.



**Table 6. Outcomes for Cervical Cancer Screening Strategies Over the Lifetime of Screening (Screening End Age 70 Years)\***

| #  | Strategy                              | Per 1,000 women |           |              |        |                 |                 |                  |          |           |            |
|----|---------------------------------------|-----------------|-----------|--------------|--------|-----------------|-----------------|------------------|----------|-----------|------------|
|    |                                       | Cyto tests      | HPV tests | Total tests† | Colpos | CIN2,3 detected | CIN3+ detected‡ | False positives§ | CC cases | CC deaths | Life-years |
| 0  | No screening                          | 0               | 0         | 0            | 0      | 0               | 0               | 0                | 18.86    | 8.34      | 63921.34   |
| 1  | CYTO-3Y, 21-70                        | 15149           | 855       | 16004        | 689    | 166             | 48              | 522              | 1.98     | 0.55      | 64184.58   |
| 2  | CYTO-3Y, 21 / COTEST-5Y, 30-70        | 12173           | 9128      | 21301        | 1705   | 207             | 55              | 1498             | 0.88     | 0.19      | 64194.07   |
| 3  | CYTO-4Y, 21 / HPV-3Y (16/18), 25-70   | 1940            | 16024     | 17965        | 2620   | 224             | 58              | 2395             | 0.56     | 0.13      | 64196.60   |
| 4  | CYTO-3Y, 21 / HPV-3Y (16/18), 27-70   | 2914            | 15055     | 17968        | 2373   | 220             | 57              | 2153             | 0.64     | 0.14      | 64196.14   |
| 5  | CYTO-3Y, 21 / HPV-3Y (16/18), 30-70   | 3861            | 13721     | 17582        | 2074   | 210             | 56              | 1863             | 0.82     | 0.17      | 64194.60   |
| 6  | CYTO-4Y, 21 / HPV-5Y (16/18), 25-70   | 1729            | 10807     | 12536        | 2134   | 216             | 56              | 1917             | 0.60     | 0.15      | 64196.47   |
| 7  | CYTO-3Y, 21 / HPV-5Y (16/18), 27-70   | 2722            | 10049     | 12771        | 1930   | 212             | 56              | 1718             | 0.68     | 0.17      | 64195.84   |
| 8  | CYTO-3Y, 21 / HPV-5Y (16/18), 30-70   | 3698            | 9216      | 12914        | 1700   | 204             | 55              | 1495             | 0.85     | 0.18      | 64194.50   |
| 9  | CYTO-4Y, 21 / HPV-3Y (cyto), 25-65/70 | 2327            | 15979     | 18306        | 2283   | 222             | 58              | 2060             | 0.58     | 0.13      | 64196.52   |
| 10 | CYTO-3Y, 21 / HPV-3Y (cyto), 27-70    | 3260            | 15035     | 18295        | 2073   | 218             | 57              | 1854             | 0.66     | 0.14      | 64195.91   |
| 11 | CYTO-3Y, 21 / HPV-3Y (cyto), 30-70    | 4157            | 13704     | 17861        | 1816   | 209             | 56              | 1606             | 0.85     | 0.17      | 64194.31   |
| 12 | CYTO-4Y, 21 / HPV-5Y (cyto), 25-70    | 2028            | 10790     | 12818        | 1881   | 214             | 56              | 1667             | 0.62     | 0.15      | 64196.41   |
| 13 | CYTO-3Y, 21 / HPV-5Y (cyto), 27-70    | 2986            | 10032     | 13017        | 1706   | 210             | 56              | 1496             | 0.71     | 0.17      | 64195.57   |
| 14 | CYTO-3Y, 21 / HPV-5Y (cyto), 30-70    | 3923            | 9198      | 13121        | 1508   | 203             | 55              | 1304             | 0.88     | 0.18      | 64194.18   |
| 15 | CYTO-4Y, 21 / COTEST-3Y, 25-70        | 16929           | 15899     | 32829        | 2640   | 229             | 58              | 2411             | 0.58     | 0.14      | 64196.49   |
| 16 | CYTO-3Y, 21 / COTEST-3Y, 27-70        | 17021           | 14979     | 32000        | 2412   | 224             | 58              | 2188             | 0.65     | 0.15      | 64195.78   |
| 17 | CYTO-3Y, 21 / COTEST-3Y, 30-70        | 16722           | 13677     | 30399        | 2131   | 215             | 56              | 1916             | 0.84     | 0.17      | 64194.21   |
| 18 | CYTO-4Y, 21 / COTEST-5Y, 25-70        | 11696           | 10666     | 22361        | 2105   | 219             | 57              | 1886             | 0.63     | 0.15      | 64196.31   |
| 19 | CYTO-3Y, 21 / COTEST-5Y, 27-70        | 12038           | 9995      | 22033        | 1922   | 215             | 56              | 1707             | 0.69     | 0.17      | 64195.50   |

\* Outcomes calculated from age 20 to 100 years.

† Total number of tests, irrespective of primary, triage, or surveillance context.

‡ CIN3+ includes CIN3 and cervical cancer detected through screening (excludes clinically-detected cancer).

§ Total number of colposcopies that did not result in CIN2, CIN3, or cancer detection.

Abbreviations: CC: cervical cancer; CIN: cervical intraepithelial neoplasia; Colpos: colposcopies; Cyto: cytology; HPV: Human papillomavirus.

**Table 7. Outcomes for Cervical Cancer Screening Strategies Over the Lifetime of Screening (Screening End Age 75 Years)\***

| #  | Strategy                            | Per 1,000 women |           |              |        |                 |                 |                  |          |           |            |
|----|-------------------------------------|-----------------|-----------|--------------|--------|-----------------|-----------------|------------------|----------|-----------|------------|
|    |                                     | Cyto tests      | HPV tests | Total tests† | Colpos | CIN2,3 detected | CIN3+ detected‡ | False positives§ | CC cases | CC deaths | Life-years |
| 0  | No screening                        | 0               | 0         | 0            | 0      | 0               | 0               | 0                | 18.86    | 8.34      | 63921.34   |
| 1  | CYTO-3Y, 21-75                      | 16213           | 913       | 17127        | 724    | 170             | 49              | 554              | 1.76     | 0.42      | 64185.93   |
| 2  | CYTO-3Y, 21 / COTEST-5Y, 30-75      | 12848           | 9803      | 22651        | 1767   | 211             | 56              | 1556             | 0.77     | 0.13      | 64194.54   |
| 3  | CYTO-4Y, 21 / HPV-3Y (16/18), 25-75 | 1967            | 17051     | 19018        | 2687   | 228             | 59              | 2459             | 0.47     | 0.09      | 64197.01   |
| 4  | CYTO-3Y, 21 / HPV-3Y (16/18), 27-75 | 2942            | 16156     | 19098        | 2446   | 224             | 58              | 2222             | 0.54     | 0.09      | 64196.58   |
| 5  | CYTO-3Y, 21 / HPV-3Y (16/18), 30-75 | 3890            | 14825     | 18715        | 2147   | 215             | 57              | 1931             | 0.72     | 0.12      | 64195.03   |
| 6  | CYTO-4Y, 21 / HPV-5Y (16/18), 25-75 | 1748            | 11473     | 13222        | 2186   | 220             | 58              | 1966             | 0.49     | 0.09      | 64196.93   |
| 7  | CYTO-3Y, 21 / HPV-5Y (16/18), 27-75 | 2741            | 10730     | 13472        | 1984   | 216             | 57              | 1768             | 0.58     | 0.11      | 64196.36   |
| 8  | CYTO-3Y, 21 / HPV-5Y (16/18), 30-75 | 3717            | 9880      | 13597        | 1752   | 208             | 56              | 1544             | 0.75     | 0.13      | 64194.93   |
| 9  | CYTO-4Y, 21 / HPV-3Y (cyto), 25-75  | 2366            | 17013     | 19380        | 2340   | 226             | 59              | 2114             | 0.49     | 0.09      | 64196.93   |
| 10 | CYTO-3Y, 21 / HPV-3Y (cyto), 27-75  | 3301            | 16112     | 19413        | 2132   | 223             | 58              | 1910             | 0.56     | 0.10      | 64196.32   |
| 11 | CYTO-3Y, 21 / HPV-3Y (cyto), 30-75  | 4198            | 14784     | 18983        | 1876   | 213             | 57              | 1662             | 0.75     | 0.12      | 64194.71   |
| 12 | CYTO-4Y, 21 / HPV-5Y (cyto), 25-75  | 2057            | 11455     | 13512        | 1926   | 218             | 57              | 1708             | 0.51     | 0.09      | 64196.89   |
| 13 | CYTO-3Y, 21 / HPV-5Y (cyto), 27-75  | 3015            | 10712     | 13728        | 1752   | 215             | 57              | 1537             | 0.59     | 0.11      | 64196.12   |
| 14 | CYTO-3Y, 21 / HPV-5Y (cyto), 30-75  | 3952            | 9861      | 13813        | 1552   | 207             | 56              | 1345             | 0.78     | 0.13      | 64194.63   |
| 15 | CYTO-4Y, 21 / COTEST-3Y, 25-75      | 18009           | 16979     | 34988        | 2726   | 233             | 60              | 2493             | 0.49     | 0.09      | 64196.92   |
| 16 | CYTO-3Y, 21 / COTEST-3Y, 27-75      | 18113           | 16071     | 34184        | 2500   | 229             | 59              | 2271             | 0.55     | 0.10      | 64196.20   |
| 17 | CYTO-3Y, 21 / COTEST-3Y, 30-75      | 17813           | 14768     | 32581        | 2218   | 219             | 58              | 1999             | 0.75     | 0.12      | 64194.61   |
| 18 | CYTO-4Y, 21 / COTEST-5Y, 25-75      | 12372           | 11342     | 23714        | 2166   | 223             | 58              | 1943             | 0.52     | 0.10      | 64196.81   |
| 19 | CYTO-3Y, 21 / COTEST-5Y, 27-75      | 12724           | 10681     | 23405        | 1985   | 219             | 57              | 1766             | 0.58     | 0.11      | 64196.01   |

\* Outcomes calculated from age 20 to 100 years.

† Total number of tests, irrespective of primary, triage, or surveillance context.

‡ CIN3+ includes CIN3 and cervical cancer detected through screening (excludes clinically-detected cancer).

§ Total number of colposcopies that did not result in CIN2, CIN3, or cancer detection.

Abbreviations: CC: cervical cancer; CIN: cervical intraepithelial neoplasia; Colpos: colposcopies; Cyto: cytology; HPV: Human papillomavirus.

**Table 8. Sensitivity Analysis Summary: Impact of Uncertainty on Colposcopies per Life-Year Gained**

| #  | Strategy*                           | Base case | Screen end age 70 years | Screen end age 75 years | Cyto sens 81.5%, spec 88.0% | Cyto sens 51.4%, spec 93.6% | HPV relative sens 1.15 | Imperfect colpo | CIN treat efficacy 82% | LSIL threshold to colpo† | 6-month follow up‡ | 24-month follow up‡ | Direct colpo referral§ | 16/18 vacc |
|----|-------------------------------------|-----------|-------------------------|-------------------------|-----------------------------|-----------------------------|------------------------|-----------------|------------------------|--------------------------|--------------------|---------------------|------------------------|------------|
| 0  | No Screening¶                       | --        | --                      | --                      | --                          | --                          | --                     | --              | --                     | --                       | --                 | --                  | --                     | --         |
| 1  | CYTO-3Y, 21-65                      | 3         | 3                       | 3                       | 3                           | 2                           | 2                      | 3               | 3                      | 2                        | 2                  | 2                   | 3                      | 5          |
| 14 | CYTO-3Y, 21 / HPV-5Y (cyto), 30-65  | 73        | 85                      | 95                      | 80                          | 38                          | 66                     | 42              | 69                     | 71                       | 90                 | 48                  | 73                     | 113        |
| 2  | CYTO-3Y, 21 / COTEST-5Y, 30-65      | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 8  | CYTO-3Y, 21 / HPV-5Y (16/18), 30-65 | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 13 | CYTO-3Y, 21 / HPV-5Y (cyto), 27-65  | 143       | 143                     | 135                     | 167                         | 67                          | 116                    | 89              | 115                    | 128                      | 222                | 109                 | 143                    | 402        |
| 11 | CYTO-3Y, 21 / HPV-3Y (cyto), 30-65  | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 12 | CYTO-4Y, 21 / HPV-5Y (cyto), 25-65  | 195       | 208                     | 225                     | 323                         | 104                         | x                      | 136             | 212                    | 184                      | 255                | 114                 | 195                    | 463        |
| 19 | CYTO-3Y, 21 / COTEST-5Y, 27-65      | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 7  | CYTO-3Y, 21 / HPV-5Y (16/18), 27-65 | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 5  | CYTO-3Y, 21 / HPV-3Y (16/18), 30-65 | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 10 | CYTO-3Y, 21 / HPV-3Y (cyto), 27-65  | x         | x                       | x                       | x                           | x                           | 127                    | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 17 | CYTO-3Y, 21 / COTEST-3Y, 30-65      | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 18 | CYTO-4Y, 21 / COTEST-5Y, 25-65      | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 6  | CYTO-4Y, 21 / HPV-5Y (16/18), 25-65 | x         | x                       | x                       | x                           | 873                         | x                      | 1,251           | x                      | x                        | x                  | x                   | x                      | x          |
| 9  | CYTO-4Y, 21 / HPV-3Y (cyto), 25-65  | 2,188     | 3,758                   | x                       | 2,590                       | x                           | 234                    | 1,741           | 2,016                  | 1,789                    | x                  | 1,369               | 2,188                  | 8,163      |
| 4  | CYTO-3Y, 21 / HPV-3Y (16/18), 27-65 | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 16 | CYTO-3Y, 21 / COTEST-3Y, 27-65      | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 3  | CYTO-4Y, 21 / HPV-3Y (16/18), 25-65 | 3,822     | 4,014                   | 6,239                   | 3,347                       | 2,147                       | 459                    | 2,064           | 4,661                  | 2,783                    | x                  | 2,074               | 3,822                  | x          |
| 15 | CYTO-4Y, 21 / COTEST-3Y, 25-65      | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | 3,887              | x                   | x                      | 10,884     |
| 25 | CYTO-3Y, 21 / HPV-5Y (colpo), 30-65 | na        | na                      | na                      | na                          | na                          | na                     | na              | na                     | na                       | na                 | na                  | x                      | na         |
| 24 | CYTO-3Y, 21 / HPV-5Y (colpo), 27-65 | na        | na                      | na                      | na                          | na                          | na                     | na              | na                     | na                       | na                 | na                  | x                      | na         |
| 22 | CYTO-3Y, 21 / HPV-3Y (colpo), 30-65 | na        | na                      | na                      | na                          | na                          | na                     | na              | na                     | na                       | na                 | na                  | x                      | na         |
| 23 | CYTO-4Y, 21 / HPV-5Y (colpo), 25-65 | na        | na                      | na                      | na                          | na                          | na                     | na              | na                     | na                       | na                 | na                  | x                      | na         |
| 21 | CYTO-3Y, 21 / HPV-3Y (colpo), 27-65 | na        | na                      | na                      | na                          | na                          | na                     | na              | na                     | na                       | na                 | na                  | x                      | na         |
| 20 | CYTO-4Y, 21 / HPV-3Y (colpo), 25-65 | na        | na                      | na                      | na                          | na                          | na                     | na              | na                     | na                       | na                 | na                  | x                      | na         |

\* Strategies are listed in order of increasing colposcopies in the base-case analysis. "HPV (colpo)" strategies involved direct colposcopy referral for all HPV-positive women, which were only included in one sensitivity analysis.

† Among HPV-positive women who received cytology triage, LSIL was used as a threshold for colposcopy referral (base-case analysis assumed ASC-US); see **Figure 8**.

‡ Among HPV-positive women who received cytology triage and were referred for follow up testing, interval was varied 6 to 24 months (base-case analysis assumed 12 months); see **Figure 8**.

§ All strategies involving direct colposcopy referral for all HPV-positive women were inefficient.

|| Screening was evaluated in women fully vaccinated with HPV-16/18 vaccines in preadolescence.

¶ No screening was the baseline comparator strategy in all analyses.

Abbreviations: Colpo: colposcopy; Cyto: cytology; HPV: Human papillomavirus; LSIL: low-grade squamous intraepithelial lesion; na: not analyzed; sens: sensitivity; spec: specificity; vacc: vaccinated. "x" marks strategies that were inefficient (i.e., more harmful and less beneficial than another strategy, or had a higher harm-to-benefit ratio than a strategy with greater harms).

**Table 9. Sensitivity Analysis Summary: Impact of Uncertainty on Tests per Life-Year Gained**

| #  | Strategy*                           | Base case | Screen end age 70 years | Screen end age 75 years | Cyto sens 81.5%, spec 88.0% | Cyto sens 51.4%, spec 93.6% | HPV relative sens 1.15 | Imperfect colpo | CIN treat efficacy 82% | LSIL threshold to colpo† | 6-month follow up‡ | 24-month follow up‡ | Direct colpo referral§ | 16/18 vacc |
|----|-------------------------------------|-----------|-------------------------|-------------------------|-----------------------------|-----------------------------|------------------------|-----------------|------------------------|--------------------------|--------------------|---------------------|------------------------|------------|
| 0  | No Screening¶                       | --        | --                      | --                      | --                          | --                          | --                     | --              | --                     | --                       | --                 | --                  | --                     | --         |
| 6  | CYTO-4Y, 21 / HPV-5Y (16/18), 25-65 | 43        | 46                      | 48                      | 43                          | 43                          | 42                     | 42              | 43                     | 43                       | 44                 | 41                  | x                      | x          |
| 7  | CYTO-3Y, 21 / HPV-5Y (16/18), 27-65 | x         | x                       | X                       | x                           | x                           | x                      | x               | x                      | x                        |                    | x                   | x                      | x          |
| 12 | CYTO-4Y, 21 / HPV-5Y (cyto), 25-65  | x         | x                       | X                       | x                           | x                           | x                      | x               | x                      | x                        | 6,359              | x                   | x                      | 107        |
| 8  | CYTO-3Y, 21 / HPV-5Y (16/18), 30-65 | x         | x                       | X                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 13 | CYTO-3Y, 21 / HPV-5Y (cyto), 27-65  | x         | x                       | X                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 14 | CYTO-3Y, 21 / HPV-5Y (cyto), 30-65  | x         | x                       | X                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 1  | CYTO-3Y, 21 -65                     | x         | x                       | X                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 5  | CYTO-3Y, 21 / HPV-3Y (16/18), 30-65 | x         | x                       | X                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 11 | CYTO-3Y, 21 / HPV-3Y (cyto), 30-65  | x         | x                       | X                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 4  | CYTO-3Y, 21 / HPV-3Y (16/18), 27-65 | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 3  | CYTO-4Y, 21 / HPV-3Y (16/18), 25-65 | 22,335    | 41,852                  | 69,063                  | 23,126                      | 23,131                      | 3,111                  | 24,006          | 24,871                 | 21,594                   | 45,833             | 25,769              | 35,241                 | x          |
| 10 | CYTO-3Y, 21 / HPV-3Y (cyto), 27-65  | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 9  | CYTO-4Y, 21 / HPV-3Y (cyto), 25-65  | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | 159,953    |
| 2  | CYTO-3Y, 21 / COTEST-5Y, 30-65      | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 19 | CYTO-3Y, 21 / COTEST-5Y, 27-65      | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 18 | CYTO-4Y, 21 / COTEST-5Y, 25-65      | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 17 | CYTO-3Y, 21 / COTEST-3Y, 30-65      | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 16 | CYTO-3Y, 21 / COTEST-3Y, 27-65      | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 15 | CYTO-4Y, 21 / COTEST-3Y, 25-65      | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | 134,110            | x                   | x                      | 429,590    |
| 23 | CYTO-4Y, 21 / HPV-5Y (colpo), 25-65 | na        | na                      | na                      | na                          | na                          | na                     | na              | na                     | na                       | na                 | na                  | 42                     | na         |
| 24 | CYTO-3Y, 21 / HPV-5Y (colpo), 27-65 | na        | na                      | na                      | na                          | na                          | na                     | na              | na                     | na                       | na                 | na                  | x                      | na         |
| 25 | CYTO-3Y, 21 / HPV-5Y (colpo), 30-65 | na        | na                      | na                      | na                          | na                          | na                     | na              | na                     | na                       | na                 | na                  | x                      | na         |
| 22 | CYTO-3Y, 21 / HPV-3Y (colpo), 30-65 | na        | na                      | na                      | na                          | na                          | na                     | na              | na                     | na                       | na                 | na                  | x                      | na         |
| 20 | CYTO-4Y, 21 / HPV-3Y (colpo), 25-65 | na        | na                      | na                      | na                          | na                          | na                     | na              | na                     | na                       | na                 | na                  | 29,184                 | na         |
| 21 | CYTO-3Y, 21 / HPV-3Y (colpo), 27-65 | na        | na                      | na                      | na                          | na                          | na                     | na              | na                     | na                       | na                 | na                  | x                      | na         |

\* Strategies are listed in order of increasing tests in the base-case analysis. "HPV (colpo)" strategies involved direct colposcopy referral for all HPV-positive women, which were only included in one sensitivity analysis.

† Among HPV-positive women who received cytology triage, LSIL was used as a threshold for colposcopy referral (base-case analysis assumed ASC-US); see **Figure 8**.

‡ Among HPV-positive women who received cytology triage and were referred for follow up testing, interval was varied 6 to 24 months (base-case analysis assumed 12 months); see **Figure 8**.

§ Rank order of efficient strategies in scenario including colposcopy referral for all HPV-positive women differed from base-case analysis.

|| Screening was evaluated in women fully vaccinated with HPV-16/18 vaccines in preadolescence.

¶ No screening was the baseline comparator strategy in all analyses.

Abbreviations: Colpo: colposcopy; Cyto: cytology; HPV: Human papillomavirus; LSIL: low-grade squamous intraepithelial lesion; na: not analyzed; sens: sensitivity; spec: specificity; vacc: vaccinated. "x" marks strategies that were inefficient (i.e., more harmful and less beneficial than another strategy, or had a higher harm-to-benefit ratio than a strategy with greater harms).

**Table 10. Sensitivity Analysis Summary: Impact of Uncertainty on Colposcopies per Cervical Cancer Case Averted**

| #  | Strategy*                         | Base case | Screen end age 70 years | Screen end age 75 years | Cyto sens 81.5%, spec 88.0% | Cyto sens 51.4%, spec 93.6% | HPV relative sens 1.15 | Imperfect colpo | CIN treat efficacy 82% | LSIL threshold to colpo† | 6-month follow up‡ | 24-month follow up‡ | Direct colpo referral§ | 16/18 vacc |
|----|-----------------------------------|-----------|-------------------------|-------------------------|-----------------------------|-----------------------------|------------------------|-----------------|------------------------|--------------------------|--------------------|---------------------|------------------------|------------|
| 0  | No Screening¶                     | --        | --                      | --                      | --                          | --                          | --                     | --              | --                     | --                       | --                 | --                  | --                     | --         |
| 1  | CYTO-3Y, 21-65                    | 39        | 41                      | 42                      | 54                          | 33                          | 38                     | 42              | 41                     | 39                       | 39                 | 39                  | 39                     | 77         |
| 14 | CYTO-3Y, 21/HPV-5Y (cyto), 30-65  | 640       | 748                     | 847                     | 638                         | 399                         | 551                    | 375             | 601                    | 624                      | 792                | 423                 | 640                    | 900        |
| 2  | CYTO-3Y, 21/COTEST-5Y, 30-65      | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 8  | CYTO-3Y, 21/HPV-5Y (16/18), 30-65 | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 13 | CYTO-3Y, 21/HPV-5Y (cyto), 27-65  | 1,161     | 1,134                   | 1,081                   | 1,495                       | 647                         | 1,138                  | 681             | 1,028                  | 1,118                    | 1,601              | 879                 | 1,161                  | 2,900      |
| 11 | CYTO-3Y, 21/HPV-3Y (cyto), 30-65  | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 12 | CYTO-4Y, 21/HPV-5Y (cyto), 25-65  | 1,735     | 1,914                   | 2,064                   | 2,017                       | 974                         | x                      | 1,101           | 1,543                  | 1,682                    | 2,023              | 1,449               | 1,735                  | 3,247      |
| 19 | CYTO-3Y, 21/COTEST-5Y, 27-65      | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 7  | CYTO-3Y, 21/HPV-5Y (16/18), 27-65 | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 5  | CYTO-3Y, 21/HPV-3Y (16/18), 30-65 | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 10 | CYTO-3Y, 21/HPV-3Y (cyto), 27-65  | x         | x                       | x                       | x                           | x                           | 1,307                  | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 17 | CYTO-3Y, 21/COTEST-3Y, 30-65      | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 18 | CYTO-4Y, 21/COTEST-5Y, 25-65      | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 6  | CYTO-4Y, 21/HPV-5Y (16/18), 25-65 | x         | x                       | 15,899                  | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 9  | CYTO-4Y, 21/HPV-3Y (cyto), 25-65  | 7,018     | 9,848                   | 19,645                  | 7,240                       | 6,223                       | 1,757                  | 6,261           | 6,879                  | 6,132                    | 17,353             | 4,839               | 7,018                  | 19,066     |
| 4  | CYTO-3Y, 21/HPV-3Y (16/18), 27-65 | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 16 | CYTO-3Y, 21/COTEST-3Y, 27-65      | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | x                  | x                   | x                      | x          |
| 3  | CYTO-4Y, 21/HPV-3Y (16/18), 25-65 | 23,974    | 23,361                  | 25,112                  | 37,553                      | 11,659                      | 4,500                  | 16,287          | 15,224                 | 12,647                   | x                  | 7,646               | 23,974                 | x          |
| 15 | CYTO-4Y, 21/COTEST-3Y, 25-65      | x         | x                       | x                       | x                           | x                           | x                      | x               | x                      | x                        | 27,217             | x                   | x                      | 418,870    |
| 23 | CYTO-4Y, 21/HPV-5Y (colpo), 25-65 | na        | na                      | na                      | na                          | na                          | na                     | na              | na                     | na                       | na                 | na                  | x                      | na         |
| 24 | CYTO-3Y, 21/HPV-5Y (colpo), 27-65 | na        | na                      | na                      | na                          | na                          | na                     | na              | na                     | na                       | na                 | na                  | x                      | na         |
| 25 | CYTO-3Y, 21/HPV-5Y (colpo), 30-65 | na        | na                      | na                      | na                          | na                          | na                     | na              | na                     | na                       | na                 | na                  | x                      | na         |
| 22 | CYTO-3Y, 21/HPV-3Y (colpo), 30-65 | na        | na                      | na                      | na                          | na                          | na                     | na              | na                     | na                       | na                 | na                  | x                      | na         |
| 20 | CYTO-4Y, 21/HPV-3Y (colpo), 25-65 | na        | na                      | na                      | na                          | na                          | na                     | na              | na                     | na                       | na                 | na                  | x                      | na         |
| 21 | CYTO-3Y, 21/HPV-3Y (colpo), 27-65 | na        | na                      | na                      | na                          | na                          | na                     | na              | na                     | na                       | na                 | na                  | x                      | na         |

\* Strategies are listed in order of increasing colposcopies in the base-case analysis. "HPV (colpo)" strategies involved direct colposcopy referral for all HPV-positive women, which were only included in one sensitivity analysis.

† Among HPV-positive women who received cytology triage, LSIL was used as a threshold for colposcopy referral (base-case analysis assumed ASC-US); see **Figure 8**.

‡ Among HPV-positive women who received cytology triage and were referred for follow up testing, interval was varied 6 to 24 months (base-case analysis assumed 12 months); see **Figure 8**.

§ All strategies involving direct colposcopy referral for HPV-positive women were inefficient.

|| Screening was evaluated in women fully vaccinated with HPV-16/18 vaccines in preadolescence.

¶ No screening was the baseline comparator strategy in all analyses.

Abbreviations: Colpo: colposcopy; Cyto: cytology; HPV: Human papillomavirus; LSIL: low-grade squamous intraepithelial lesion; na: not analyzed; sens: sensitivity; spec: specificity; vacc: vaccinated. "x" marks strategies that were inefficient (i.e., more harmful and less beneficial than another strategy, or had a higher harm-to-benefit ratio than a strategy with greater harms).

**Appendix Table 1. Natural History Model Parameters, Post-Calibration**

| <b>Model parameter</b>                  | <b>Range of values for 50 best-fitting sets*</b> |
|---|--|
| <b>HPV-16 incidence<sup>12,13</sup></b> |  |
| Age <20 years                           | 0.00003-0.01018                                  |
| Age 20-24 years                         | 0.00170-0.00842                                  |
| Age 25-29 years                         | 0.00147-0.00416                                  |
| Age 30-34 years                         | 0.00136-0.00373                                  |
| Age 35-39 years                         | 0.00132-0.00344                                  |
| Age 40-44 years                         | 0.00124-0.00328                                  |
| Age 45-49 years                         | 0.00113-0.00312                                  |
| Age 50-54 years                         | 0.00102-0.00282                                  |
| Age 55-59 years                         | 0.00093-0.00255                                  |
| Age 60+ years                           | 0.00041-0.00231                                  |
| <b>HPV-18 incidence<sup>12,13</sup></b> |  |
| Age <20 years                           | 0.00001-0.00328                                  |
| Age 20-24 years                         | 0.00101-0.00348                                  |
| Age 25-29 years                         | 0.00074-0.00297                                  |
| Age 30-34 years                         | 0.00062-0.00172                                  |
| Age 35-39 years                         | 0.00048-0.00142                                  |
| Age 40-44 years                         | 0.00041-0.00113                                  |
| Age 45-49 years                         | 0.00037-0.00097                                  |
| Age 50-54 years                         | 0.00034-0.00088                                  |
| Age 55-59 years                         | 0.00030-0.00079                                  |
| Age 60+ years                           | 0.00013-0.00072                                  |
| <b>HPV-31 incidence<sup>12,13</sup></b> |  |
| Age <20 years                           | 0.00001-0.00515                                  |
| Age 20-24 years                         | 0.00188-0.00505                                  |
| Age 25-29 years                         | 0.00126-0.00290                                  |
| Age 30-34 years                         | 0.00088-0.00198                                  |
| Age 35-39 years                         | 0.00070-0.00137                                  |
| Age 40-44 years                         | 0.00068-0.00120                                  |
| Age 45-49 years                         | 0.00063-0.00118                                  |
| Age 50-54 years                         | 0.00057-0.00106                                  |
| Age 55-59 years                         | 0.00052-0.00096                                  |
| Age 60+ years                           | 0.00023-0.00087                                  |
| <b>HPV-33 incidence<sup>12,13</sup></b> |  |
| Age <20 years                           | 0.00001-0.00243                                  |
| Age 20-24 years                         | 0.00056-0.00228                                  |
| Age 25-29 years                         | 0.00034-0.00111                                  |
| Age 30-34 years                         | 0.00025-0.00068                                  |
| Age 35-39 years                         | 0.00022-0.00054                                  |
| Age 40-44 years                         | 0.00020-0.00048                                  |
| Age 45-49 years                         | 0.00018-0.00044                                  |
| Age 50-54 years                         | 0.00016-0.00040                                  |
| Age 55-59 years                         | 0.00015-0.00036                                  |
| Age 60+ years                           | 0.00007-0.00034                                  |
| <b>HPV-45 incidence<sup>12,13</sup></b> |  |
| Age <20 years                           | 0.00001-0.00314                                  |
| Age 20-24 years                         | 0.00057-0.00295                                  |
| Age 25-29 years                         | 0.00044-0.00116                                  |
| Age 30-34 years                         | 0.00038-0.00096                                  |
| Age 35-39 years                         | 0.00031-0.00082                                  |
| Age 40-44 years                         | 0.00026-0.00068                                  |
| Age 45-49 years                         | 0.00023-0.00054                                  |

**Appendix Table 1. Natural History Model Parameters, Post-Calibration**

| <b>Model parameter</b>                              | <b>Range of values for 50 best-fitting sets*</b> |
|---|--|
| Age 50-54 years                                     | 0.00019-0.00047                                  |
| Age 55-59 years                                     | 0.00017-0.00042                                  |
| Age 60+ years                                       | 0.00008-0.00038                                  |
| <b>HPV-52 incidence<sup>12,13</sup></b>             |  |
| Age <20 years                                       | 0.00002-0.00711                                  |
| Age 20-24 years                                     | 0.00108-0.00623                                  |
| Age 25-29 years                                     | 0.00074-0.00152                                  |
| Age 30-34 years                                     | 0.00067-0.00104                                  |
| Age 35-39 years                                     | 0.00063-0.00099                                  |
| Age 40-44 years                                     | 0.00060-0.00093                                  |
| Age 45-49 years                                     | 0.00060-0.00096                                  |
| Age 50-54 years                                     | 0.00054-0.00089                                  |
| Age 55-59 years                                     | 0.00049-0.00080                                  |
| Age 60+ years                                       | 0.00022-0.00073                                  |
| <b>HPV-58 incidence<sup>12,13</sup></b>             |  |
| Age <20 years                                       | 0.00001-0.00376                                  |
| Age 20-24 years                                     | 0.00087-0.00366                                  |
| Age 25-29 years                                     | 0.00069-0.00174                                  |
| Age 30-34 years                                     | 0.00059-0.00142                                  |
| Age 35-39 years                                     | 0.00050-0.00123                                  |
| Age 40-44 years                                     | 0.00040-0.00101                                  |
| Age 45-49 years                                     | 0.00036-0.00083                                  |
| Age 50-54 years                                     | 0.00033-0.00076                                  |
| Age 55-59 years                                     | 0.00029-0.00068                                  |
| Age 60+ years                                       | 0.00008-0.00062                                  |
| <b>Other carcinogenic incidence<sup>12,13</sup></b> |  |
| Age <20 years                                       | 0.00003-0.02953                                  |
| Age 20-24 years                                     | 0.01021-0.02870                                  |
| Age 25-29 years                                     | 0.00679-0.02033                                  |
| Age 30-34 years                                     | 0.00498-0.01315                                  |
| Age 35-39 years                                     | 0.00419-0.00980                                  |
| Age 40-44 years                                     | 0.00368-0.00861                                  |
| Age 45-49 years                                     | 0.00341-0.00770                                  |
| Age 50-54 years                                     | 0.00308-0.00705                                  |
| Age 55-59 years                                     | 0.00279-0.00638                                  |
| Age 60+ years                                       | 0.00123-0.00578                                  |
| <b>Noncarcinogenic incidence<sup>12,13</sup></b>    |  |
| Age <20 years                                       | 0.00001-0.03840                                  |
| Age 20-24 years                                     | 0.00260-0.02043                                  |
| Age 25-29 years                                     | 0.00204-0.01950                                  |
| Age 30-34 years                                     | 0.00158-0.01471                                  |
| Age 35-39 years                                     | 0.00143-0.01194                                  |
| Age 40-44 years                                     | 0.00124-0.01083                                  |
| Age 45-49 years                                     | 0.00086-0.00891                                  |
| Age 50-54 years                                     | 0.00065-0.00633                                  |
| Age 55-59 years                                     | 0.00047-0.00474                                  |
| Age 60+ years                                       | 0.00022-0.00356                                  |
| <b>Natural immunity<sup>12†</sup></b>               | 0.502-0.803                                      |
| <b>HPV-16 clearance<sup>12,14‡</sup></b>            |  |
| Year 1  | 0.04189  |
| Year 2  | 0.04075  |
| Year 3  | 0.03390  |

**Appendix Table 1. Natural History Model Parameters, Post-Calibration**

| <b>Model parameter</b>                               | <b>Range of values for 50 best-fitting sets*</b> |
|--|--|
| Year 4   | 0.03189  |
| Year 5+  | 0.01985  |
| <b>HPV-18 clearance<sup>12,14‡</sup></b>             |  |
| Year 1   | 0.07334  |
| Year 2   | 0.06324  |
| Year 3   | 0.05360  |
| Year 4   | 0.02062  |
| Year 5+  | 0.02062  |
| <b>HPV-31 clearance<sup>12,14‡</sup></b>             |  |
| Year 1   | 0.06345  |
| Year 2   | 0.03383  |
| Year 3   | 0.03383  |
| Year 4   | 0.03383  |
| Year 5+  | 0.03383  |
| <b>HPV-33 clearance<sup>12,14‡</sup></b>             |  |
| Year 1   | 0.08345  |
| Year 2   | 0.04496  |
| Year 3   | 0.03616  |
| Year 4   | 0.03616  |
| Year 5+  | 0.03616  |
| <b>HPV-45 clearance<sup>12,14‡</sup></b>             |  |
| Year 1   | 0.07852  |
| Year 2   | 0.04258  |
| Year 3   | 0.04168  |
| Year 4   | 0.03013  |
| Year 5+  | 0.01507  |
| <b>HPV-52 clearance<sup>12,14‡</sup></b>             |  |
| Year 1   | 0.06300  |
| Year 2   | 0.04440  |
| Year 3   | 0.04440  |
| Year 4   | 0.03933  |
| Year 5+  | 0.03933  |
| <b>HPV-58 clearance<sup>12,14‡</sup></b>             |  |
| Year 1   | 0.06557  |
| Year 2   | 0.05443  |
| Year 3   | 0.05397  |
| Year 4   | 0.03332  |
| Year 5+  | 0.01666  |
| <b>Other carcinogenic clearance<sup>12,14‡</sup></b> |  |
| Year 1   | 0.08077  |
| Year 2   | 0.06663  |
| Year 3   | 0.05397  |
| Year 4   | 0.04923  |
| Year 5+  | 0.00509  |
| <b>Noncarcinogenic clearance<sup>12,14,15‡</sup></b> |  |
| Year 1   | 0.05189  |
| Year 2   | 0.05001  |
| Year 3   | 0.03465  |
| Year 4   | 0.03465  |
| Year 5+  | 0.02861  |
| <b>HPV-16 progression to CIN2<sup>12,14‡</sup></b>   |  |
| Year 1   | 0.00171  |



**Appendix Table 1. Natural History Model Parameters, Post-Calibration**

| <b>Model parameter</b>                             | <b>Range of values for 50 best-fitting sets*</b> |
|--|--|
| Year 2   | 0.00242  |
| Year 3   | 0.00258  |
| Year 4   | 0.00552  |
| Year 5   | 0.01500  |
| Years 6-10   | 0.0152-0.04489                                   |
| Years 11+  | 0.0195-0.07437                                   |
| <b>HPV-16 progression to CIN3<sup>12,14†</sup></b> |  |
| Year 1   | 0.00057  |
| Year 2   | 0.00081  |
| Year 3   | 0.00086  |
| Year 4   | 0.00184  |
| Year 5   | 0.00502  |
| Years 6-10   | 0.00509-0.01504                                  |
| Years 11+  | 0.00653-0.02491                                  |
| <b>HPV-18 progression to CIN2<sup>12,14†</sup></b> |  |
| Year 1   | 0.00004  |
| Year 2   | 0.00019  |
| Year 3   | 0.00019  |
| Year 4   | 0.00773  |
| Year 5   | 0.00773  |
| Years 6-10   | 0.00784-0.02314                                  |
| Years 11+  | 0.01005-0.03835                                  |
| <b>HPV-18 progression to CIN3<sup>12,14†</sup></b> |  |
| Year 1   | 0.00001  |
| Year 2   | 0.00005  |
| Year 3   | 0.00005  |
| Year 4   | 0.00194  |
| Year 5   | 0.00194  |
| Years 6-10   | 0.00196-0.00580                                  |
| Years 11+  | 0.00252-0.00961                                  |
| <b>HPV-31 progression to CIN2<sup>12,14†</sup></b> |  |
| Year 1   | 0.00026  |
| Year 2   | 0.00278  |
| Year 3   | 0.00309  |
| Year 4   | 0.00693  |
| Year 5   | 0.00693  |
| Years 6-10   | 0.00702-0.02074                                  |
| Years 11+  | 0.00901-0.03437                                  |
| <b>HPV-31 progression to CIN3<sup>12,14†</sup></b> |  |
| Year 1   | 0.00007  |
| Year 2   | 0.00070  |
| Year 3   | 0.00077  |
| Year 4   | 0.00174  |
| Year 5   | 0.00174  |
| Years 6-10   | 0.00176-0.00520                                  |
| Years 11+  | 0.00226-0.00861                                  |
| <b>HPV-33 progression to CIN2<sup>12,14†</sup></b> |  |
| Year 1   | 0.00072  |
| Year 2   | 0.00072  |
| Year 3   | 0.00494  |
| Year 4   | 0.00494  |
| Year 5   | 0.00494  |

**Appendix Table 1. Natural History Model Parameters, Post-Calibration**

| <b>Model parameter</b>                             | <b>Range of values for 50 best-fitting sets*</b> |
|--|--|
| Years 6-10   | 0.00500-0.01478                                  |
| Years 11+  | 0.00642-0.02449                                  |
| <b>HPV-33 progression to CIN3<sup>12,14†</sup></b> |  |
| Year 1   | 0.00018  |
| Year 2   | 0.00018  |
| Year 3   | 0.00124  |
| Year 4   | 0.00124  |
| Year 5   | 0.00124  |
| Years 6-10   | 0.00125-0.00370                                  |
| Years 11+  | 0.00161-0.00613                                  |
| <b>HPV-45 progression to CIN2<sup>12,14†</sup></b> |  |
| Year 1   | 0.00000  |
| Year 2   | 0.00000  |
| Year 3   | 0.00226  |
| Year 4   | 0.00533  |
| Year 5   | 0.00533  |
| Years 6-10   | 0.00540-0.01596                                  |
| Years 11+  | 0.00693-0.02645                                  |
| <b>HPV-45 progression to CIN3<sup>12,14†</sup></b> |  |
| Year 1   | 0.00000  |
| Year 2   | 0.00000  |
| Year 3   | 0.00056  |
| Year 4   | 0.00134  |
| Year 5   | 0.00134  |
| Years 6-10   | 0.00135-0.00400                                  |
| Years 11+  | 0.00174-0.00662                                  |
| <b>HPV-52 progression to CIN2<sup>12,14†</sup></b> |  |
| Year 1   | 0.00088  |
| Year 2   | 0.00168  |
| Year 3   | 0.00168  |
| Year 4   | 0.00198  |
| Year 5   | 0.00568  |
| Years 6-10   | 0.00575-0.01699                                  |
| Years 11+  | 0.00738-0.02816                                  |
| <b>HPV-52 progression to CIN3<sup>12,14†</sup></b> |  |
| Year 1   | 0.00022  |
| Year 2   | 0.00042  |
| Year 3   | 0.00042  |
| Year 4   | 0.00049  |
| Year 5   | 0.00142  |
| Years 6-10   | 0.00144-0.00426                                  |
| Years 11+  | 0.00185-0.00705                                  |
| <b>HPV-58 progression to CIN2<sup>12,14†</sup></b> |  |
| Year 1   | 0.00059  |
| Year 2   | 0.00247  |
| Year 3   | 0.00247  |
| Year 4   | 0.00462  |
| Year 5   | 0.01025  |
| Years 6-10   | 0.01038-0.03066                                  |
| Years 11+  | 0.01332-0.05080                                  |
| <b>HPV-58 progression to CIN3<sup>12,14†</sup></b> |  |
| Year 1   | 0.00015  |

**Appendix Table 1. Natural History Model Parameters, Post-Calibration**

| <b>Model parameter</b>   | <b>Range of values for 50 best-fitting sets*</b> |
|--|--|
| Year 2   | 0.00062  |
| Year 3   | 0.00062  |
| Year 4   | 0.00116  |
| Year 5   | 0.00257  |
| Years 6-10   | 0.00261-0.00770                                  |
| Years 11+  | 0.00334-0.01275                                  |
| <b>Other carcinogenic HPV progression to CIN2<sup>12,14†</sup></b>   |  |
| Year 1   | 0.00013  |
| Year 2   | 0.00037  |
| Year 3   | 0.00196  |
| Year 4   | 0.00196  |
| Year 5   | 0.00196  |
| Years 6-10   | 0.00199-0.00587                                  |
| Years 11+  | 0.00255-0.00972                                  |
| <b>Other carcinogenic HPV progression to CIN3<sup>12,14†</sup></b>   |  |
| Year 1   | 0.00003  |
| Year 2   | 0.00009  |
| Year 3   | 0.00049  |
| Year 4   | 0.00049  |
| Year 5   | 0.00049  |
| Years 6-10   | 0.00050-0.00147                                  |
| Years 11+  | 0.00064-0.00243                                  |
| <b>Noncarcinogenic HPV progression to CIN2<sup>12,14†</sup></b>      |  |
| Year 1   | 0.00021  |
| Year 2   | 0.00029  |
| Year 3   | 0.00031  |
| Year 4   | 0.00066  |
| Year 5   | 0.00066  |
| Years 6-10   | 0.00067-0.00199                                  |
| Years 11+  | 0.00086-0.00329                                  |
| <b>Noncarcinogenic HPV progression to CIN3<sup>12,14†</sup></b>      |  |
| Year 1   | 0.00002  |
| Year 2   | 0.00003  |
| Year 3   | 0.00003  |
| Year 4   | 0.00007  |
| Year 5   | 0.00007  |
| Years 6-10   | 0.00007-0.00022                                  |
| Years 11+  | 0.00010-0.00037                                  |
| <b>Regression of CIN2 related to HPV-16<sup>12,16-20§</sup></b>      |  |
| Years 1-5  | 0.04500  |
| Years 6-10   | 0.03600  |
| Years 11-20  | 0.02700  |
| Years 21-30  | 0.00180  |
| Years 31-40  | 0.00090  |
| Years 41+  | 0.00045  |
| <b>Regression of CIN2 related to other hrHPV<sup>12,16-20§</sup></b> |  |
| Years 1-5  | 0.05000  |
| Years 6-10   | 0.04000  |
| Years 11-20  | 0.03000  |
| Years 21-30  | 0.00200  |
| Years 31-40  | 0.00100  |
| Years 41+  | 0.00050  |

**Appendix Table 1. Natural History Model Parameters, Post-Calibration**

| <b>Model parameter</b>  | <b>Range of values for 50 best-fitting sets*</b> |
|---|--|
| <b>Regression of CIN3 related to HPV-16<sup>12,16-20§</sup></b>                           |  |
| Years 1-5   | 0.02250  |
| Years 6-10  | 0.01800  |
| Years 11-20   | 0.01350  |
| Years 21-30   | 0.00090  |
| Years 31-40   | 0.00045  |
| Years 41+   | 0.00023  |
| <b>Regression of CIN3 related to other hrHPV<sup>12,16-20§</sup></b>                      |  |
| Years 1-5   | 0.02500  |
| Years 6-10  | 0.02000  |
| Years 11-20   | 0.01500  |
| Years 21-30   | 0.00100  |
| Years 31-40   | 0.00050  |
| Years 41+   | 0.00025  |
| <b>Progression of CIN2 to invasive cancer related to HPV-16<sup>12,21¶</sup></b>          |  |
| Years 1-5   | 0.00003-0.00004                                  |
| Years 6-10  | 0.00004-0.00005                                  |
| Years 11-20   | 0.00086-0.00111                                  |
| Years 21-29   | 0.00251-0.00325                                  |
| Years 30-34   | 0.00502-0.00650                                  |
| Years 35-39   | 0.00544-0.00704                                  |
| Years 40-44   | 0.01171-0.01516                                  |
| Years 45-49   | 0.01255-0.01624                                  |
| Years 50+   | 0.07512-0.09724                                  |
| <b>Progression of CIN2 to invasive cancer related to HPV-18<sup>2,21¶</sup></b>           |  |
| Years 1-5   | 0.00003-0.00004                                  |
| Years 6-10  | 0.00004-0.00005                                  |
| Years 11-20   | 0.00086-0.00110                                  |
| Years 21-29   | 0.00252-0.00323                                  |
| Years 30-34   | 0.00504-0.00646                                  |
| Years 35-39   | 0.00546-0.00700                                  |
| Years 40-44   | 0.01177-0.01508                                  |
| Years 45-49   | 0.01261-0.01615                                  |
| Years 50+   | 0.07547-0.09671                                  |
| <b>Progression of CIN2 to invasive cancer related to HPV-33<sup>12,21¶</sup></b>          |  |
| Years 1-5   | 0.00003-0.00004                                  |
| Years 6-10  | 0.00004-0.00005                                  |
| Years 11-20   | 0.00087-0.00111                                  |
| Years 21-29   | 0.00254-0.00324                                  |
| Years 30-34   | 0.00509-0.00648                                  |
| Years 35-39   | 0.00551-0.00702                                  |
| Years 40-44   | 0.01187-0.01511                                  |
| Years 45-49   | 0.01272-0.01619                                  |
| Years 50+   | 0.07613-0.09693                                  |
| <b>Progression of CIN2 to invasive cancer related to HPV-31/45/52/58<sup>12,21¶</sup></b> |  |
| Years 1-5   | 0.00002  |
| Years 6-10  | 0.00002  |
| Years 11-20   | 0.00057  |
| Years 21-29   | 0.00167  |
| Years 30-34   | 0.00334  |
| Years 35-39   | 0.00362  |
| Years 40-44   | 0.00780  |

**Appendix Table 1. Natural History Model Parameters, Post-Calibration**

| <b>Model parameter</b>  | <b>Range of values for 50 best-fitting sets*</b> |
|---|--|
| Years 45-49   | 0.00835  |
| Years 50+   | 0.05000  |
| <b>Progression of CIN2 to invasive cancer related to other hrHPV<sup>12,21¶</sup></b>     |  |
| Years 1-5   | 0.00000  |
| Years 6-10  | 0.00000  |
| Years 11-20   | 0.00006-0.00011                                  |
| Years 21-29   | 0.00017-0.00033                                  |
| Years 30-34   | 0.00034-0.00067                                  |
| Years 35-39   | 0.00037-0.00072                                  |
| Years 40-44   | 0.00079-0.00156                                  |
| Years 45-49   | 0.00085-0.00167                                  |
| Years 50+   | 0.00085-0.00167                                  |
| <b>Progression of CIN3 to invasive cancer related to HPV-16<sup>12,21¶</sup></b>          |  |
| Years 1-5   | 0.00017-0.00021                                  |
| Years 6-10  | 0.00018-0.00023                                  |
| Years 11-20   | 0.00430-0.00556                                  |
| Years 21-29   | 0.01259-0.01625                                  |
| Years 30-34   | 0.02518-0.03251                                  |
| Years 35-39   | 0.02728-0.03522                                  |
| Years 40-44   | 0.05875-0.07585                                  |
| Years 45-49   | 0.06294-0.08127                                  |
| Years 50+   | 0.07536-0.09731                                  |
| <b>Progression of CIN3 to invasive cancer related to HPV-18<sup>12,21¶</sup></b>          |  |
| Years 1-5   | 0.00017-0.00021                                  |
| Years 6-10  | 0.00018-0.00023                                  |
| Years 11-20   | 0.00431-0.00556                                  |
| Years 21-29   | 0.01259-0.01626                                  |
| Years 30-34   | 0.02519-0.03253                                  |
| Years 35-39   | 0.02729-0.03524                                  |
| Years 40-44   | 0.05877-0.07590                                  |
| Years 45-49   | 0.06297-0.08132                                  |
| Years 50+   | 0.07539-0.09737                                  |
| <b>Progression of CIN3 to invasive cancer related to HPV-33<sup>12,21¶</sup></b>          |  |
| Years 1-5   | 0.00016-0.00021                                  |
| Years 6-10  | 0.00018-0.00023                                  |
| Years 11-20   | 0.00429-0.00553                                  |
| Years 21-29   | 0.01254-0.01618                                  |
| Years 30-34   | 0.02507-0.03237                                  |
| Years 35-39   | 0.02716-0.03506                                  |
| Years 40-44   | 0.05850-0.07552                                  |
| Years 45-49   | 0.06268-0.08092                                  |
| Years 50+   | 0.07505-0.09688                                  |
| <b>Progression of CIN3 to invasive cancer related to HPV-31/45/52/58<sup>12,21¶</sup></b> |  |
| Years 1-5   | 0.00011  |
| Years 6-10  | 0.00012  |
| Years 11-20   | 0.00286  |
| Years 21-29   | 0.00835  |
| Years 30-34   | 0.01670  |
| Years 35-39   | 0.01810  |
| Years 40-44   | 0.03898  |
| Years 45-49   | 0.04176  |
| Years 50+   | 0.05000  |

**Appendix Table 1. Natural History Model Parameters, Post-Calibration**

| <b>Model parameter</b>  | <b>Range of values for 50 best-fitting sets*</b> |
|---|--|
| <b>Progression of CIN3 to invasive cancer related to other hrHPV<sup>†‡</sup></b> |  |
| Years 1-5   | 0.00001-0.00002                                  |
| Years 6-10  | 0.00001-0.00002                                  |
| Years 11-20   | 0.00029-0.00056                                  |
| Years 21-29   | 0.00084-0.00165                                  |
| Years 30-34   | 0.00168-0.00330                                  |
| Years 35-39   | 0.00182-0.00358                                  |
| Years 40-44   | 0.00392-0.00770                                  |
| Years 45-49   | 0.00420-0.00825                                  |
| Years 50+   | 0.00420-0.00825                                  |
| <b>Progression of invasive cancer stages<sup>†,‡‡</sup></b>                       |  |
| Local to regional   | 0.02000  |
| Regional to distant   | 0.02500  |
| <b>Invasive cancer mortality<sup>††</sup></b>                                     |  |
| Local   |  |
| Year 1  | 0.00159  |
| Years 2-3   | 0.00141  |
| Years 4-20  | 0.00094  |
| Regional  |  |
| Year 1  | 0.00946  |
| Years 2-3   | 0.00781  |
| Years 4-20  | 0.00362  |
| Distant   |  |
| Year 1  | 0.02934  |
| Years 2-3   | 0.01947  |
| Years 4-20  | 0.00760  |
| <b>Probability of symptom detection<sup>††,‡‡</sup></b>                           |  |
| Local   | 0.01740  |
| Regional  | 0.07350  |
| Distant   | 0.17460  |

\* Values represent monthly probabilities, unless otherwise noted; values represent the range of probabilities across the 50 best-fitting sets; parameters without a range of values were held constant across the 50 best-fitting sets.

† Natural immunity represents the percentage reduction in risk of subsequent, type-specific infection after a woman has cleared a carcinogenic infection with the same type. Risk reduction is assumed to be constant across age, time, and genotype.

‡ HPV clearance and progression probabilities are a function of time since infection (i.e., persistence).

§ Precancer regression probabilities decrease by time since lesion onset and are constant across carcinogenic HPV types. Given limited data, we assumed that the monthly CIN3 regression probability is 50% of CIN2 regression; 50% regress to type-specific HPV-infected health states and 50% regress to the normal health state.

¶ Precancer progression probabilities increase by time since lesion onset and are constant across carcinogenic HPV types. CIN2 progression is set at 20% of CIN3 progression (for carcinogenic types only).

|| In addition to time since diagnosis, cancer mortality was adjusted for age at diagnosis by applying stage-specific multipliers to the baseline probabilities that ranged from 0.30 to 7.39 for local cancer; 0.39 to 1.30 for regional cancer; and 0.002 to 15.16 for distant cancer.

Abbreviations: CIN2, cervical intraepithelial neoplasia grade 2; CIN3, cervical intraepithelial neoplasia grade 3; HPV, human papillomavirus; HR, high-risk.

**Appendix Table 2. Efficient Cervical Cancer Screening Strategies (Screening End Age 65) in Terms of Colposcopies per Life-Year Gained**

| #, Strategy                         | Colpos | Δ Colpos* | Life-years | Δ Life-years* | Efficiency ratio (Δcolpo/ΔLY) |
|-------------------------------------|--------|-----------|------------|---------------|-------------------------------|
| <b>Scenario A</b>                   |        |           |            |               |                               |
| 0, No screening                     | 0      |           | 63921.34   |               |                               |
| 1, CYTO-3Y, 21                      | 645    | 645       | 64181.89   | 260.56        | 3                             |
| 8, CYTO-3Y, 21 / HPV-5Y (16/18), 30 | 1635   | 990       | 64193.38   | 11.45         | 86                            |
| 7, CYTO-3Y, 21 / HPV-5Y (16/18), 27 | 1861   | 227       | 64194.69   | 1.31          | 173                           |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25 | 2068   | 206       | 64195.39   | 0.69          | 297                           |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25 | 2528   | 461       | 64195.61   | 0.22          | 2,082                         |
| <b>Scenario B</b>                   |        |           |            |               |                               |
| 0, No screening                     | 0      |           | 63921.34   |               |                               |
| 1, CYTO-3Y, 21                      | 645    | 645       | 64181.89   | 260.56        | 3                             |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30 | 1452   | 807       | 64193.07   | 11.14         | 73                            |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27 | 1648   | 196       | 64194.44   | 1.37          | 143                           |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25 | 1826   | 177       | 64195.35   | 0.91          | 195                           |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25  | 2209   | 384       | 64195.53   | 0.18          | 2,188                         |
| <b>Scenario C</b>                   |        |           |            |               |                               |
| 0, No screening                     | 0      |           | 63921.34   |               |                               |
| 1, CYTO-3Y, 21                      | 645    | 645       | 64181.89   | 260.56        | 3                             |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30 | 1452   | 807       | 64193.07   | 11.14         | 73                            |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27 | 1648   | 196       | 64194.44   | 1.37          | 143                           |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25 | 1826   | 177       | 64195.35   | 0.91          | 19                            |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25  | 2209   | 384       | 64195.53   | 0.18          | 2,188                         |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25 | 2530   | 321       | 64195.61   | 0.08          | 3,822                         |
| <b>Scenario D</b>                   |        |           |            |               |                               |
| 0, No screening                     | 0      |           | 63921.34   |               |                               |
| 1, CYTO-3Y, 21                      | 645    | 645       | 64181.89   | 260.56        | 3                             |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30 | 1452   | 807       | 64193.07   | 11.14         | 73                            |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27 | 1648   | 196       | 64194.44   | 1.37          | 143                           |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25 | 1826   | 177       | 64195.35   | 0.91          | 195                           |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25  | 2209   | 384       | 64195.53   | 0.18          | 2,188                         |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25 | 2530   | 321       | 64195.61   | 0.08          | 3,822                         |

\* Incremental values may be slightly different due to rounding.

Abbreviations: Colpos: colposcopies; Cyto: cytology; HPV: human papillomavirus; LY: life-year.

**Appendix Table 3. Efficient Cervical Cancer Screening Strategies (Screening End Age 65 Years) in Terms of Tests per Life-Year Gained**

| #, Strategy                         | Screening tests | Δ Tests* | Life-years | Δ Life-years* | Efficiency ratio (Δtests/ΔLY) |
|-------------------------------------|-----------------|----------|------------|---------------|-------------------------------|
| <b>Scenario A</b>                   |                 |          |            |               |                               |
| 0, No screening                     | 0               |          | 63921.34   |               |                               |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25 | 11771           | 11771    | 64195.39   | 274.05        | 43                            |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25 | 16712           | 4942     | 64195.61   | 0.22          | 22,335                        |
| <b>Scenario B</b>                   |                 |          |            |               |                               |
| 0, No screening                     | 0               |          | 63921.34   |               |                               |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25 | 12042           | 12042    | 64195.35   | 274.01        | 44                            |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25  | 17067           | 5025     | 64195.53   | 0.18          | 28,636                        |
| <b>Scenario C</b>                   |                 |          |            |               |                               |
| 0, No screening                     | 0               |          | 63921.34   |               |                               |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25 | 11771           | 11771    | 64195.39   | 274.05        | 43                            |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25 | 16712           | 4942     | 64195.61   | 0.22          | 22,335                        |
| <b>Scenario D</b>                   |                 |          |            |               |                               |
| 0, No screening                     | 0               |          | 63921.34   |               |                               |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25 | 11771           | 11771    | 64195.39   | 274.05        | 43                            |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25 | 16712           | 4942     | 64195.61   | 0.22          | 22,335                        |

\* Incremental values may be slightly different due to rounding.

Abbreviations: Colpos: colposcopies; Cyto: cytology; HPV: human papillomavirus; LY: life-year.



**Appendix Table 4. Efficient Cervical Cancer Screening Strategies (Screening End Age 65 Years) in Terms of Colposcopies per Cervical Cancer Case Averted\***

| #, Strategy                         | Colpos | Δ Colpos† | Cancer cases averted | Δ Cases averted† | Efficiency ratio (Δcolpo/Δcases averted) |
|-------------------------------------|--------|-----------|----------------------|------------------|--|
| <b>Scenario A</b>                   |        |           |                      |                  |  |
| 0, No screening                     | 0      |           | 0.00                 |                  |  |
| 1, CYTO-3Y, 21                      | 645    | 645       | 16.52                | 16.52            | 39                                       |
| 8, CYTO-3Y, 21 / HPV-5Y (16/18), 30 | 1635   | 990       | 17.81                | 1.29             | 766                                      |
| 7, CYTO-3Y, 21 / HPV-5Y (16/18), 27 | 1861   | 227       | 17.97                | 0.16             | 1,432                                    |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25 | 2068   | 206       | 18.07                | 0.10             | 2,120                                    |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25 | 2530   | 463       | 18.12                | 0.05             | 8,580                                    |
| <b>Scenario B</b>                   |        |           |                      |                  |  |
| 0, No screening                     | 0      |           | 0.00                 |                  |  |
| 1, CYTO-3Y, 21                      | 645    | 645       | 16.52                | 16.52            | 39                                       |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30 | 1452   | 807       | 17.78                | 1.26             | 640                                      |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27 | 1648   | 196       | 17.95                | 0.17             | 1,161                                    |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25 | 1826   | 177       | 18.05                | 0.10             | 1,735                                    |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25  | 2209   | 384       | 18.11                | 0.05             | 7,018                                    |
| <b>Scenario C</b>                   |        |           |                      |                  |  |
| 0, No screening                     | 0      |           | 0.00                 |                  |  |
| 1, CYTO-3Y, 21                      | 645    | 645       | 16.52                | 16.52            | 39                                       |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30 | 1452   | 807       | 17.78                | 1.26             | 640                                      |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27 | 1648   | 196       | 17.95                | 0.17             | 1,161                                    |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25 | 1826   | 177       | 18.05                | 0.10             | 1,735                                    |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25  | 2209   | 384       | 18.11                | 0.05             | 7,018                                    |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25 | 2530   | 321       | 18.12                | 0.01             | 23,974                                   |
| <b>Scenario D</b>                   |        |           |                      |                  |  |
| 0, No screening                     | 0      |           | 0.00                 |                  |  |
| 1, CYTO-3Y, 21                      | 645    | 645       | 16.52                | 16.52            | 39                                       |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30 | 1452   | 807       | 17.78                | 1.26             | 640                                      |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27 | 1648   | 196       | 17.95                | 0.17             | 1,161                                    |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25 | 1826   | 177       | 18.05                | 0.10             | 1,735                                    |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25  | 2209   | 384       | 18.11                | 0.05             | 7,018                                    |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25 | 2530   | 321       | 18.12                | 0.01             | 23,974                                   |

\* Cervical cancer cases averted, compared to no intervention (i.e., natural history).

† Incremental values may be slightly different due to rounding.

Abbreviations: Colpos: colposcopies; Cyto: cytology; HPV: human papillomavirus.

**Appendix Table 5. Efficient Cervical Cancer Screening Strategies (Screening End Age 70 Years) in Terms of Colposcopies per Life-Year Gained**

| #, Strategy                            | Colpos | Δ Colpos* | Life-years | Δ Life-years* | Efficiency ratio (Δcolpo/ΔLY) |
|--|--------|-----------|------------|---------------|-------------------------------|
| <b>Scenario A</b>                      |        |           |            |               |                               |
| 0, No screening                        | 0      |           | 63921.34   |               |                               |
| 1, CYTO-3Y, 21-70                      | 689    | 689       | 64184.58   | 263.24        | 3                             |
| 8, CYTO-3Y, 21 / HPV-5Y (16/18), 30-70 | 1700   | 1011      | 64194.50   | 9.93          | 102                           |
| 7, CYTO-3Y, 21 / HPV-5Y (16/18), 27-70 | 1930   | 230       | 64195.84   | 1.34          | 172                           |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-70 | 2134   | 203       | 64196.47   | 0.63          | 321                           |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-70 | 2620   | 486       | 64196.60   | 0.13          | 3,747                         |
| <b>Scenario B</b>                      |        |           |            |               |                               |
| 0, No screening                        | 0      |           | 63921.34   |               |                               |
| 1, CYTO-3Y, 21-70                      | 689    | 689       | 64184.58   | 263.24        | 3                             |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30-70 | 1508   | 819       | 64194.18   | 9.61          | 85                            |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27-70 | 1706   | 199       | 64195.57   | 1.39          | 143                           |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25-70 | 1881   | 175       | 64196.41   | 0.84          | 208                           |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25-70  | 2283   | 401       | 64196.52   | 0.11          | 3,758                         |
| <b>Scenario C</b>                      |        |           |            |               |                               |
| 0, No screening                        | 0      |           | 63921.34   |               |                               |
| 1, CYTO-3Y, 21-70                      | 689    | 689       | 64184.58   | 263.24        | 3                             |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30-70 | 1508   | 819       | 64194.18   | 9.61          | 85                            |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27-70 | 1706   | 199       | 64195.57   | 1.39          | 143                           |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25-70 | 1881   | 175       | 64196.41   | 0.84          | 208                           |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25-70  | 2283   | 401       | 64196.52   | 0.11          | 3,758                         |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-70 | 2620   | 337       | 64196.60   | 0.08          | 4,014                         |
| <b>Scenario D</b>                      |        |           |            |               |                               |
| 0, No screening                        | 0      |           | 63921.34   |               |                               |
| 1, CYTO-3Y, 21-70                      | 689    | 689       | 64184.58   | 263.24        | 3                             |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30-70 | 1508   | 819       | 64194.18   | 9.61          | 85                            |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27-70 | 1706   | 199       | 64195.57   | 1.39          | 143                           |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25-70 | 1881   | 175       | 64196.41   | 0.84          | 208                           |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25-70  | 2283   | 401       | 64196.52   | 0.11          | 3,758                         |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-70 | 2620   | 337       | 64196.60   | 0.08          | 4,014                         |

\* Incremental values may be slightly different due to rounding.

Abbreviations: Colpos: colposcopies; Cyto: cytology; HPV: human papillomavirus; LY: life-year.

**Appendix Table 6. Efficient Cervical Cancer Screening Strategies (Screening End Age 70 Years) in Terms of Tests per Life-Year Gained**

| #, Strategy                            | Screening tests | Δ Tests* | Life-years | Δ Life-years* | Efficiency ratio (Δtests/ΔLY) |
|--|-----------------|----------|------------|---------------|-------------------------------|
| <b>Scenario A</b>                      |                 |          |            |               |                               |
| 0, No screening                        | 0               |          | 63921.34   |               |                               |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-70 | 12536           | 12536    | 64196.47   | 275.14        | 46                            |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-70 | 17965           | 5428     | 64196.60   | 0.13          | 41,852                        |
| <b>Scenario B</b>                      |                 |          |            |               |                               |
| 0, No screening                        | 0               |          | 63921.34   |               |                               |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25-70 | 12818           | 12818    | 64196.41   | 275.07        | 47                            |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25-70  | 18306           | 5488     | 64196.52   | 0.11          | 51,383                        |
| <b>Scenario C</b>                      |                 |          |            |               |                               |
| 0, No screening                        | 0               |          | 63921.34   |               |                               |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-70 | 12536           | 12536    | 64196.47   | 275.14        | 46                            |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-70 | 17965           | 5428     | 64196.60   | 0.13          | 41,852                        |
| <b>Scenario D</b>                      |                 |          |            |               |                               |
| 0, No screening                        |                 |          | 63921.34   |               |                               |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-70 | 12536           | 12536    | 64196.47   | 275.14        | 46                            |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-70 | 17965           | 5428     | 64196.60   | 0.13          | 41,852                        |

\* Incremental values may be slightly different due to rounding.

Abbreviations: Colpos: colposcopies; Cyto: cytology; HPV: human papillomavirus; LY: life-year.

**Appendix Table 7. Efficient Cervical Cancer Screening Strategies (Screening End Age 70 Years) in Terms of Colposcopies per Cervical Cancer Case Averted\***

| #, Strategy                            | Colpos | Δ Colpos† | Cancer cases averted | Δ Cases averted† | Efficiency ratio (Δcolpo/Δcases averted) |
|--|--------|-----------|----------------------|------------------|--|
| <b>Scenario A</b>                      |        |           |                      |                  |  |
| 0, No screening                        | 0      |           | 0.00                 |                  |  |
| 1, CYTO-3Y, 21-70                      | 689    | 689       | 16.88                | 16.88            | 41                                       |
| 8, CYTO-3Y, 21 / HPV-5Y (16/18), 30-70 | 1700   | 1011      | 18.01                | 1.13             | 898                                      |
| 7, CYTO-3Y, 21 / HPV-5Y (16/18), 27-70 | 1930   | 230       | 18.17                | 0.17             | 1,391                                    |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-70 | 2134   | 203       | 18.26                | 0.09             | 2,375                                    |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-70 | 2620   | 486       | 18.30                | 0.04             | 12,631                                   |
| <b>Scenario B</b>                      |        |           |                      |                  |  |
| 0, No screening                        | 0      |           | 0.00                 |                  |  |
| 1, CYTO-3Y, 21-70                      | 689    | 689       | 16.88                | 16.88            | 41                                       |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30-70 | 1508   | 819       | 17.97                | 1.09             | 748                                      |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27-70 | 1706   | 199       | 18.15                | 0.18             | 1,134                                    |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25-70 | 1881   | 175       | 18.24                | 0.09             | 1,914                                    |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25-70  | 2283   | 401       | 18.28                | 0.04             | 9,848                                    |
| <b>Scenario C</b>                      |        |           |                      |                  |  |
| 0, No screening                        | 0      |           | 0.00                 |                  |  |
| 1, CYTO-3Y, 21-70                      | 689    | 689       | 16.88                | 16.88            | 41                                       |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30-70 | 1508   | 819       | 17.97                | 1.09             | 748                                      |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27-70 | 1706   | 199       | 18.15                | 0.18             | 1,134                                    |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25-70 | 1881   | 175       | 18.24                | 0.09             | 1,914                                    |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25-70  | 2283   | 401       | 18.28                | 0.04             | 9,848                                    |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-70 | 2620   | 337       | 18.30                | 0.01             | 23,361                                   |
| <b>Scenario D</b>                      |        |           |                      |                  |  |
| 0, No screening                        | 0      |           | 0.00                 |                  |  |
| 1, CYTO-3Y, 21-70                      | 689    | 689       | 16.88                | 16.88            | 41                                       |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30-70 | 1508   | 819       | 17.97                | 1.09             | 748                                      |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27-70 | 1706   | 199       | 18.15                | 0.18             | 1,134                                    |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25-70 | 1881   | 175       | 18.24                | 0.09             | 1,914                                    |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25-70  | 2283   | 401       | 18.28                | 0.04             | 9,848                                    |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-70 | 2620   | 337       | 18.30                | 0.01             | 23,361                                   |

\* Cervical cancer cases averted, compared to no intervention (i.e., natural history).

† Incremental values may be slightly different due to rounding.

Abbreviations: Colpos: colposcopies; Cyto: cytology; HPV: human papillomavirus.

**Appendix Table 8. Efficient Cervical Cancer Screening Strategies (Screening End Age 75 Years) in Terms of Colposcopies per Life-Year Gained**

| #, Strategy                            | Colpos | Δ Colpos* | Life-years | Δ Life-years* | Efficiency ratio (Δcolpo/ΔLY) |
|--|--------|-----------|------------|---------------|-------------------------------|
| <b>Scenario A</b>                      |        |           |            |               |                               |
| 0, No screening                        | 0      |           | 63921.34   |               |                               |
| 1, CYTO-3Y, 21-75                      | 724    | 724       | 64185.93   | 264.59        | 3                             |
| 8, CYTO-3Y, 21 / HPV-5Y (16/18), 30-75 | 1752   | 1028      | 64194.93   | 9.00          | 114                           |
| 7, CYTO-3Y, 21 / HPV-5Y (16/18), 27-75 | 1984   | 232       | 64196.36   | 1.43          | 163                           |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-75 | 2186   | 202       | 64196.93   | 0.57          | 353                           |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-75 | 2687   | 501       | 64197.01   | 0.08          | 5,972                         |
| <b>Scenario B</b>                      |        |           |            |               |                               |
| 0, No screening                        | 0      |           | 63921.34   |               |                               |
| 1, CYTO-3Y, 21-75                      | 724    | 724       | 64185.93   | 264.59        | 3                             |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30-75 | 1552   | 827       | 64194.63   | 8.71          | 95                            |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27-75 | 1752   | 200       | 64196.12   | 1.49          | 135                           |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25-75 | 1926   | 174       | 64196.89   | 0.77          | 225                           |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25-75  | 2340   | 414       | 64196.93   | 0.04          | 10,854                        |
| <b>Scenario C</b>                      |        |           |            |               |                               |
| 0, No screening                        | 0      |           | 63921.34   |               |                               |
| 1, CYTO-3Y, 21-75                      | 724    | 724       | 64185.93   | 264.59        | 3                             |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30-75 | 1552   | 827       | 64194.63   | 8.71          | 95                            |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27-75 | 1752   | 200       | 64196.12   | 1.49          | 135                           |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25-75 | 1926   | 174       | 64196.89   | 0.77          | 225                           |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-75 | 2687   | 762       | 64197.01   | 0.12          | 6,239                         |
| <b>Scenario D</b>                      |        |           |            |               |                               |
| 0, No screening                        | 0      |           | 63921.34   |               |                               |
| 1, CYTO-3Y, 21-75                      | 724    | 724       | 64185.93   | 264.59        | 3                             |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30-75 | 1552   | 827       | 64194.63   | 8.71          | 95                            |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27-75 | 1752   | 200       | 64196.12   | 1.49          | 135                           |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25-75 | 1926   | 174       | 64196.89   | 0.77          | 225                           |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-75 | 2687   | 762       | 64197.01   | 0.12          | 6,239                         |

\* Incremental values may be slightly different due to rounding.

Abbreviations: Colpos: colposcopies; Cyto: cytology; HPV: human papillomavirus; LY: life-year.

**Appendix Table 9. Efficient Cervical Cancer Screening Strategies (Screening End Age 75 Years) in Terms of Tests per Life-Year Gained**

| #, Strategy                            | Screening tests | Δ Tests* | Life-years | Δ Life-years* | Efficiency ratio (Δtests/ΔLY) |
|--|-----------------|----------|------------|---------------|-------------------------------|
| <b>Scenario A</b>                      |                 |          |            |               |                               |
| 0, No screening                        | 0               |          | 63921.34   |               |                               |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-75 | 13222           | 13222    | 64196.93   | 275.59        | 48                            |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-75 | 19018           | 5796     | 64197.01   | 0.08          | 69,063                        |
| <b>Scenario B</b>                      |                 |          |            |               |                               |
| 0, No screening                        | 0               |          | 63921.34   |               |                               |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25-75 | 13512           | 13512    | 64196.89   | 275.55        | 49                            |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25-75  | 19380           | 5868     | 64196.93   | 0.04          | 153,816                       |
| <b>Scenario C</b>                      |                 |          |            |               |                               |
| 0, No screening                        | 0               |          | 63921.34   |               |                               |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-75 | 13222           | 13222    | 64196.93   | 275.59        | 48                            |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-75 | 19018           | 5796     | 64197.01   | 0.08          | 69,063                        |
| <b>Scenario D</b>                      |                 |          |            |               |                               |
| 0, No screening                        | 0               |          | 63921.34   |               |                               |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-75 | 13222           | 13222    | 64196.93   | 275.59        | 48                            |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-75 | 19018           | 5796     | 64197.01   | 0.08          | 69,063                        |

\* Incremental values may be slightly different due to rounding.

Abbreviations: Colpos: colposcopies; Cyto: cytology; HPV: human papillomavirus; LY: life-year.

**Appendix Table 10. Efficient Cervical Cancer Screening Strategies (Screening End Age 75 Years) in Terms of Colposcopies per Cervical Cancer Case Averted\***

| #, Strategy                            | Colpos | Δ Colpos† | Cancer cases averted | Δ Cases averted† | Efficiency ratio (Δcolpo/Δcases averted) |
|--|--------|-----------|----------------------|------------------|--|
| <b>Scenario A</b>                      |        |           |                      |                  |  |
| 0, No screening                        | 0      |           | 0.00                 |                  | --                                       |
| 1, CYTO-3Y, 21-75                      | 724    | 724       | 17.10                | 17.10            | 42                                       |
| 8, CYTO-3Y, 21 / HPV-5Y (16/18), 30-75 | 1752   | 1028      | 18.11                | 1.01             | 1,020                                    |
| 7, CYTO-3Y, 21 / HPV-5Y (16/18), 27-75 | 1984   | 232       | 18.28                | 0.17             | 1,333                                    |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-75 | 2186   | 202       | 18.36                | 0.08             | 2,479                                    |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-75 | 2687   | 501       | 18.38                | 0.02             | 23,138                                   |
| <b>Scenario B</b>                      |        |           |                      |                  |  |
| 0, No screening                        | 0      |           | 0.00                 |                  |  |
| 1, CYTO-3Y, 21-75                      | 724    | 724       | 17.10                | 17.10            | 42                                       |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30-75 | 1552   | 827       | 18.08                | 0.98             | 847                                      |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27-75 | 1752   | 200       | 18.26                | 0.19             | 1,081                                    |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25-75 | 1926   | 174       | 18.35                | 0.08             | 2,064                                    |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25-75  | 2340   | 414       | 18.37                | 0.02             | 17,110                                   |
| <b>Scenario C</b>                      |        |           |                      |                  |  |
| 0, No screening                        | 0      |           | 0.00                 |                  |  |
| 1, CYTO-3Y, 21-75                      | 724    | 724       | 17.10                | 17.10            | 42                                       |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30-75 | 1552   | 827       | 18.08                | 0.98             | 847                                      |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27-75 | 1752   | 200       | 18.26                | 0.19             | 1,081                                    |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25-75 | 1926   | 174       | 18.35                | 0.08             | 2,064                                    |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-75 | 2186   | 260       | 18.36                | 0.02             | 15,899                                   |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25-75  | 2340   | 154       | 18.37                | 0.01             | 19,645                                   |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-75 | 2687   | 348       | 18.38                | 0.01             | 25,112                                   |
| <b>Scenario D</b>                      |        |           |                      |                  |  |
| 0, No screening                        | 0      |           | 0.00                 |                  |  |
| 1, CYTO-3Y, 21-75                      | 724    | 724       | 17.10                | 17.10            | 42                                       |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30-75 | 1552   | 827       | 18.08                | 0.98             | 847                                      |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27-75 | 1752   | 200       | 18.26                | 0.19             | 1,081                                    |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25-75 | 1926   | 174       | 18.35                | 0.08             | 2,064                                    |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-75 | 2186   | 260       | 18.36                | 0.02             | 15,899                                   |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25-75  | 2340   | 154       | 18.37                | 0.01             | 19,645                                   |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-75 | 2687   | 348       | 18.38                | 0.01             | 25,112                                   |

\* Cervical cancer cases averted, compared to no intervention (i.e., natural history).

† Incremental values may be slightly different due to rounding.

Abbreviations: Colpos: colposcopies; Cyto: cytology; HPV: human papillomavirus.

**Appendix Table 11. Efficient Cervical Cancer Screening Strategies, Varying HPV Testing Switch Age, Interval, and Screening End Age in Terms of Colposcopies per Life-Year Gained**

| #, Strategy                            | Colpos | Δ Colpos* | Life-years | Δ Life-years* | Efficiency ratio (Δcolpo/ΔLY) |
|--|--------|-----------|------------|---------------|-------------------------------|
| <b>Scenario A</b>                      |        |           |            |               |                               |
| 0, No screening                        | 0      |           | 63921.34   |               |                               |
| 1, CYTO-3Y, 21-65                      | 645    | 645       | 64181.89   | 260.56        | 3                             |
| 1, CYTO-3Y, 21-70                      | 689    | 44        | 64184.58   | 2.65          | 17                            |
| 1, CYTO-3Y, 21-75                      | 724    | 36        | 64185.93   | 1.35          | 26                            |
| 8, CYTO-3Y, 21 / HPV-5Y (16/18), 30-70 | 1700   | 976       | 64194.50   | 8.58          | 114                           |
| 8, CYTO-3Y, 21 / HPV-5Y (16/18), 30-75 | 1752   | 52        | 64194.93   | 0.43          | 123                           |
| 7, CYTO-3Y, 21 / HPV-5Y (16/18), 27-75 | 1984   | 232       | 64196.36   | 1.43          | 163                           |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-75 | 2186   | 202       | 64196.93   | 0.57          | 353                           |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-75 | 2687   | 501       | 64197.01   | 0.08          | 5,972                         |
| <b>Scenario B</b>                      |        |           |            |               |                               |
| 0, No screening                        | 0      |           | 63921.34   |               |                               |
| 1, CYTO-3Y, 21-65                      | 645    | 645       | 64181.89   | 260.56        | 3                             |
| 1, CYTO-3Y, 21-70                      | 689    | 44        | 64184.58   | 2.65          | 17                            |
| 1, CYTO-3Y, 21-75                      | 724    | 36        | 64185.93   | 1.35          | 26                            |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30-70 | 1508   | 783       | 64194.18   | 8.26          | 95                            |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30-75 | 1552   | 44        | 64194.63   | 0.45          | 99                            |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27-75 | 1752   | 200       | 64196.12   | 1.49          | 135                           |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25-75 | 1926   | 174       | 64196.89   | 0.77          | 225                           |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25-75  | 2340   | 414       | 64196.93   | 0.04          | 10,854                        |
| <b>Scenario C</b>                      |        |           |            |               |                               |
| 0, No screening                        | 0      |           | 63921.34   |               |                               |
| 1, CYTO-3Y, 21-65                      | 645    | 645       | 64181.89   | 260.56        | 3                             |
| 1, CYTO-3Y, 21-70                      | 689    | 44        | 64184.58   | 2.65          | 17                            |
| 1, CYTO-3Y, 21-75                      | 724    | 36        | 64185.93   | 1.35          | 26                            |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30-70 | 1508   | 783       | 64194.18   | 8.26          | 95                            |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30-75 | 1552   | 44        | 64194.63   | 0.45          | 99                            |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27-75 | 1752   | 200       | 64196.12   | 1.49          | 135                           |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25-75 | 1926   | 174       | 64196.89   | 0.77          | 225                           |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-75 | 2687   | 762       | 64197.01   | 0.12          | 6,239                         |
| <b>Scenario D</b>                      |        |           |            |               |                               |
| 0, No screening                        | 0      |           | 63921.34   |               |                               |
| 1, CYTO-3Y, 21-65                      | 645    | 645       | 64181.89   | 260.56        | 3                             |
| 1, CYTO-3Y, 21-70                      | 689    | 44        | 64184.58   | 2.65          | 17                            |
| 1, CYTO-3Y, 21-75                      | 724    | 36        | 64185.93   | 1.35          | 26                            |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30-70 | 1508   | 783       | 64194.18   | 8.26          | 95                            |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30-75 | 1552   | 44        | 64194.63   | 0.45          | 99                            |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27-75 | 1752   | 200       | 64196.12   | 1.49          | 135                           |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25-75 | 1926   | 174       | 64196.89   | 0.77          | 225                           |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-75 | 2687   | 762       | 64197.01   | 0.12          | 6,239                         |

\* Incremental values may be slightly different due to rounding.

Abbreviations: Colpos: colposcopies; Cyto: cytology; HPV: human papillomavirus; LY: life-year.



**Appendix Table 12. Efficient Cervical Cancer Screening Strategies, Varying HPV Testing Switch Age, Interval, and Screening End Age in Terms of Tests per Life-Year Gained**

| #, Strategy                            | Screening tests | Δ Tests* | Life-years | Δ Life-years* | Efficiency ratio (Δtests/ΔLY) |
|--|-----------------|----------|------------|---------------|-------------------------------|
| <b>Scenario A</b>                      |                 |          |            |               |                               |
| 0, No screening                        | 0               |          | 63921.34   |               |                               |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-65 | 11771           | 11771    | 64195.39   | 274.05        | 43                            |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-70 | 12536           | 766      | 64196.47   | 1.08          | 707                           |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-75 | 13222           | 685      | 64196.93   | 0.46          | 1,497                         |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-75 | 19018           | 5796     | 64197.01   | 0.08          | 69,063                        |
| <b>Scenario B</b>                      |                 |          |            |               |                               |
| 0, No screening                        | 0               |          | 63921.34   |               |                               |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25-65 | 12042           | 12042    | 64195.35   | 274.01        | 44                            |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25-70 | 12818           | 776      | 64196.41   | 1.06          | 731                           |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25-75 | 13512           | 694      | 64196.89   | 0.48          | 1,444                         |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25-75  | 19380           | 5868     | 64196.93   | 0.04          | 153,816                       |
| <b>Scenario C</b>                      |                 |          |            |               |                               |
| 0, No screening                        | 0               |          | 63921.34   |               |                               |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-65 | 11771           | 11771    | 64195.39   | 274.05        | 43                            |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-70 | 12536           | 766      | 64196.47   | 1.08          | 707                           |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-75 | 13222           | 685      | 64196.93   | 0.46          | 1,497                         |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-75 | 19018           | 5796     | 64197.01   | 0.08          | 69,063                        |
| <b>Scenario D</b>                      |                 |          |            |               |                               |
| 0, No screening                        | 0               |          | 63921.34   |               |                               |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-65 | 11771           | 11771    | 64195.39   | 274.05        | 43                            |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-70 | 12536           | 766      | 64196.47   | 1.08          | 707                           |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-75 | 13222           | 685      | 64196.93   | 0.46          | 1,497                         |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-75 | 19018           | 5796     | 64197.01   | 0.08          | 69,063                        |

\* Incremental values may be slightly different due to rounding.

Abbreviations: Colpos: colposcopies; Cyto: cytology; HPV: human papillomavirus; LY: life-year.

**Appendix Table 13. Efficient Cervical Cancer Screening Strategies, Varying HPV Testing Switch Age, Interval, and Screening End Age in Terms of Colposcopies per Cervical Cancer Case Averted\***

| #, Strategy                            | Colpos | Δ Colpos† | Cancer cases averted | Δ Cases averted† | Efficiency ratio (Δcolpo/Δcases averted) |
|--|--------|-----------|----------------------|------------------|--|
| <b>Scenario A</b>                      |        |           |                      |                  |  |
| 0, No screening                        | 0      |           | 0.00                 |                  |  |
| 1, CYTO-3Y, 21-65                      | 645    | 645       | 16.52                | 16.52            | 39                                       |
| 1, CYTO-3Y, 21-70                      | 689    | 44        | 16.88                | 0.36             | 122                                      |
| 1, CYTO-3Y, 21-75                      | 724    | 36        | 17.10                | 0.22             | 162                                      |
| 8, CYTO-3Y, 21 / HPV-5Y (16/18), 30-75 | 1752   | 1028      | 18.11                | 1.01             | 1,020                                    |
| 7, CYTO-3Y, 21 / HPV-5Y (16/18), 27-75 | 1984   | 232       | 18.28                | 0.17             | 1,333                                    |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-75 | 2186   | 202       | 18.36                | 0.08             | 2,479                                    |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-75 | 2687   | 501       | 18.38                | 0.02             | 23,138                                   |
| <b>Scenario B</b>                      |        |           |                      |                  |  |
| 0, No screening                        | 0      |           | 0.00                 |                  |  |
| 1, CYTO-3Y, 21-65                      | 645    | 645       | 16.52                | 16.52            | 39                                       |
| 1, CYTO-3Y, 21-70                      | 689    | 44        | 16.88                | 0.36             | 122                                      |
| 1, CYTO-3Y, 21-75                      | 724    | 36        | 17.10                | 0.22             | 162                                      |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30-75 | 1552   | 827       | 18.08                | 0.98             | 847                                      |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27-75 | 1752   | 200       | 18.26                | 0.19             | 1,081                                    |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25-75 | 1926   | 174       | 18.35                | 0.08             | 2,064                                    |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25-75  | 2340   | 414       | 18.37                | 0.02             | 17,110                                   |
| <b>Scenario C</b>                      |        |           |                      |                  |  |
| 0, No screening                        | 0      |           | 0.00                 |                  |  |
| 1, CYTO-3Y, 21-65                      | 645    | 645       | 16.52                | 16.52            | 39                                       |
| 1, CYTO-3Y, 21-70                      | 689    | 44        | 16.88                | 0.36             | 122                                      |
| 1, CYTO-3Y, 21-75                      | 724    | 36        | 17.10                | 0.22             | 162                                      |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30-75 | 1552   | 827       | 18.08                | 0.98             | 847                                      |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27-75 | 1752   | 200       | 18.26                | 0.19             | 1,081                                    |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25-75 | 1926   | 174       | 18.35                | 0.08             | 2,064                                    |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-75 | 2186   | 260       | 18.36                | 0.02             | 15,899                                   |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25-75  | 2340   | 154       | 18.37                | 0.01             | 19,645                                   |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-75 | 2687   | 348       | 18.38                | 0.01             | 25,112                                   |
| <b>Scenario D</b>                      |        |           |                      |                  |  |
| 0, No screening                        | 0      |           | 0.00                 |                  |  |
| 1, CYTO-3Y, 21-65                      | 645    | 645       | 16.52                | 16.52            | 39                                       |
| 1, CYTO-3Y, 21-70                      | 689    | 44        | 16.88                | 0.36             | 122                                      |
| 1, CYTO-3Y, 21-75                      | 724    | 36        | 17.10                | 0.22             | 162                                      |
| 14, CYTO-3Y, 21 / HPV-5Y (cyto), 30-75 | 1552   | 827       | 18.08                | 0.98             | 847                                      |
| 13, CYTO-3Y, 21 / HPV-5Y (cyto), 27-75 | 1752   | 200       | 18.26                | 0.19             | 1,081                                    |
| 12, CYTO-4Y, 21 / HPV-5Y (cyto), 25-75 | 1926   | 174       | 18.35                | 0.08             | 2,064                                    |
| 6, CYTO-4Y, 21 / HPV-5Y (16/18), 25-75 | 2186   | 260       | 18.36                | 0.02             | 15,899                                   |
| 9, CYTO-4Y, 21 / HPV-3Y (cyto), 25-75  | 2340   | 154       | 18.37                | 0.01             | 19,645                                   |
| 3, CYTO-4Y, 21 / HPV-3Y (16/18), 25-75 | 2687   | 348       | 18.38                | 0.01             | 25,112                                   |

\* Cervical cancer cases averted, compared to no intervention (i.e., natural history).

† Incremental values may be slightly different due to rounding.

Abbreviations: Colpos: colposcopies; Cyto: cytology; HPV: human papillomavirus.

**Appendix Table 14. Range of Outcomes for Cervical Cancer Screening Strategies Over Lifetime of Screening (Screening End Age 65 Years) Across 50 Best-Fitting Parameter Sets**

| #  | Strategy                               | Per 1,000 women        |                        |                        |                     |                  |                 |                     |                        |                      |                                 |
|----|--|------------------------|------------------------|------------------------|---------------------|------------------|-----------------|---------------------|------------------------|----------------------|---------------------------------|
|    |  | Cyto tests             | HPV tests              | Total tests*           | Colpos              | CIN2,3 detected  | CIN3+ detected† | False positives‡    | CC cases               | CC deaths            | Life-years                      |
| 0  | No screening                           | 0<br>(0-0)             | 0<br>(0-0)             | 0<br>(0-0)             | 0<br>(0-0)          | 0<br>(0-0)       | 0<br>(0-0)      | 0<br>(0-0)          | 18.86<br>(13.05-24.07) | 8.34<br>(5.78-10.76) | 63921.34<br>(63844.9-64006.01)  |
| 1  | CYTO-3Y, 21                            | 13877<br>(13806-13939) | 786<br>(773-797)       | 14662<br>(14579-14736) | 645<br>(593-691)    | 160<br>(121-197) | 46<br>(35-57)   | 484<br>(473-495)    | 2.34<br>(1.75-2.84)    | 0.76<br>(0.54-0.99)  | 64181.89<br>(64178.07-64185.61) |
| 2  | CYTO-3Y, 21 /<br>COTEST-5Y, 30         | 11425<br>(11185-11641) | 8380<br>(8139-8602)    | 19806<br>(19324-20243) | 1630<br>(1393-1864) | 201<br>(150-248) | 54<br>(41-66)   | 1429<br>(1243-1616) | 1.08<br>(0.8-1.34)     | 0.3<br>(0.2-0.43)    | 64192.97<br>(64191.52-64194.54) |
| 3  | CYTO-4Y, 21 /<br>HPV-3Y (16/18),<br>25 | 1905<br>(1798-2020)    | 14807<br>(14432-15116) | 16712<br>(16230-17136) | 2530<br>(2159-2889) | 218<br>(164-269) | 57<br>(43-70)   | 2312<br>(1995-2620) | 0.74<br>(0.52-0.93)    | 0.23<br>(0.13-0.34)  | 64195.61<br>(64194.38-64196.64) |
| 4  | CYTO-3Y, 21 /<br>HPV-3Y (16/18),<br>27 | 2876<br>(2780-2989)    | 13772<br>(13423-14066) | 16648<br>(16203-17056) | 2278<br>(1942-2611) | 214<br>(161-264) | 56<br>(44-70)   | 2063<br>(1781-2348) | 0.83<br>(0.6-1.02)     | 0.25<br>(0.15-0.36)  | 64195.08<br>(64193.92-64196.34) |
| 5  | CYTO-3Y, 21 /<br>HPV-3Y (16/18),<br>30 | 3824<br>(3740-3929)    | 12428<br>(12124-12692) | 16252<br>(15865-16621) | 1978<br>(1688-2273) | 205<br>(153-252) | 54<br>(41-67)   | 1773<br>(1535-2021) | 1.01<br>(0.76-1.26)    | 0.27<br>(0.18-0.38)  | 64193.51<br>(64192.11-64195.08) |
| 6  | CYTO-4Y, 21 /<br>HPV-5Y (16/18),<br>25 | 1706<br>(1625-1793)    | 10065<br>(9703-10377)  | 11771<br>(11328-12170) | 2068<br>(1743-2386) | 211<br>(159-259) | 55<br>(42-68)   | 1857<br>(1585-2127) | 0.79<br>(0.54-1.02)    | 0.25<br>(0.14-0.37)  | 64195.39<br>(64194.02-64196.61) |
| 7  | CYTO-3Y, 21 /<br>HPV-5Y (16/18),<br>27 | 2697<br>(2626-2785)    | 9290<br>(8947-9588)    | 11987<br>(11573-12373) | 1861<br>(1568-2157) | 207<br>(155-255) | 55<br>(42-67)   | 1655<br>(1413-1902) | 0.89<br>(0.63-1.12)    | 0.28<br>(0.16-0.41)  | 64194.69<br>(64193.29-64196.03) |
| 8  | CYTO-3Y, 21 /<br>HPV-5Y (16/18),<br>30 | 3675<br>(3609-3759)    | 8476<br>(8195-8735)    | 12151<br>(11805-12495) | 1635<br>(1382-1897) | 199<br>(149-246) | 53<br>(41-66)   | 1435<br>(1233-1652) | 1.05<br>(0.77-1.31)    | 0.29<br>(0.18-0.4)   | 64193.38<br>(64191.97-64194.94) |
| 9  | CYTO-4Y, 21 /<br>HPV-3Y (cyto),<br>25  | 2277<br>(2085-2415)    | 14790<br>(14425-15090) | 17067<br>(16510-17506) | 2209<br>(1855-2530) | 217<br>(163-267) | 56<br>(43-69)   | 1992<br>(1693-2263) | 0.75<br>(0.53-0.94)    | 0.23<br>(0.13-0.33)  | 64195.53<br>(64194.5-64196.58)  |
| 10 | CYTO-3Y, 21 /<br>HPV-3Y (cyto),<br>27  | 3205<br>(3021-3339)    | 13738<br>(13395-14026) | 16943<br>(16416-17364) | 1992<br>(1673-2288) | 213<br>(160-262) | 56<br>(42-70)   | 1779<br>(1513-2027) | 0.85<br>(0.62-1.06)    | 0.25<br>(0.15-0.36)  | 64194.82<br>(64193.64-64196.11) |
| 11 | CYTO-3Y, 21 /<br>HPV-3Y (cyto),<br>30  | 4102<br>(3931-4226)    | 12397<br>(12097-12655) | 16499<br>(16029-16882) | 1734<br>(1459-1996) | 203<br>(153-251) | 54<br>(41-67)   | 1530<br>(1307-1746) | 1.04<br>(0.78-1.3)     | 0.28<br>(0.18-0.39)  | 64193.19<br>(64191.76-64194.63) |
| 12 | CYTO-4Y, 21 /<br>HPV-5Y (cyto),<br>25  | 1993<br>(1851-2101)    | 10049<br>(9696-10351)  | 12042<br>(11547-12452) | 1826<br>(1515-2114) | 209<br>(157-257) | 55<br>(42-67)   | 1617<br>(1358-1857) | 0.81<br>(0.56-1.04)    | 0.25<br>(0.14-0.38)  | 64195.35<br>(64194.07-64196.59) |
| 13 | CYTO-3Y, 21 /<br>HPV-5Y (cyto),<br>27  | 2950<br>(2810-3055)    | 9273<br>(8938-9564)    | 12223<br>(11748-12619) | 1648<br>(1368-1915) | 205<br>(154-253) | 54<br>(41-67)   | 1443<br>(1215-1663) | 0.91<br>(0.64-1.15)    | 0.28<br>(0.16-0.41)  | 64194.44<br>(64193.09-64195.79) |

**Appendix Table 14. Range of Outcomes for Cervical Cancer Screening Strategies Over Lifetime of Screening (Screening End Age 65 Years) Across 50 Best-Fitting Parameter Sets**

| #  | Strategy                        | Per 1,000 women        |                        |                        |                     |                  |                 |                     |                     |                     |                                 |
|----|---------------------------------|------------------------|------------------------|------------------------|---------------------|------------------|-----------------|---------------------|---------------------|---------------------|---------------------------------|
|    |                                 | Cyto tests             | HPV tests              | Total tests*           | Colpos              | CIN2,3 detected  | CIN3+ detected† | False positives‡    | CC cases            | CC deaths           | Life-years                      |
| 14 | CYTO-3Y, 21 / HPV-5Y (cyto), 30 | 3888<br>(3757-3989)    | 8459<br>(8183-8713)    | 12348<br>(11940-12701) | 1452<br>(1211-1690) | 198<br>(148-244) | 53<br>(40-66)   | 1254<br>(1063-1446) | 1.08<br>(0.79-1.35) | 0.29<br>(0.19-0.42) | 64193.07<br>(64191.49-64194.47) |
| 15 | CYTO-4Y, 21 / COTEST-3Y, 25     | 15723<br>(15411-15979) | 14693<br>(14383-14950) | 30416<br>(29794-30929) | 2535<br>(2188-2851) | 223<br>(167-275) | 57<br>(43-70)   | 2312<br>(2022-2577) | 0.76<br>(0.54-0.95) | 0.23<br>(0.13-0.34) | 64195.5<br>(64194.52-64196.49)  |
| 16 | CYTO-3Y, 21 / COTEST-3Y, 27     | 15765<br>(15470-16009) | 13723<br>(13432-13969) | 29488<br>(28901-29978) | 2303<br>(1990-2596) | 218<br>(163-269) | 57<br>(43-70)   | 2084<br>(1827-2327) | 0.83<br>(0.6-1.03)  | 0.25<br>(0.15-0.36) | 64194.75<br>(64193.45-64195.79) |
| 17 | CYTO-3Y, 21 / COTEST-3Y, 30     | 15456<br>(15192-15674) | 12411<br>(12156-12635) | 27867<br>(27348-28309) | 2021<br>(1750-2280) | 209<br>(156-257) | 55<br>(42-68)   | 1812<br>(1595-2023) | 1.03<br>(0.77-1.28) | 0.27<br>(0.18-0.38) | 64193.17<br>(64191.94-64194.73) |
| 18 | CYTO-4Y, 21 / COTEST-5Y, 25     | 10944<br>(10638-11204) | 9914<br>(9609-10175)   | 20859<br>(20247-21380) | 2029<br>(1723-2313) | 213<br>(160-263) | 55<br>(42-68)   | 1816<br>(1564-2051) | 0.82<br>(0.58-1.06) | 0.26<br>(0.14-0.39) | 64195.26<br>(64194.02-64196.45) |
| 19 | CYTO-3Y, 21 / COTEST-5Y, 27     | 11275<br>(10985-11524) | 9233<br>(8942-9485)    | 20508<br>(19927-21008) | 1846<br>(1570-2110) | 209<br>(156-258) | 55<br>(42-68)   | 1637<br>(1414-1853) | 0.89<br>(0.64-1.11) | 0.27<br>(0.17-0.4)  | 64194.4<br>(64192.85-64195.58)  |

\* Total number of tests, irrespective of primary, triage, or surveillance context.

†CIN3+ includes CIN3 and cervical cancer detected through screening (excludes clinically-detected cancer).

‡Total number of colposcopies that did not result in CIN2, CIN3, or cancer detection.

Abbreviations: CC: cervical cancer; CIN: cervical intraepithelial neoplasia; Colpos: colposcopies; Cyto: cytology; HPV: human papillomavirus.

**Appendix Table 15. Outcomes for Cervical Cancer Screening Strategies Over Lifetime of Screening (Screening End Age 65 Years) Assuming Cytology Sensitivity of 81.5% and Specificity of 88.0%**

| #  | Strategy                         | Per 1,000 women |           |              |        |                 |                  |          |           |            |
|----|----------------------------------|-----------------|-----------|--------------|--------|-----------------|------------------|----------|-----------|------------|
|    |                                  | Cyto tests      | HPV tests | Total tests* | Colpos | CIN2,3 detected | False positives† | CC cases | CC deaths | Life-years |
| 1  | CYTO-3Y, 21                      | 14116           | 1164      | 15280        | 904    | 173             | 730              | 1.99     | 0.66      | 64185.81   |
| 2  | CYTO-3Y, 21 / COTEST-5Y, 30      | 11574           | 8537      | 20111        | 1818   | 208             | 1610             | 0.99     | 0.29      | 64193.80   |
| 3  | CYTO-4Y, 21 / HPV-3Y (16/18), 25 | 1939            | 14829     | 16768        | 2585   | 221             | 2364             | 0.72     | 0.23      | 64195.66   |
| 4  | CYTO-3Y, 21 / HPV-3Y (16/18), 27 | 2934            | 13821     | 16755        | 2352   | 218             | 2134             | 0.79     | 0.24      | 64195.35   |
| 5  | CYTO-3Y, 21 / HPV-3Y (16/18), 30 | 3903            | 12504     | 16407        | 2071   | 210             | 1860             | 0.92     | 0.26      | 64194.18   |
| 6  | CYTO-4Y, 21 / HPV-5Y (16/18), 25 | 1740            | 10088     | 11828        | 2115   | 213             | 1902             | 0.77     | 0.25      | 64195.45   |
| 7  | CYTO-3Y, 21 / HPV-5Y (16/18), 27 | 2755            | 9342      | 12097        | 1930   | 211             | 1718             | 0.85     | 0.27      | 64194.97   |
| 8  | CYTO-3Y, 21 / HPV-5Y (16/18), 30 | 3754            | 8540      | 12294        | 1720   | 205             | 1515             | 0.96     | 0.28      | 64194.05   |
| 9  | CYTO-4Y, 21 / HPV-3Y (cyto), 25  | 2310            | 14812     | 17122        | 2279   | 220             | 2059             | 0.73     | 0.23      | 64195.57   |
| 10 | CYTO-3Y, 21 / HPV-3Y (cyto), 27  | 3261            | 13790     | 17051        | 2079   | 217             | 1862             | 0.80     | 0.24      | 64195.29   |
| 11 | CYTO-3Y, 21 / HPV-3Y (cyto), 30  | 4180            | 12475     | 16656        | 1838   | 209             | 1628             | 0.94     | 0.27      | 64194.00   |
| 12 | CYTO-4Y, 21 / HPV-5Y (cyto), 25  | 2027            | 10072     | 12099        | 1884   | 212             | 1671             | 0.78     | 0.25      | 64195.42   |
| 13 | CYTO-3Y, 21 / HPV-5Y (cyto), 27  | 3006            | 9326      | 12333        | 1726   | 210             | 1516             | 0.86     | 0.27      | 64194.93   |
| 14 | CYTO-3Y, 21 / HPV-5Y (cyto), 30  | 3966            | 8523      | 12490        | 1546   | 204             | 1342             | 0.98     | 0.28      | 64193.86   |
| 15 | CYTO-4Y, 21 / COTEST-3Y, 25      | 15893           | 14859     | 30752        | 2771   | 227             | 2544             | 0.73     | 0.23      | 64195.59   |
| 16 | CYTO-3Y, 21 / COTEST-3Y, 27      | 15952           | 13911     | 29864        | 2548   | 224             | 2324             | 0.78     | 0.24      | 64195.27   |
| 17 | CYTO-3Y, 21 / COTEST-3Y, 30      | 15650           | 12613     | 28263        | 2268   | 215             | 2053             | 0.92     | 0.26      | 64194.05   |
| 18 | CYTO-4Y, 21 / COTEST-5Y, 25      | 11080           | 10046     | 21126        | 2196   | 217             | 1979             | 0.80     | 0.26      | 64195.34   |
| 19 | CYTO-3Y, 21 / COTEST-5Y, 27      | 11439           | 9398      | 20837        | 2027   | 215             | 1812             | 0.84     | 0.27      | 64194.89   |

\* Total number of tests, irrespective of primary, triage, or surveillance context.

† Total number of colposcopies that did not result in CIN2, CIN3, or cancer detection.

Abbreviations: CC: cervical cancer; CIN: cervical intraepithelial neoplasia; Colpos: colposcopies; Cyto: cytology; HPV: human papillomavirus.

**Appendix Table 16. Outcomes for Cervical Cancer Screening Strategies Over Lifetime of Screening (Screening End Age 65 Years) Assuming Cytology Sensitivity of 51.4% and Specificity of 93.6%**

| #  | Strategy                         | Per 1,000 women |           |              |        |                 |                  |          |           |            |
|----|----------------------------------|-----------------|-----------|--------------|--------|-----------------|------------------|----------|-----------|------------|
|    |                                  | Cyto tests      | HPV tests | Total tests* | Colpos | CIN2,3 detected | False positives† | CC cases | CC deaths | Life-years |
| 1  | CYTO-3Y, 21                      | 13723           | 614       | 14336        | 506    | 128             | 377              | 3.64     | 1.16      | 64166.02   |
| 2  | CYTO-3Y, 21 / COTEST-5Y, 30      | 11357           | 8327      | 19684        | 1542   | 185             | 1356             | 1.39     | 0.33      | 64190.18   |
| 3  | CYTO-4Y, 21 / HPV-3Y (16/18), 25 | 1881            | 14805     | 16686        | 2505   | 213             | 2292             | 0.79     | 0.23      | 64195.32   |
| 4  | CYTO-3Y, 21 / HPV-3Y (16/18), 27 | 2835            | 13757     | 16592        | 2240   | 204             | 2036             | 0.95     | 0.26      | 64193.95   |
| 5  | CYTO-3Y, 21 / HPV-3Y (16/18), 30 | 3767            | 12400     | 16168        | 1929   | 191             | 1738             | 1.29     | 0.31      | 64190.86   |
| 6  | CYTO-4Y, 21 / HPV-5Y (16/18), 25 | 1682            | 10063     | 11745        | 2046   | 205             | 1841             | 0.84     | 0.26      | 64195.11   |
| 7  | CYTO-3Y, 21 / HPV-5Y (16/18), 27 | 2656            | 9275      | 11931        | 1828   | 197             | 1631             | 1.01     | 0.29      | 64193.55   |
| 8  | CYTO-3Y, 21 / HPV-5Y (16/18), 30 | 3619            | 8458      | 12077        | 1590   | 185             | 1405             | 1.32     | 0.32      | 64190.74   |
| 9  | CYTO-4Y, 21 / HPV-3Y (cyto), 25  | 2253            | 14792     | 17045        | 2177   | 209             | 1968             | 0.82     | 0.24      | 64195.01   |
| 10 | CYTO-3Y, 21 / HPV-3Y (cyto), 27  | 3165            | 13726     | 16891        | 1949   | 201             | 1747             | 1.01     | 0.27      | 64193.42   |
| 11 | CYTO-3Y, 21 / HPV-3Y (cyto), 30  | 4047            | 12371     | 16418        | 1681   | 188             | 1492             | 1.36     | 0.32      | 64190.10   |
| 12 | CYTO-4Y, 21 / HPV-5Y (cyto), 25  | 1970            | 10051     | 12021        | 1799   | 201             | 1598             | 0.88     | 0.26      | 64194.82   |
| 13 | CYTO-3Y, 21 / HPV-5Y (cyto), 27  | 2910            | 9262      | 12172        | 1611   | 194             | 1416             | 1.07     | 0.30      | 64193.02   |
| 14 | CYTO-3Y, 21 / HPV-5Y (cyto), 30  | 3834            | 8445      | 12280        | 1404   | 183             | 1221             | 1.39     | 0.34      | 64189.96   |
| 15 | CYTO-4Y, 21 / COTEST-3Y, 25      | 15652           | 14632     | 30284        | 2431   | 214             | 2216             | 0.83     | 0.24      | 64195.03   |
| 16 | CYTO-3Y, 21 / COTEST-3Y, 27      | 15680           | 13651     | 29331        | 2192   | 206             | 1986             | 0.99     | 0.26      | 64193.57   |
| 17 | CYTO-3Y, 21 / COTEST-3Y, 30      | 15361           | 12332     | 27693        | 1905   | 192             | 1712             | 1.34     | 0.31      | 64190.34   |
| 18 | CYTO-4Y, 21 / COTEST-5Y, 25      | 10890           | 9870      | 20760        | 1956   | 204             | 1751             | 0.88     | 0.27      | 64194.80   |
| 19 | CYTO-3Y, 21 / COTEST-5Y, 27      | 11202           | 9174      | 20376        | 1764   | 197             | 1567             | 1.05     | 0.29      | 64193.24   |

\* Total number of tests, irrespective of primary, triage, or surveillance context.

† Total number of colposcopies that did not result in CIN2, CIN3, or cancer detection.

Abbreviations: CC: cervical cancer; CIN: cervical intraepithelial neoplasia; Colpos: colposcopies; Cyto: cytology; HPV: human papillomavirus.

**Appendix Table 17. Outcomes for Cervical Cancer Screening Strategies Over Lifetime of Screening (Screening End Age 65 Years), Assuming Lower HPV Sensitivity (Relative Sensitivity of 1.15)**

| #  | Strategy                         | Per 1,000 women |           |              |        |                 |                  |          |           |            |
|----|----------------------------------|-----------------|-----------|--------------|--------|-----------------|------------------|----------|-----------|------------|
|    |                                  | Cyto tests      | HPV tests | Total tests* | Colpos | CIN2,3 detected | False positives† | CC cases | CC deaths | Life-years |
| 1  | CYTO-3Y, 21                      | 13864           | 785       | 14649        | 632    | 158             | 473              | 2.43     | 0.79      | 64180.92   |
| 2  | CYTO-3Y, 21 / COTEST-5Y, 30      | 11128           | 8088      | 19217        | 1308   | 185             | 1123             | 1.52     | 0.47      | 64189.04   |
| 3  | CYTO-4Y, 21 / HPV-3Y (16/18), 25 | 1907            | 14528     | 16435        | 2120   | 206             | 1913             | 0.91     | 0.28      | 64194.24   |
| 4  | CYTO-3Y, 21 / HPV-3Y (16/18), 27 | 2876            | 13519     | 16395        | 1915   | 203             | 1712             | 1.01     | 0.30      | 64193.56   |
| 5  | CYTO-3Y, 21 / HPV-3Y (16/18), 30 | 3820            | 12214     | 16034        | 1671   | 195             | 1476             | 1.19     | 0.33      | 64191.86   |
| 6  | CYTO-4Y, 21 / HPV-5Y (16/18), 25 | 1705            | 9769      | 11474        | 1701   | 195             | 1506             | 1.12     | 0.34      | 64192.65   |
| 7  | CYTO-3Y, 21 / HPV-5Y (16/18), 27 | 2695            | 9025      | 11719        | 1540   | 192             | 1347             | 1.21     | 0.37      | 64191.67   |
| 8  | CYTO-3Y, 21 / HPV-5Y (16/18), 30 | 3670            | 8260      | 11930        | 1362   | 186             | 1175             | 1.33     | 0.38      | 64190.44   |
| 9  | CYTO-4Y, 21 / HPV-3Y (cyto), 25  | 2280            | 14489     | 16770        | 1791   | 203             | 1588             | 0.99     | 0.30      | 64193.53   |
| 10 | CYTO-3Y, 21 / HPV-3Y (cyto), 27  | 3204            | 13466     | 16670        | 1622   | 200             | 1422             | 1.08     | 0.32      | 64192.81   |
| 11 | CYTO-3Y, 21 / HPV-3Y (cyto), 30  | 4098            | 12166     | 16264        | 1423   | 192             | 1230             | 1.26     | 0.35      | 64191.07   |
| 12 | CYTO-4Y, 21 / HPV-5Y (cyto), 25  | 1990            | 9726      | 11715        | 1452   | 190             | 1262             | 1.25     | 0.38      | 64191.17   |
| 13 | CYTO-3Y, 21 / HPV-5Y (cyto), 27  | 2943            | 8986      | 11929        | 1322   | 188             | 1133             | 1.31     | 0.40      | 64190.44   |
| 14 | CYTO-3Y, 21 / HPV-5Y (cyto), 30  | 3880            | 8222      | 12103        | 1176   | 183             | 992              | 1.44     | 0.41      | 64189.18   |
| 15 | CYTO-4Y, 21 / COTEST-3Y, 25      | 15371           | 14344     | 29716        | 2069   | 208             | 1861             | 1.07     | 0.34      | 64193.27   |
| 16 | CYTO-3Y, 21 / COTEST-3Y, 27      | 15447           | 13409     | 28856        | 1890   | 205             | 1684             | 1.12     | 0.35      | 64192.70   |
| 17 | CYTO-3Y, 21 / COTEST-3Y, 30      | 15184           | 12144     | 27328        | 1671   | 197             | 1474             | 1.30     | 0.38      | 64190.99   |
| 18 | CYTO-4Y, 21 / COTEST-5Y, 25      | 10545           | 9517      | 20062        | 1595   | 192             | 1402             | 1.36     | 0.44      | 64190.84   |
| 19 | CYTO-3Y, 21 / COTEST-5Y, 27      | 10920           | 8881      | 19801        | 1464   | 190             | 1273             | 1.39     | 0.45      | 64190.12   |

\* Total number of tests, irrespective of primary, triage, or surveillance context.

† Total number of colposcopies that did not result in CIN2, CIN3, or cancer detection.

Abbreviations: CC: cervical cancer; CIN: cervical intraepithelial neoplasia; Colpos: colposcopies; Cyto: cytology; HPV: human papillomavirus.

**Appendix Table 18. Outcomes for Cervical Cancer Screening Strategies Over Lifetime of Screening (Screening End Age 65 Years), Assuming Imperfect Colposcopy/Biopsy Performance**

| #  | Strategy                         | Per 1,000 women |           |              |        |                 |                  |          |           |            |
|----|----------------------------------|-----------------|-----------|--------------|--------|-----------------|------------------|----------|-----------|------------|
|    |                                  | Cyto tests      | HPV tests | Total tests* | Colpos | CIN2,3 detected | False positives† | CC cases | CC deaths | Life-years |
| 1  | CYTO-3Y, 21                      | 13912           | 793       | 14705        | 672    | 138             | 534              | 2.71     | 0.85      | 64178.64   |
| 2  | CYTO-3Y, 21 / COTEST-5Y, 30      | 11331           | 8282      | 19613        | 1428   | 153             | 1275             | 1.19     | 0.31      | 64192.14   |
| 3  | CYTO-4Y, 21 / HPV-3Y (16/18), 25 | 1918            | 14520     | 16438        | 2142   | 151             | 1991             | 0.77     | 0.23      | 64195.37   |
| 4  | CYTO-3Y, 21 / HPV-3Y (16/18), 27 | 2888            | 13517     | 16405        | 1939   | 152             | 1787             | 0.89     | 0.25      | 64194.63   |
| 5  | CYTO-3Y, 21 / HPV-3Y (16/18), 30 | 3836            | 12212     | 16048        | 1698   | 149             | 1548             | 1.13     | 0.28      | 64192.79   |
| 6  | CYTO-4Y, 21 / HPV-5Y (16/18), 25 | 1715            | 9778      | 11493        | 1739   | 150             | 1589             | 0.82     | 0.26      | 64195.17   |
| 7  | CYTO-3Y, 21 / HPV-5Y (16/18), 27 | 2705            | 9041      | 11747        | 1577   | 151             | 1426             | 0.95     | 0.29      | 64194.24   |
| 8  | CYTO-3Y, 21 / HPV-5Y (16/18), 30 | 3684            | 8264      | 11949        | 1399   | 148             | 1251             | 1.16     | 0.30      | 64192.68   |
| 9  | CYTO-4Y, 21 / HPV-3Y (cyto), 25  | 2295            | 14561     | 16856        | 1859   | 153             | 1705             | 0.79     | 0.23      | 64195.24   |
| 10 | CYTO-3Y, 21 / HPV-3Y (cyto), 27  | 3221            | 13534     | 16755        | 1686   | 154             | 1533             | 0.92     | 0.26      | 64194.35   |
| 11 | CYTO-3Y, 21 / HPV-3Y (cyto), 30  | 4117            | 12225     | 16343        | 1484   | 151             | 1333             | 1.15     | 0.29      | 64192.43   |
| 12 | CYTO-4Y, 21 / HPV-5Y (cyto), 25  | 2006            | 9818      | 11824        | 1529   | 152             | 1377             | 0.84     | 0.26      | 64195.00   |
| 13 | CYTO-3Y, 21 / HPV-5Y (cyto), 27  | 2960            | 9070      | 12031        | 1392   | 152             | 1240             | 0.97     | 0.28      | 64193.99   |
| 14 | CYTO-3Y, 21 / HPV-5Y (cyto), 30  | 3900            | 8292      | 12193        | 1242   | 150             | 1092             | 1.19     | 0.30      | 64192.31   |
| 15 | CYTO-4Y, 21 / COTEST-3Y, 25      | 15594           | 14561     | 30154        | 2192   | 158             | 2034             | 0.78     | 0.23      | 64195.34   |
| 16 | CYTO-3Y, 21 / COTEST-3Y, 27      | 15651           | 13606     | 29257        | 2005   | 158             | 1847             | 0.90     | 0.25      | 64194.23   |
| 17 | CYTO-3Y, 21 / COTEST-3Y, 30      | 15364           | 12315     | 27679        | 1776   | 155             | 1621             | 1.13     | 0.28      | 64192.33   |
| 18 | CYTO-4Y, 21 / COTEST-5Y, 25      | 10803           | 9770      | 20573        | 1739   | 155             | 1583             | 0.85     | 0.27      | 64195.08   |
| 19 | CYTO-3Y, 21 / COTEST-5Y, 27      | 11146           | 9102      | 20248        | 1595   | 155             | 1440             | 0.96     | 0.28      | 64193.86   |

\* Total number of tests, irrespective of primary, triage, or surveillance context.

† Total number of colposcopies that did not result in CIN2, CIN3, or cancer detection.

Abbreviations: CC: cervical cancer; CIN: cervical intraepithelial neoplasia; Colpos: colposcopies; Cyto: cytology; HPV: human papillomavirus.



**Appendix Table 19. Outcomes for Cervical Cancer Screening Strategies Over Lifetime of Screening (Screening End Age 65 Years), Assuming CIN2/CIN3 Treatment Effectiveness of 82%**

| #  | Strategy                         | Per 1,000 women |           |              |        |                 |                  |          |           |            |
|----|----------------------------------|-----------------|-----------|--------------|--------|-----------------|------------------|----------|-----------|------------|
|    |                                  | Cyto tests      | HPV tests | Total tests* | Colpos | CIN2,3 detected | False positives† | CC cases | CC deaths | Life-years |
| 1  | CYTO-3Y, 21                      | 13891           | 791       | 14682        | 668    | 183             | 484              | 2.62     | 0.81      | 64179.95   |
| 2  | CYTO-3Y, 21 / COTEST-5Y, 30      | 11458           | 8412      | 19870        | 1675   | 231             | 1443             | 1.24     | 0.32      | 64191.94   |
| 3  | CYTO-4Y, 21 / HPV-3Y (16/18), 25 | 1906            | 14841     | 16746        | 2584   | 252             | 2331             | 0.85     | 0.24      | 64194.91   |
| 4  | CYTO-3Y, 21 / HPV-3Y (16/18), 27 | 2877            | 13804     | 16682        | 2328   | 247             | 2081             | 0.96     | 0.26      | 64194.44   |
| 5  | CYTO-3Y, 21 / HPV-3Y (16/18), 30 | 3827            | 12457     | 16284        | 2023   | 235             | 1787             | 1.17     | 0.29      | 64192.55   |
| 6  | CYTO-4Y, 21 / HPV-5Y (16/18), 25 | 1707            | 10106     | 11813        | 2121   | 244             | 1877             | 0.91     | 0.26      | 64194.71   |
| 7  | CYTO-3Y, 21 / HPV-5Y (16/18), 27 | 2699            | 9328      | 12026        | 1911   | 238             | 1672             | 1.02     | 0.29      | 64194.07   |
| 8  | CYTO-3Y, 21 / HPV-5Y (16/18), 30 | 3679            | 8511      | 12190        | 1680   | 229             | 1450             | 1.21     | 0.30      | 64192.42   |
| 9  | CYTO-4Y, 21 / HPV-3Y (cyto), 25  | 2276            | 14823     | 17100        | 2264   | 250             | 2013             | 0.87     | 0.24      | 64194.84   |
| 10 | CYTO-3Y, 21 / HPV-3Y (cyto), 27  | 3205            | 13770     | 16975        | 2042   | 245             | 1797             | 0.99     | 0.26      | 64194.17   |
| 11 | CYTO-3Y, 21 / HPV-3Y (cyto), 30  | 4105            | 12425     | 16530        | 1780   | 234             | 1545             | 1.21     | 0.30      | 64192.14   |
| 12 | CYTO-4Y, 21 / HPV-5Y (cyto), 25  | 1994            | 10088     | 12082        | 1879   | 241             | 1637             | 0.93     | 0.27      | 64194.65   |
| 13 | CYTO-3Y, 21 / HPV-5Y (cyto), 27  | 2950            | 9309      | 12260        | 1698   | 237             | 1461             | 1.05     | 0.29      | 64193.79   |
| 14 | CYTO-3Y, 21 / HPV-5Y (cyto), 30  | 3892            | 8493      | 12385        | 1497   | 228             | 1269             | 1.24     | 0.31      | 64192.05   |
| 15 | CYTO-4Y, 21 / COTEST-3Y, 25      | 15755           | 14724     | 30479        | 2589   | 257             | 2332             | 0.88     | 0.24      | 64194.91   |
| 16 | CYTO-3Y, 21 / COTEST-3Y, 27      | 15796           | 13753     | 29549        | 2354   | 251             | 2102             | 0.97     | 0.26      | 64194.13   |
| 17 | CYTO-3Y, 21 / COTEST-3Y, 30      | 15485           | 12438     | 27923        | 2066   | 240             | 1826             | 1.18     | 0.29      | 64192.16   |
| 18 | CYTO-4Y, 21 / COTEST-5Y, 25      | 10982           | 9951      | 20934        | 2082   | 246             | 1836             | 0.95     | 0.27      | 64194.68   |
| 19 | CYTO-3Y, 21 / COTEST-5Y, 27      | 11310           | 9267      | 20577        | 1896   | 241             | 1654             | 1.03     | 0.29      | 64193.76   |

\* Total number of tests, irrespective of primary, triage, or surveillance context.

† Total number of colposcopies that did not result in CIN2, CIN3, or cancer detection.

Abbreviations: CC: cervical cancer; CIN: cervical intraepithelial neoplasia; Colpos: colposcopies; Cyto: cytology; HPV: human papillomavirus.

**Appendix Table 20. Outcomes for Cervical Cancer Screening Strategies Over Lifetime of Screening (Screening End Age 65 Years), Assuming Referral to Colposcopy for LSIL or Worse Result on Cytology Triage**

| #  | Strategy                         | Per 1,000 women |           |              |        |                 |                  |          |           |            |
|----|----------------------------------|-----------------|-----------|--------------|--------|-----------------|------------------|----------|-----------|------------|
|    |                                  | Cyto tests      | HPV tests | Total tests* | Colpos | CIN2,3 detected | False positives† | CC cases | CC deaths | Life-years |
| 1  | CYTO-3Y, 21                      | 13877           | 786       | 14662        | 645    | 160             | 484              | 2.34     | 0.76      | 64181.89   |
| 2  | CYTO-3Y, 21 / COTEST-5Y, 30      | 11425           | 8380      | 19806        | 1630   | 201             | 1429             | 1.08     | 0.30      | 64192.97   |
| 3  | CYTO-4Y, 21 / HPV-3Y (16/18), 25 | 1905            | 14799     | 16703        | 2488   | 217             | 2271             | 0.74     | 0.23      | 64195.53   |
| 4  | CYTO-3Y, 21 / HPV-3Y (16/18), 27 | 2876            | 13761     | 16636        | 2240   | 213             | 2027             | 0.83     | 0.25      | 64195.01   |
| 5  | CYTO-3Y, 21 / HPV-3Y (16/18), 30 | 3824            | 12419     | 16242        | 1947   | 204             | 1743             | 1.02     | 0.27      | 64193.44   |
| 6  | CYTO-4Y, 21 / HPV-5Y (16/18), 25 | 1705            | 10056     | 11761        | 2036   | 209             | 1827             | 0.80     | 0.25      | 64195.30   |
| 7  | CYTO-3Y, 21 / HPV-5Y (16/18), 27 | 2697            | 9282      | 11978        | 1834   | 205             | 1629             | 0.90     | 0.28      | 64194.63   |
| 8  | CYTO-3Y, 21 / HPV-5Y (16/18), 30 | 3675            | 8469      | 12144        | 1612   | 198             | 1413             | 1.06     | 0.29      | 64193.31   |
| 9  | CYTO-4Y, 21 / HPV-3Y (cyto), 25  | 2275            | 14780     | 17056        | 2148   | 214             | 1934             | 0.77     | 0.23      | 64195.41   |
| 10 | CYTO-3Y, 21 / HPV-3Y (cyto), 27  | 3204            | 13725     | 16929        | 1938   | 210             | 1727             | 0.87     | 0.25      | 64194.65   |
| 11 | CYTO-3Y, 21 / HPV-3Y (cyto), 30  | 4101            | 12386     | 16487        | 1689   | 201             | 1487             | 1.07     | 0.28      | 64192.92   |
| 12 | CYTO-4Y, 21 / HPV-5Y (cyto), 25  | 1992            | 10038     | 12030        | 1779   | 206             | 1574             | 0.83     | 0.26      | 64195.21   |
| 13 | CYTO-3Y, 21 / HPV-5Y (cyto), 27  | 2948            | 9263      | 12211        | 1608   | 202             | 1405             | 0.93     | 0.28      | 64194.27   |
| 14 | CYTO-3Y, 21 / HPV-5Y (cyto), 30  | 3888            | 8450      | 12338        | 1419   | 196             | 1222             | 1.10     | 0.30      | 64192.79   |
| 15 | CYTO-4Y, 21 / COTEST-3Y, 25      | 15723           | 14693     | 30416        | 2535   | 223             | 2312             | 0.76     | 0.23      | 64195.50   |
| 16 | CYTO-3Y, 21 / COTEST-3Y, 27      | 15765           | 13723     | 29488        | 2303   | 218             | 2084             | 0.83     | 0.25      | 64194.75   |
| 17 | CYTO-3Y, 21 / COTEST-3Y, 30      | 15456           | 12411     | 27867        | 2021   | 209             | 1812             | 1.03     | 0.27      | 64193.17   |
| 18 | CYTO-4Y, 21 / COTEST-5Y, 25      | 10944           | 9914      | 20859        | 2029   | 213             | 1816             | 0.82     | 0.26      | 64195.26   |
| 19 | CYTO-3Y, 21 / COTEST-5Y, 27      | 11275           | 9233      | 20508        | 1846   | 209             | 1637             | 0.89     | 0.27      | 64194.40   |

\* Total number of tests, irrespective of primary, triage, or surveillance context.

† Total number of colposcopies that did not result in CIN2, CIN3, or cancer detection.

Abbreviations: CC: cervical cancer; CIN: cervical intraepithelial neoplasia; Colpos: colposcopies; Cyto: cytology; HPV: human papillomavirus.

**Appendix Table 21. Outcomes for Cervical Cancer Screening Strategies Over Lifetime of Screening (Screening End Age 65 Years), Assuming 6-Month Followup for HPV-Positive, Cytology-Negative Women**

| #  | Strategy                         | Per 1,000 women |           |              |        |                 |                  |          |           |            |
|----|----------------------------------|-----------------|-----------|--------------|--------|-----------------|------------------|----------|-----------|------------|
|    |                                  | Cyto tests      | HPV tests | Total tests* | Colpos | CIN2,3 detected | False positives† | CC cases | CC deaths | Life-years |
| 1  | CYTO-3Y, 21                      | 13877           | 786       | 14662        | 645    | 160             | 484              | 2.34     | 0.76      | 64181.89   |
| 2  | CYTO-3Y, 21 / COTEST-5Y, 30      | 11720           | 8674      | 20394        | 1865   | 203             | 1661             | 1.04     | 0.29      | 64193.28   |
| 3  | CYTO-4Y, 21 / HPV-3Y (16/18), 25 | 1909            | 15226     | 17135        | 2825   | 220             | 2604             | 0.72     | 0.23      | 64195.63   |
| 4  | CYTO-3Y, 21 / HPV-3Y (16/18), 27 | 2881            | 14176     | 17057        | 2537   | 216             | 2321             | 0.79     | 0.24      | 64195.19   |
| 5  | CYTO-3Y, 21 / HPV-3Y (16/18), 30 | 3828            | 12769     | 16596        | 2194   | 206             | 1988             | 0.98     | 0.26      | 64193.60   |
| 6  | CYTO-4Y, 21 / HPV-5Y (16/18), 25 | 1711            | 10442     | 12153        | 2297   | 213             | 2083             | 0.75     | 0.24      | 64195.47   |
| 7  | CYTO-3Y, 21 / HPV-5Y (16/18), 27 | 2700            | 9579      | 12279        | 2055   | 208             | 1847             | 0.86     | 0.27      | 64194.82   |
| 8  | CYTO-3Y, 21 / HPV-5Y (16/18), 30 | 3679            | 8755      | 12434        | 1800   | 201             | 1599             | 1.01     | 0.28      | 64193.57   |
| 9  | CYTO-4Y, 21 / HPV-3Y (cyto), 25  | 2285            | 15385     | 17670        | 2639   | 220             | 2419             | 0.73     | 0.23      | 64195.59   |
| 10 | CYTO-3Y, 21 / HPV-3Y (cyto), 27  | 3214            | 14307     | 17521        | 2372   | 215             | 2156             | 0.80     | 0.24      | 64194.96   |
| 11 | CYTO-3Y, 21 / HPV-3Y (cyto), 30  | 4109            | 12880     | 16989        | 2053   | 206             | 1847             | 0.99     | 0.26      | 64193.55   |
| 12 | CYTO-4Y, 21 / HPV-5Y (cyto), 25  | 2004            | 10585     | 12589        | 2159   | 213             | 1946             | 0.76     | 0.24      | 64195.53   |
| 13 | CYTO-3Y, 21 / HPV-5Y (cyto), 27  | 2956            | 9694      | 12650        | 1933   | 208             | 1725             | 0.87     | 0.27      | 64194.65   |
| 14 | CYTO-3Y, 21 / HPV-5Y (cyto), 30  | 3896            | 8863      | 12759        | 1696   | 200             | 1495             | 1.02     | 0.27      | 64193.58   |
| 15 | CYTO-4Y, 21 / COTEST-3Y, 25      | 16245           | 15215     | 31460        | 2959   | 226             | 2733             | 0.72     | 0.22      | 64195.74   |
| 16 | CYTO-3Y, 21 / COTEST-3Y, 27      | 16212           | 14170     | 30382        | 2675   | 221             | 2453             | 0.79     | 0.24      | 64194.96   |
| 17 | CYTO-3Y, 21 / COTEST-3Y, 30      | 15836           | 12790     | 28626        | 2333   | 211             | 2122             | 0.98     | 0.26      | 64193.52   |
| 18 | CYTO-4Y, 21 / COTEST-5Y, 25      | 11335           | 10306     | 21641        | 2349   | 216             | 2132             | 0.78     | 0.25      | 64195.48   |
| 19 | CYTO-3Y, 21 / COTEST-5Y, 27      | 11588           | 9545      | 21133        | 2122   | 212             | 1910             | 0.86     | 0.27      | 64194.57   |

\* Total number of tests, irrespective of primary, triage, or surveillance context.

† Total number of colposcopies that did not result in CIN2, CIN3, or cancer detection.

Abbreviations: CC: cervical cancer; CIN: cervical intraepithelial neoplasia; Colpos: colposcopies; Cyto: cytology; HPV: human papillomavirus.

**Appendix Table 22. Outcomes for Cervical Cancer Screening Strategies Over Lifetime of Screening (Screening End Age 65 Years), Assuming 24-Month Followup for HPV-Positive, Cytology-Negative Women**

| #  | Strategy                         | Per 1,000 women |           |              |        |                 |                  |          |           |            |
|----|----------------------------------|-----------------|-----------|--------------|--------|-----------------|------------------|----------|-----------|------------|
|    |                                  | Cyto tests      | HPV tests | Total tests* | Colpos | CIN2,3 detected | False positives† | CC cases | CC deaths | Life-years |
| 1  | CYTO-3Y, 21                      | 13878           | 786       | 14663        | 646    | 161             | 484              | 2.35     | 0.76      | 64181.83   |
| 2  | CYTO-3Y, 21 / COTEST-5Y, 30      | 11098           | 8052      | 19151        | 1343   | 196             | 1146             | 1.14     | 0.30      | 64192.47   |
| 3  | CYTO-4Y, 21 / HPV-3Y (16/18), 25 | 1894            | 14199     | 16094        | 2176   | 214             | 1962             | 0.77     | 0.24      | 64195.40   |
| 4  | CYTO-3Y, 21 / HPV-3Y (16/18), 27 | 2870            | 13319     | 16189        | 1975   | 211             | 1764             | 0.85     | 0.25      | 64194.93   |
| 5  | CYTO-3Y, 21 / HPV-3Y (16/18), 30 | 3820            | 12050     | 15870        | 1721   | 203             | 1518             | 1.03     | 0.27      | 64193.46   |
| 6  | CYTO-4Y, 21 / HPV-5Y (16/18), 25 | 1700            | 9675      | 11375        | 1797   | 207             | 1590             | 0.82     | 0.26      | 64195.21   |
| 7  | CYTO-3Y, 21 / HPV-5Y (16/18), 27 | 2693            | 8939      | 11632        | 1627   | 203             | 1424             | 0.92     | 0.28      | 64194.53   |
| 8  | CYTO-3Y, 21 / HPV-5Y (16/18), 30 | 3672            | 8199      | 11872        | 1441   | 197             | 1244             | 1.07     | 0.29      | 64193.30   |
| 9  | CYTO-4Y, 21 / HPV-3Y (cyto), 25  | 2255            | 13905     | 16160        | 1686   | 209             | 1477             | 0.84     | 0.24      | 64195.16   |
| 10 | CYTO-3Y, 21 / HPV-3Y (cyto), 27  | 3192            | 13060     | 16252        | 1536   | 206             | 1330             | 0.90     | 0.26      | 64194.24   |
| 11 | CYTO-3Y, 21 / HPV-3Y (cyto), 30  | 4093            | 11834     | 15927        | 1353   | 199             | 1153             | 1.09     | 0.28      | 64192.81   |
| 12 | CYTO-4Y, 21 / HPV-5Y (cyto), 25  | 1982            | 9469      | 11451        | 1424   | 201             | 1223             | 0.89     | 0.26      | 64194.97   |
| 13 | CYTO-3Y, 21 / HPV-5Y (cyto), 27  | 2940            | 8753      | 11693        | 1299   | 198             | 1100             | 0.98     | 0.29      | 64193.87   |
| 14 | CYTO-3Y, 21 / HPV-5Y (cyto), 30  | 3883            | 8041      | 11924        | 1160   | 193             | 967              | 1.14     | 0.30      | 64192.60   |
| 15 | CYTO-4Y, 21 / COTEST-3Y, 25      | 14994           | 13964     | 28958        | 2018   | 215             | 1803             | 0.82     | 0.24      | 64195.09   |
| 16 | CYTO-3Y, 21 / COTEST-3Y, 27      | 15153           | 13111     | 28264        | 1848   | 212             | 1636             | 0.89     | 0.25      | 64194.24   |
| 17 | CYTO-3Y, 21 / COTEST-3Y, 30      | 14947           | 11902     | 26849        | 1640   | 204             | 1436             | 1.08     | 0.28      | 64192.63   |
| 18 | CYTO-4Y, 21 / COTEST-5Y, 25      | 10486           | 9457      | 19943        | 1636   | 205             | 1430             | 0.88     | 0.27      | 64194.76   |
| 19 | CYTO-3Y, 21 / COTEST-5Y, 27      | 10849           | 8806      | 19655        | 1501   | 203             | 1298             | 0.97     | 0.29      | 64193.82   |

\* Total number of tests, irrespective of primary, triage, or surveillance context.

† Total number of colposcopies that did not result in CIN2, CIN3, or cancer detection.

Abbreviations: CC: cervical cancer; CIN: cervical intraepithelial neoplasia; Colpos: colposcopies; Cyto: cytology; HPV: human papillomavirus.

**Appendix Table 23. Outcomes for Cervical Cancer Screening Strategies Over Lifetime of Screening (Screening End Age 65 Years) Assuming Colposcopy Referral for All HPV-Positive Women**

| #  | Strategy                              | Per 1,000 women |           |              |        |                 |                  |          |           |            |
|----|---------------------------------------|-----------------|-----------|--------------|--------|-----------------|------------------|----------|-----------|------------|
|    |                                       | Cyto tests      | HPV tests | Total tests* | Colpos | CIN2,3 detected | False positives† | CC cases | CC deaths | Life-years |
| 1  | CYTO-3Y, 21                           | 13876           | 786       | 14662        | 645    | 160             | 484              | 2.34     | 0.76      | 64181.93   |
| 2  | CYTO-3Y, 21 / COTEST-5Y, 30*          | 11424           | 8379      | 19803        | 1629   | 201             | 1428             | 1.08     | 0.30      | 64192.98   |
| 20 | CYTO-3Y, 21 / HPV-3Y (colposcopy), 25 | 1145            | 15029     | 16175        | 3261   | 220             | 3041             | 0.74     | 0.23      | 64195.59   |
| 21 | CYTO-3Y, 21 / HPV-3Y (colposcopy), 27 | 2258            | 14041     | 16299        | 2923   | 216             | 2706             | 0.81     | 0.24      | 64195.10   |
| 22 | CYTO-3Y, 21 / HPV-3Y (colposcopy), 30 | 3364            | 12652     | 16016        | 2515   | 207             | 2308             | 0.98     | 0.27      | 64193.61   |
| 23 | CYTO-3Y, 21 / HPV-5Y (colposcopy), 25 | 1145            | 10353     | 11499        | 2632   | 214             | 2417             | 0.78     | 0.25      | 64195.43   |
| 24 | CYTO-3Y, 21 / HPV-5Y (colposcopy), 27 | 2258            | 9514      | 11772        | 2341   | 209             | 2132             | 0.87     | 0.28      | 64194.72   |
| 25 | CYTO-3Y, 21 / HPV-5Y (colposcopy), 30 | 3364            | 8687      | 12051        | 2037   | 201             | 1835             | 1.02     | 0.28      | 64193.50   |

\* Total number of tests, irrespective of primary, triage, or surveillance context.

† Total number of colposcopies that did not result in CIN2, CIN3, or cancer detection.

Abbreviations: CC: cervical cancer; CIN: cervical intraepithelial neoplasia; Colpos: colposcopies; Cyto: cytology; HPV: human papillomavirus.

**Appendix Table 24. Outcomes for Cervical Cancer Screening Strategies Over Lifetime of Screening (Screening End Age 65 Years) in Women Vaccinated Against HPV-16/18\***

| #  | Strategy                         | Per 1,000 women |           |              |        |                 |                  |          |           |            |
|----|----------------------------------|-----------------|-----------|--------------|--------|-----------------|------------------|----------|-----------|------------|
|    |                                  | Cyto tests      | HPV tests | Total tests† | Colpos | CIN2,3 detected | False positives‡ | CC cases | CC deaths | Life-years |
| 1  | CYTO-3Y, 21                      | 13742           | 763       | 14504        | 539    | 87              | 452              | 0.89     | 0.28      | 64193.31   |
| 2  | CYTO-3Y, 21 / COTEST-5Y, 30      | 10909           | 7922      | 18831        | 1221   | 110             | 1111             | 0.34     | 0.07      | 64197.65   |
| 3  | CYTO-4Y, 21 / HPV-3Y (16/18), 25 | 1911            | 14012     | 15923        | 1549   | 116             | 1433             | 0.24     | 0.06      | 64198.41   |
| 4  | CYTO-3Y, 21 / HPV-3Y (16/18), 27 | 2852            | 13013     | 15866        | 1396   | 114             | 1282             | 0.27     | 0.07      | 64198.18   |
| 5  | CYTO-3Y, 21 / HPV-3Y (16/18), 30 | 3775            | 11783     | 15558        | 1217   | 109             | 1108             | 0.33     | 0.07      | 64197.74   |
| 6  | CYTO-4Y, 21 / HPV-5Y (16/18), 25 | 1699            | 9341      | 11041        | 1300   | 111             | 1188             | 0.25     | 0.06      | 64198.38   |
| 7  | CYTO-3Y, 21 / HPV-5Y (16/18), 27 | 2663            | 8601      | 11263        | 1173   | 109             | 1063             | 0.29     | 0.08      | 64198.08   |
| 8  | CYTO-3Y, 21 / HPV-5Y (16/18), 30 | 3620            | 7950      | 11571        | 1040   | 106             | 934              | 0.34     | 0.07      | 64197.75   |
| 9  | CYTO-4Y, 21 / HPV-3Y (cyto), 25  | 1910            | 14012     | 15923        | 1549   | 115             | 1434             | 0.24     | 0.06      | 64198.39   |
| 10 | CYTO-3Y, 21 / HPV-3Y (cyto), 27  | 2853            | 13015     | 15868        | 1398   | 113             | 1284             | 0.27     | 0.07      | 64198.19   |
| 11 | CYTO-3Y, 21 / HPV-3Y (cyto), 30  | 3775            | 11783     | 15558        | 1217   | 109             | 1108             | 0.33     | 0.07      | 64197.74   |
| 12 | CYTO-4Y, 21 / HPV-5Y (cyto), 25  | 1700            | 9342      | 11041        | 1300   | 111             | 1188             | 0.25     | 0.06      | 64198.36   |
| 13 | CYTO-3Y, 21 / HPV-5Y (cyto), 27  | 2663            | 8601      | 11264        | 1173   | 109             | 1063             | 0.29     | 0.08      | 64198.08   |
| 14 | CYTO-3Y, 21 / HPV-5Y (cyto), 30  | 3621            | 7952      | 11572        | 1041   | 106             | 935              | 0.34     | 0.07      | 64197.75   |
| 15 | CYTO-4Y, 21 / COTEST-3Y, 25      | 15019           | 14014     | 29033        | 1881   | 123             | 1759             | 0.24     | 0.06      | 64198.42   |
| 16 | CYTO-3Y, 21 / COTEST-3Y, 27      | 15097           | 13097     | 28194        | 1711   | 120             | 1591             | 0.27     | 0.06      | 64198.19   |
| 17 | CYTO-3Y, 21 / COTEST-3Y, 30      | 14868           | 11881     | 26749        | 1506   | 114             | 1392             | 0.32     | 0.07      | 64197.74   |
| 18 | CYTO-4Y, 21 / COTEST-5Y, 25      | 10282           | 9276      | 19558        | 1508   | 116             | 1391             | 0.26     | 0.07      | 64198.30   |
| 19 | CYTO-3Y, 21 / COTEST-5Y, 27      | 10634           | 8635      | 19269        | 1370   | 114             | 1256             | 0.29     | 0.07      | 64198.10   |

\* HPV-16/18 vaccination is assumed to provide 100% protection against HPV-16/18 infections over the lifetime.

† Total number of tests, irrespective of primary, triage, or surveillance context.

‡ Total number of colposcopies that did not result in CIN2, CIN3, or cancer detection.

Abbreviations: CC: cervical cancer; CIN: cervical intraepithelial neoplasia; Colpos: colposcopies; Cyto: cytology; HPV: human papillomavirus.