




Canagliflozin

Revised: December 3, 2018.

CASRN: 928672-86-0



Drug Levels and Effects

Summary of Use during Lactation

No information is available on the clinical use of canagliflozin during breastfeeding. Canagliflozin is an uncharged molecule that is 99% protein bound in plasma, so it is unlikely to pass into breastmilk in clinically important amounts. The manufacturer does not recommend canagliflozin during breastfeeding because of a theoretical risk to the infant's developing kidney. An alternate drug may be preferred, especially while nursing a newborn or preterm infant.

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site .

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Alternate Drugs to Consider

Acarbose, Glipizide, Glyburide, Insulin, Metformin, Miglitol

Substance Identification

Substance Name

Canagliflozin

CAS Registry Number

928672-86-0

Drug Class

Breast Feeding

Lactation

Hypoglycemic Agents

Sodium-Glucose Cotransporter 2 Inhibitors

SGLT2 Inhibitors