




## Leflunomide

Revised: December 3, 2018.

CASRN: 75706-12-6



## Drug Levels and Effects

### Summary of Use during Lactation

No information is available on the use of leflunomide during breastfeeding, so an alternate drug may be preferred, especially while nursing a newborn or preterm infant.[1][2]

### Drug Levels

*Maternal Levels.* Relevant published information was not found as of the revision date.

*Infant Levels.* Relevant published information was not found as of the revision date.

### Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

**Disclaimer:** Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site .

## Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

## Alternate Drugs to Consider

(Rheumatoid Arthritis) Auranofin, Gold Sodium Thiomalate, Hydroxychloroquine, Infliximab, Methotrexate, Penicillamine, Sulfasalazine

## References

1. Sammaritano LR, Bermas BL. Rheumatoid arthritis medications and lactation. *Curr Opin Rheumatol*. 2014;26:354-60. PubMed PMID: 24614280.
2. Flint J, Panchal S, Hurrell A et al. BSR and BHPR guideline on prescribing drugs in pregnancy and breastfeeding-Part I: standard and biologic disease modifying anti-rheumatic drugs and corticosteroids. *Rheumatology (Oxford)*. 2016;55:1693-7. PubMed PMID: 26750124.

## Substance Identification

### Substance Name

Leflunomide

### CAS Registry Number

75706-12-6

### Drug Class

Breast Feeding

Lactation

Antirheumatic Agents

Immunosuppressive Agents