




Apixaban

Revised: December 3, 2018.

CASRN: 503612-47-3



Drug Levels and Effects

Summary of Use during Lactation

Because no information is available on the use of apixaban during breastfeeding and the drug is orally absorbable, an alternate drug is preferred while nursing a newborn or preterm infant.[1][2][3]

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site .

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Alternate Drugs to Consider

Dalteparin, Enoxaparin, Heparin, Warfarin

References

1. Bates SM, Greer IA, Middeldorp S et al. VTE, thrombophilia, antithrombotic therapy, and pregnancy: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest*. 2012;141 (2 Suppl):e691S-736S. PubMed PMID: 22315276.
2. Vanassche T, Vandenbrielle C, Peerlinck K, Verhamme P. Pharmacotherapy with oral Xa inhibitors for venous thromboembolism. *Expert Opin Pharmacother*. 2015;16:645-58. PubMed PMID: 25554350.
3. Cohen H, Arachchillage DR, Beyer-Westendorf J et al. Direct oral anticoagulants and women. *Semin Thromb Hemost*. 2016. PubMed PMID: 27706531.

Substance Identification

Substance Name

Apixaban

CAS Registry Number

503612-47-3

Drug Class

Breast Feeding

Lactation

Anticoagulants

Antithrombins