




Ceftolozane and Tazobactam

Revised: December 3, 2018.

CASRN: 1613740-46-7



Drug Levels and Effects

Summary of Use during Lactation

No information is available on the clinical use of ceftolozane-tazobactam during breastfeeding. No serious adverse effects have been reported for other cephalosporin antibiotics during breastfeeding. Occasionally disruption of the infant's gastrointestinal flora, resulting in diarrhea or thrush have been reported with cephalosporins, but these effects have not been adequately evaluated. Ceftolozane-tazobactam is acceptable in nursing mothers.

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site .

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Substance Identification

Substance Name

Ceftolozane and Tazobactam

CAS Registry Number

1613740-46-7

Drug Class

Breast Feeding

Lactation

Anti-Infective Agents

Antibacterial Agents

Cephalosporins