




Bivalirudin

Revised: December 3, 2018.

CASRN: 128270-60-0



Drug Levels and Effects

Summary of Use during Lactation

Because no information is available on the use of bivalirudin during breastfeeding, an alternate drug is preferred. [1]

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site .

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Alternate Drugs to Consider

Acenocoumarol, Dalteparin, Enoxaparin, Heparin, Warfarin

References

1. Bates SM, Greer IA, Middeldorp S et al. VTE, thrombophilia, antithrombotic therapy, and pregnancy: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. Chest. 2012;141 (2 Suppl):e691S-736S. PubMed PMID: 22315276.

Substance Identification

Substance Name

Bivalirudin

CAS Registry Number

128270-60-0

Drug Class

Breast Feeding

Lactation

Anticoagulants

Antithrombins