




Mometasone, Nasal

Revised: October 31, 2018.

CASRN: 105102-22-5



Drug Levels and Effects

Summary of Use during Lactation

Neither inhaled mometasone nor mometasone nasal implants have been studied during breastfeeding. Although not measured, the amounts of nasal corticosteroids absorbed into the maternal bloodstream and excreted into breastmilk are probably too small to affect a breastfed infant. Reviewers and an expert panel consider inhaled corticosteroids acceptable to use during breastfeeding.[1][2][3]

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Effects in Breastfed Infants

None reported with any corticosteroid.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Alternate Drugs to Consider

(Inhaled) Beclomethasone, Budesonide

References

1. Greenberger PA, Patterson R. The management of asthma during pregnancy and lactation. Clin Rev Allergy. 1987;5:317-24. PubMed PMID: 3319123.
2. Ellsworth A. Pharmacotherapy of asthma while breastfeeding. J Hum Lact. 1994;10:39-41. PubMed PMID: 7619245.
3. National Heart, Lung, and Blood Institute, National Asthma Education and Prevention Program Asthma and Pregnancy Working Group. NAEPP expert panel report. Managing asthma during pregnancy: recommendations for pharmacologic treatment-2004 update. 2004;1-57. Available at: <http://www.nhlbi.nih.gov/health/prof/lung/asthma/astpreg.htm>

Substance Identification

Substance Name

Mometasone, Nasal

CAS Registry Number

105102-22-5

Drug Class

Breast Feeding

Lactation

Anti-allergic Agents

Anti-Asthmatic Agents

Corticosteroids, Inhaled

Glucocorticoids

Anti-Inflammatory Agents