




Lithium

Revised: March 16, 2020.

CASRN: 554-13-2



Drug Levels and Effects

Summary of Use during Lactation

Lithium excretion into breastmilk and concentrations in infant serum are highly variable. Although lithium appears on many lists of drugs contraindicated during breastfeeding,[1] many sources do not consider it an absolute contraindication, especially in infants over 2 months of age and during lithium monotherapy.[2-6] Numerous reports exist of infants who were breastfed during maternal lithium therapy without any signs of toxicity or developmental problems. Most were breastfed from birth and some continued to nurse for up to 1 year of maternal lithium therapy. Some reports suggest that lithium in milk can adversely affect the infant acutely when its elimination is impaired, as in dehydration or in newborn or premature infants. Neonates may also have transplacentally acquired serum lithium levels. Because maternal lithium requirements and dosage may be increased during pregnancy, maternal serum levels should be monitored frequently postpartum and

dosage reduced as necessary to avoid excessive infant exposure via breastmilk.[7] The long-term effects of lithium on infants are not certain, but limited data indicate no obvious problems in growth and development.[8]

Lithium may be used in mothers of fullterm infants who are willing and able to monitor their infants. Discontinuing lithium 24 to 48 hours before Cesarean section delivery or at the onset of spontaneous labor and resuming the prepregnancy lithium dose immediately after delivery should minimize the infant's serum lithium concentration at birth.[9] Some investigators recommend monitoring infant serum lithium, serum creatinine, BUN, and TSH in intervals ranging from "periodic" to every 4 to 12 weeks during breastfeeding and maternal lithium therapy.[4,10,11] One systematic review recommends infant lithium serum level, thyroid and renal function tests only at 10 days postpartum, then only if the infant serum lithium is 0.3 mEq/L or greater or if clinical signs of toxicity appear.[12] However, others recommend close pediatric follow-up of the infant and only selective laboratory monitoring (i.e., serum lithium, TSH, BUN) if clinically indicated by unusual behavior, restlessness, feeding difficulties, sedation or abnormal growth and development. Infants who are preterm, dehydrated, or have an infection, should receive hydration and assessed for lithium toxicity.[6]

Drug Levels

Maternal Levels. Most older reports of lithium levels in breastmilk did not characterized breastmilk lithium excretion in a rigorous manner. Random milk levels have been reported to range from 0.12 to 0.7 mEq/L and appear to be rather consistent at about 40 to 45% of the simultaneous maternal serum level.[13-16] The milk concentration of lithium in one woman was found to be inversely proportional to milk volume. The milk to plasma ratio was found to be directly proportional to serum lithium; the ratio was about 1 with low serum lithium and 1.5 with higher serum lithium levels, indicating disproportionately higher lithium excretion into milk with higher serum levels.[17]

From data in papers published up to 1990 concerning 6 infants, it is estimated that a fully breastfed infant would receive about 26% (range 11 to 42%) of the maternal weight-adjusted dosage of lithium.[14-19] However, many of these case reports were of poor quality.

A more recent case series of 11 mothers that used better techniques found the average infant dosage to be 12.2% (range 0 to 30%) of the maternal weight-adjusted dosage.[20] The reason for the difference between the older and more recent data is not apparent.

A case series reported 10 mothers taking lithium carbonate in an average daily dosage of 850 mg (range 600 to 1200 mg daily) for bipolar disorder. Lithium milk levels in 26 milk samples taken between 8.1 and 27.5 weeks postpartum averaged 0.35 mEq/L (range 0.19 to 0.48 mEq/L). No difference was found between concentrations in fore- and hindmilk samples.[11]

One woman who was taking 1200 mg of lithium carbonate daily (dosage schedule and product not reported) had a milk lithium concentration of 0.41 mmol/L at 20 hours after her previous dose. This was about the same as her simultaneous blood concentration.[21]

Infant Levels. The serum lithium levels in the breastfed infants of mothers taking lithium have ranged from 10 to 50% of simultaneous maternal serum lithium levels.[14,18,20,22] One infant, who previously had a serum level about 50% of the mother's, became dehydrated and developed a serum level that was double that of the mother's.[22]

A 5-day-old infant developed cyanosis, lethargy, ECG T-wave inversion probably caused by lithium in breastmilk. The infant's serum lithium concentration was 0.6 mEq/L at 5 days of age and 0.21 mEq/L on day 7, 2 days after breastfeeding was discontinued.[18]

Ten exclusively breastfed infants whose mothers were taking an average of 850 mg daily of lithium carbonate had serum levels of 0.16 mEq/L or 24% (range 11 to 56%) of maternal serum levels.[11]

Two breastfed infants (extent not stated) were reported whose mothers were taking lithium carbonate during pregnancy and postpartum. The first mother was taking 1200 mg daily and the infant's serum concentrations were 0.11 mmol/L on day 4 and undetectable (<0.1 mmol/L) on days 6 and 10 postpartum. The second mother was taking 900 mg daily and her breastfed infant's serum lithium concentration was undetectable (<0.3 mmol/L) on day 3 of life. Both infants had other serum concentration measurements that were in the therapeutic or toxic ranges, but displayed no symptoms. These samples apparently had been collected in tubes containing lithium heparin.[21]

A woman took lithium carbonate orally 800 mg daily during pregnancy and postpartum while exclusively breastfeeding her infant. The mother had a serum lithium concentration of 0.74 mmol/L at 15 days postpartum. The infant's serum lithium concentration was 0.26 mmol/L at this time. Further serum lithium concentrations in the infant were 0.23 mmol/L at 1 and 2 months of age and 0.17 mmol/L at 6 months of age.[23]

Three mothers took lithium carbonate during pregnancy and breastfeeding. One mother took a dosage of 900 mg daily during 2 pregnancies. Her infants' serum lithium concentrations were 0.08 mEq/L at 31 days and 183 days postpartum in the first pregnancy and 0.11 mEq/L at 43 days of age in the second pregnancy. These values were 11%, 17% and 15% of the mother's simultaneous serum lithium concentrations, respectively. The second mother also took a dosage of 900 mg daily. Her infant's serum lithium concentration was 0.08 mEq/L at 39 days postpartum, or 10% of the mother's simultaneous serum lithium concentration. The third mother took a dosage of 1350 mg daily. Her infant's serum lithium concentration was 0.11 mEq/L at 31 days postpartum, or 11% of the mother's simultaneous serum lithium concentration.[9]

A woman with bipolar disorder was taking lithium 600 to 900 mg daily during pregnancy and postpartum. At 10 days postpartum, her infant's serum lithium level was 0.26 mmol/L, which was 58% of the maternal trough serum level. Lithium was discontinued. First quetiapine then aripiprazole were substituted for lithium, but after 2 to 3 weeks she switched back to lithium at an unspecified dosage. After several weeks of treatment, her infant's lithium level was 0.2 mmol/L.[24]

A woman with bipolar disorder took prolonged-release lithium carbonate 400 mg (Plenur-Faes Pharma, Spain) every 12 hours during pregnancy and postpartum. She breastfed her infant exclusively for 33 days, but introduced supplements for 16 days because of slow weight gain. After the 16 days, she exclusively breastfed her infant until 2.5 months of age, when mixed feeding was begun. The infant's serum lithium levels were monitored at 17 days, 1 month, 3.5 months and 5.5 months of age. Lithium levels were less than 0.2 mmol/L at all these time points.[25]

Effects in Breastfed Infants

In older reports, at least 24 infants have been reported to have been breastfed during maternal lithium therapy without any signs of toxicity or developmental problems. All were breastfed from birth and some continued to nurse for up to 6 months of maternal lithium therapy.[11,14-16,20,21,23]

A 5-day-old infant developed cyanosis, lethargy, ECG T-wave inversion probably caused by lithium in breastmilk.[18] The mother had been receiving the long-acting diuretic chlorthalidone prior to delivery which probably decreased the infant's lithium elimination and increased the neonate's lithium serum levels. Another case of probable infant lithium intoxication appeared only after the infant had a cold which may have led to dehydration and decreased lithium excretion.[22,26] Two other infants had slight increases in thyrotropin (TSH) levels at 8 and 4 weeks of age, respectively, after lithium exposure that began during pregnancy. Elevated TSH continued until maternal lithium was stopped in one,[11] and normalized by 2 months postpartum in the other, despite continued exclusive breastfeeding.[23]

Three mothers took lithium carbonate during pregnancy and breastfeeding. The first infant was born to a mother who also took bupropion 300 mg and levothyroxine 50 to 75 mcg daily. She breastfed beyond 1 year of

age. Her infant did not regain birth weight by 15 days of age, was somewhat hypotonic at 2 months of age, and was treated for gross and fine motor delay for the first year of life. The mother had a second infant on the same drug regimen. She exclusively breastfed her infant who developed normally without hypotonia. A second mother was taking a lithium dosage of 900 mg daily. Her infant gained weight slowly, but weight gain increased with breastfeeding support and she exclusively breastfed her infant for 4 months. A third mother was taking 1350 mg of lithium daily as well as escitalopram 10 mg, levothyroxine 25 mcg and heparin (dosage not stated) daily during pregnancy and breastfeeding. Her infant was normal and was exclusively breastfed until 8 weeks of age when the maternal serum lithium concentration was excessive at 2.0 mEq/L. Breastfeeding was withheld for 2 days and the dosage lowered to 600 mg daily. She then breastfed successfully until 7 months of age.[9]

A woman with bipolar disorder took prolonged-release lithium carbonate 400 mg every 12 hours during pregnancy and postpartum. She breastfed her infant exclusively for 33 days, but introduced supplements for 16 days because of slow weight gain. After the 16 days, she exclusively breastfed her infant until 2.5 months of age, when mixed feeding was begun. The infant was monitored at 17 days, 1 month, 3.5 months and 5.5 months of age. No infant side effects were observed at any time. Lithium levels were not detectable, and serum creatinine and thyroid-stimulating hormone levels were normal.[25]

Effects on Lactation and Breastmilk

Lithium increases serum prolactin.[27-29] Galactorrhea was reported in a women taking lithium carbonate for 50 days. Lactation ceased with lithium discontinuation.[29] The prolactin level in a mother with established lactation may not affect her ability to breastfeed.

References

1. Poels EMP, Bijma HH, Galbally M, et al. Lithium during pregnancy and after delivery: A review. *Int J Bipolar Disord.* 2018;6:26. PubMed PMID: 30506447.
2. Pacchiarotti I, Leon-Caballero J, Murru A, et al. Mood stabilizers and antipsychotics during breastfeeding: Focus on bipolar disorder. *Eur Neuropsychopharmacol.* 2016;26:1562–78. PubMed PMID: 27568278.
3. Uguz F, Sharma V. Mood stabilizers during breastfeeding: A systematic review of the recent literature. *Bipolar Disord.* 2016;18:325–33. PubMed PMID: 27297617.
4. Khan SJ, Fersh ME, Ernst C, et al. Bipolar disorder in pregnancy and postpartum: Principles of management. *Curr Psychiatry Rep.* 2016;18:13. PubMed PMID: 26781551.
5. Galbally M, Bergink V, Vigod SN, et al. Breastfeeding and lithium: Is breast always best? *Lancet Psychiatry.* 2018;5:534–6. PubMed PMID: 29731410.
6. Newmark RL, Bogen DL, Wisner KL, et al. Risk-benefit assessment of infant exposure to lithium through breast milk: A systematic review of the literature. *Int Rev Psychiatry.* 2019;31:295–304. PubMed PMID: 31180257.
7. Westin AA, Brekke M, Molden E, et al. Changes in drug disposition of lithium during pregnancy: A retrospective observational study of patient data from two routine therapeutic drug monitoring services in Norway. *BMJ Open.* 2017;7:e015738. PubMed PMID: 28249852.
8. Grandjean EM, Aubry JM. Lithium: updated human knowledge using an evidence-based approach: part III: clinical safety. *CNS Drugs.* 2009;23:397–418. PubMed PMID: 19453201.
9. Bogen DL, Sit D, Genovese A, et al. Three cases of lithium exposure and exclusive breastfeeding. *Arch Womens Ment Health.* 2012;15:69–72. PubMed PMID: 22277970.
10. Ketter TA, Suppes T, Morrell MJ, et al. Reproductive health and bipolar disorder. *CNS Spectr.* 2006;11:1–16. doi: [10.1017/S1092852900025608](https://doi.org/10.1017/S1092852900025608).
11. Viguera AC, Newport DJ, Ritchie J, et al. Lithium in breast milk and nursing infants: Clinical implications. *Am J Psychiatry.* 2007;164:342–5. PubMed PMID: 17267800.
12. Imaz ML, Torra M, Soy D, et al. Clinical lactation studies of lithium: A systematic review. *Front Pharmacol.* 2019;10:1005. PubMed PMID: 31551795.

13. Schou M, Amdisen A. Lithium and pregnancy--III, lithium ingestion by children breast-fed by women on lithium treatment. *Br Med J*. 1973;2:138. PubMed PMID: 4699592.
14. Sykes PA, Quarrie J, Alexander FW. Lithium carbonate and breast-feeding. *Br Med J*. 1976;2:1299. PubMed PMID: 1000200.
15. Weinstein MR, Goldfield M. Lithium carbonate treatment during pregnancy. *Dis Nerv Syst*. 1969;30:828–32. PubMed PMID: 5369134.
16. Wise MG, Javors MA, Funderburg LG, et al. Lithium levels in bodily fluids of a nursing mother and infant. *Lithium*. 1990;1:189–91.
17. Shimizu M, Matsuda H, Sakaue N, et al. *Seishin Shinkeigaku Zasshi*. 1981;83:399–405. [A few findings on lithium levels in mother milk -- from one case which fell into a manic state after childbirth]. PubMed PMID: 7330110.
18. Tunnessen WW Jr, Hertz CG. Toxic effects of lithium in newborn infants: A commentary. *J Pediatr*. 1972;81:804–7. PubMed PMID: 5074360.
19. Fries H. Lithium in pregnancy. *Lancet* 1970;295:1233. Letter. PMID: 4192416
20. Moretti ME, Koren G, Verjee Z, et al. Monitoring lithium in breast milk: an individualized approach for breast-feeding mothers. *Ther Drug Monit*. 2003;25:364–6. PubMed PMID: 12766565.
21. Tanaka T, Moretti ME, Verjee ZH, et al. A pitfall of measuring lithium levels in neonates. *Ther Drug Monit*. 2008;30:752–4. PubMed PMID: 19057375.
22. Skausig OB, Schou M. *Ugeskr Laeger*. 1977;139:400–1. [Breast feeding during lithium therapy]. PubMed PMID: 841726.
23. Marín Gabriel MA, Olza Fernández I, Donoso E, et al. *An Pediatr (Barc)*. 2011;75:67–8. [Lithium and artificial breastmilk; or is maternal breastfeeding better?]. PubMed PMID: 21420368.
24. Frew JR. Psychopharmacology of bipolar I disorder during lactation: A case report of the use of lithium and aripiprazole in a nursing mother. *Arch Womens Ment Health*. 2015;18:135–6. PubMed PMID: 25352315.
25. Marín Gabriel MA, Malalana Martínez AM, Olza Fernández I. *Arch Argent Pediatr*. 2018;116:e319–e21. [Lithium while breastfeeding: Case report]. PubMed PMID: 29557625.
26. Schou M. Lithium treatment during pregnancy, delivery, and lactation: An update. *J Clin Psychiatry*. 1990;51:410–3. PubMed PMID: 2211538.
27. Basturk M, Karaaslan F, Esel E, et al. Effects of short and long-term lithium treatment on serum prolactin levels in patients with bipolar affective disorder. *Prog Neuropsychopharmacol Biol Psychiatry*. 2001;25:315–22. PubMed PMID: 11294478.
28. Cowen PJ, Cohen PR, McCance SL, et al. 5-HT neuroendocrine responses during psychotropic drug treatment: an investigation of the effects of lithium. *J Neurosci Methods*. 1990;34:201–5. PubMed PMID: 2259242.
29. Ohishi K, Higashimura T. A case of manic state in which lactation occurred after Li₂CO₃ administration. *Folia Psychiatr Neurol Jpn*. 1983;37:33–6. PubMed PMID: 6411540.

Substance Identification

Substance Name

Lithium

CAS Registry Number

554-13-2

Drug Class

Breast Feeding

Lactation

Lithium