




Trimipramine

Revised: February 7, 2019.

CASRN: 739-71-9



Drug Levels and Effects

Summary of Use during Lactation

Because there is no published experience with trimipramine during breastfeeding, other agents may be preferred, especially while nursing a newborn or preterm infant.

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Effects in Breastfed Infants

Published information on trimipramine was not found as of the revision date. Follow-up for 1 to 3 years in a group of 20 breastfed infants whose mothers were taking a tricyclic antidepressant found no adverse effects on growth and development.[1] Two small controlled studies indicate that other tricyclic antidepressants have no adverse effect on infant development.[2][3] In another study, 25 infants whose mothers took a tricyclic antidepressant during pregnancy and lactation were tested formally between 15 to 71 months and found to have normal growth and development.[4] However, none of the mothers in any of these studies took trimipramine.

Effects on Lactation and Breastmilk

Trimipramine increases serum prolactin levels[5] and has caused at least one case of galactorrhea.[6] The maternal prolactin level in a mother with established lactation may not affect her ability to breastfeed.

An observational study looked at outcomes of 2859 women who took an antidepressant during the 2 years prior to pregnancy. Compared to women who did not take an antidepressant during pregnancy, mothers who took an antidepressant during all 3 trimesters of pregnancy were 37% less likely to be breastfeeding upon hospital discharge. Mothers who took an antidepressant only during the third trimester were 75% less likely to be breastfeeding at discharge. Those who took an antidepressant only during the first and second trimesters did not have a reduced likelihood of breastfeeding at discharge.[7] The antidepressants used by the mothers were not specified.

A retrospective cohort study of hospital electronic medical records from 2001 to 2008 compared women who had been dispensed an antidepressant during late gestation (n = 575) to those who had a psychiatric illness but did not receive an antidepressant (n = 1552) and mothers who did not have a psychiatric diagnosis (n = 30,535). Women who received an antidepressant were 37% less likely to be breastfeeding at discharge than women without a psychiatric diagnosis, but no less likely to be breastfeeding than untreated mothers with a psychiatric diagnosis.[8] None of the mothers were taking trimipramine.

Alternate Drugs to Consider

Nortriptyline, Paroxetine, Sertraline

References

1. Misri S, Sivertz K. Tricyclic drugs in pregnancy and lactation: a preliminary report. *Int J Psychiatry Med.* 1991;21:157-71. PubMed PMID: 1894455.
2. Buist A, Janson H. Effect of exposure to dothiepin and northiaden in breast milk on child development. *Br J Psychiatry.* 1995;167:370-3. PubMed PMID: 7496646.
3. Yoshida K, Smith B, Craggs M et al. Investigation of pharmacokinetics and possible adverse effects in infants exposed to tricyclic antidepressants in breast-milk. *J Affective Disord.* 1997;43:225-37. PubMed PMID: 9186793.
4. Nulman I, Rovet J, Stewart DE et al. Child development following exposure to tricyclic antidepressants or fluoxetine throughout fetal life: a prospective, controlled study. *Am J Psychiatry.* 2002;159:1889-95. PubMed PMID: 12411224.
5. Wiegand M, Berger M. Action of trimipramine on sleep and pituitary hormone secretion. *Drugs.* 1989;38 (Suppl 1):35-42. PubMed PMID: 2612388.
6. Wolfspenger M, Greil W. Galactorrhea during treatment with trimipramine. A case report. *Pharmacopsychiatry.* 2005;38:326-7. PubMed PMID: 16342006.
7. Venkatesh KK, Castro VM, Perlis RH et al. Impact of antidepressant treatment during pregnancy on obstetric outcomes among women previously treated for depression: An observational cohort study. *J Perinatol.* 2017;37:1003-9. PubMed PMID: 28682318.

- Leggett C, Costi L, Morrison JL et al. Antidepressant use in late gestation and breastfeeding rates at discharge from hospital. *J Hum Lact.* 2017;33:701-9. PubMed PMID: 28984528.

Substance Identification

Substance Name

Trimipramine

CAS Registry Number

739-71-9

Drug Class

Breast Feeding

Lactation

Antidepressive Agents

Antidepressive Agents, Tricyclic