




Trazodone

Revised: October 31, 2018.

CASRN: 19794-93-5



Drug Levels and Effects

Summary of Use during Lactation

Limited information indicates that trazodone levels in milk are low and would not be expected to cause any adverse effects in breastfed infants, especially if the infant is older than 2 months or when doses of 100 mg or less are used at bedtime for sleep.

Drug Levels

Maternal Levels. Six women who were 3 to 8 months postpartum were given 50 mg of trazodone orally and milk samples were taken for 30 hours after administration. Peak trazodone milk levels averaging about 110 mcg/L occurred 2 hours after the dose and milk levels declined with an average half-life of 7.5 hours (range 5.2 to 12.3 hours). The authors estimated that an exclusively breastfed infant would receive a dose of 0.005 mg/kg with this

maternal dose or 0.65% of the maternal weight-adjusted dosage.[1] However, this study did not measure the active metabolite of trazodone in milk.

One woman who was 6.5 weeks postpartum and taking trazodone 75 mg daily had her milk measured at an unreported time after her dose. The trazaodone concentration in her milk was 40 mcg/L.[2]

Infant Levels. Relevant published information was not found as of the revision date.

Effects in Breastfed Infants

One infant whose mother took trazodone 200 mg daily for 12 weeks starting at 4 weeks postpartum was followed up at 12 months of age. No adverse effects on growth and development were found.[3]

One woman was 6.5 weeks postpartum and taking trazodone 75 mg, venlafaxine 75 mg and quetiapine 75 mg daily before conception, during pregnancy and during breastfeeding. Her breastfed infant's development was tested at 12 months of age with the Bayley Scales. Measurements were within normal limits on the mental, psychomotor and behavior scales.[2]

One exclusively breastfed 15-week-old infant was breastfed during maternal therapy with trazodone 100 mg daily and venlafaxine 150 mg daily. No adverse reactions were reported by the mother or found in the medical records.[4]

Effects on Lactation and Breastmilk

A nonpregnant woman with depression was treated with citalopram 20 mg daily, then 40 mg daily. Trazodone 50 mg at bedtime was added to treat insomnia and then increased to 100 mg at bedtime. One week later the patient noticed milk leakage from her breasts, which stained her clothing. Her serum prolactin was somewhat elevated, but no other abnormalities were noted. The trazodone dosage was tapered and then discontinued. One month later, the galactorrhea had resolved and her serum prolactin was in the normal range.[5]

An observational study looked at outcomes of 2859 women who took an antidepressant during the 2 years prior to pregnancy. Compared to women who did not take an antidepressant during pregnancy, mothers who took an antidepressant during all 3 trimesters of pregnancy were 37% less likely to be breastfeeding upon hospital discharge. Mothers who took an antidepressant only during the third trimester were 75% less likely to be breastfeeding at discharge. Those who took an antidepressant only during the first and second trimesters did not have a reduced likelihood of breastfeeding at discharge.[6] The antidepressants used by the mothers were not specified.

A retrospective cohort study of hospital electronic medical records from 2001 to 2008 compared women who had been dispensed an antidepressant during late gestation (n = 575) to those who had a psychiatric illness but did not receive an antidepressant (n = 1552) and mothers who did not have a psychiatric diagnosis (n = 30,535). Women who received an antidepressant were 37% less likely to be breastfeeding at discharge than women without a psychiatric diagnosis, but no less likely to be breastfeeding than untreated mothers with a psychiatric diagnosis.[7] None of the mothers were taking trazodone.

Alternate Drugs to Consider

Nortriptyline, Paroxetine, Sertraline

References

1. Verbeeck RK, Ross SG, McKenna EA. Excretion of trazodone in breast milk. *Br J Clin Pharmacol.* 1986;22:367-70. PubMed PMID: 3768252.

2. Misri S, Corral M, Wardrop AA, Kendrick K. Quetiapine augmentation in lactation: a series of case reports. *J Clin Psychopharmacol*. 2006;26:508-11. PubMed PMID: 16974194.
3. Misri S, Sivertz K. Tricyclic drugs in pregnancy and lactation: a preliminary report. *Int J Psychiatry Med*. 1991;21:157-71. PubMed PMID: 1894455.
4. Newport DJ, Ritchie JC, Knight B et al. Venlafaxine in human breast milk and nursing infant plasma: determination of exposure. *J Clin Psychiatry*. 2009;70:1304-10. PubMed PMID: 19607765.
5. Arslan FC, Uysal EK, Ozkorumak E et al. Trazodone induced galactorrhea: A case report. *Gen Hosp Psychiatry*. 2015;37:373.e1-2. PubMed PMID: 25920951.
6. Venkatesh KK, Castro VM, Perlis RH et al. Impact of antidepressant treatment during pregnancy on obstetric outcomes among women previously treated for depression: An observational cohort study. *J Perinatol*. 2017;37:1003-9. PubMed PMID: 28682318.
7. Leggett C, Costi L, Morrison JL et al. Antidepressant use in late gestation and breastfeeding rates at discharge from hospital. *J Hum Lact*. 2017;33:701-9. PubMed PMID: 28984528.

Substance Identification

Substance Name

Trazodone

CAS Registry Number

19794-93-5

Drug Class

Breast Feeding

Lactation

Antidepressive Agents

Serotonin Uptake Inhibitors