




## Triazolam

Revised: October 31, 2018.

CASRN: 28911-01-5



## Drug Levels and Effects

### Summary of Use during Lactation

Because little information is available on the use of triazolam during breastfeeding, an alternate hypnotic may be preferred, especially while nursing a newborn or preterm infant. Triazolam has a relatively short half-life, so occasional use while breastfeeding an older infant should pose little risk to the infant, but monitor the infant for excessive drowsiness.

### Drug Levels

*Maternal Levels.* Relevant published information was not found as of the revision date.

*Infant Levels.* Relevant published information was not found as of the revision date.

## Effects in Breastfed Infants

In a telephone follow-up study, 124 mothers who took a benzodiazepine while nursing reported whether their infants had any signs of sedation. One mother took triazolam while breastfeeding and reported no sedation in her infant.[1]

## Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

## Alternate Drugs to Consider

Zaleplon, Zolpidem

## References

1. Kelly LE, Poon S, Madadi P, Koren G. Neonatal benzodiazepines exposure during breastfeeding. J Pediatr. 2012;161:448-51. PubMed PMID: 22504099.

## Substance Identification

### Substance Name

Triazolam

### CAS Registry Number

28911-01-5

### Drug Class

Breast Feeding

Lactation

Hypnotics and Sedatives