




## Alclometasone

Revised: October 31, 2018.

CASRN: 67452-97-5



## Drug Levels and Effects

### Summary of Use during Lactation

Alclometasone has not been studied during breastfeeding. Since only extensive application of the most potent of these drugs cause systemic effects in the mother, it is unlikely that short-term application of topical corticosteroids would pose a risk to the breastfed infant by passage into breastmilk. However, it would be prudent to use the least potent drug on the smallest area of skin possible. It is particularly important to ensure that the infant's skin does not come into direct contact with the areas of skin that have been treated. Only the lower potency corticosteroids (e.g., hydrocortisone, triamcinolone) should be used on the nipple or areola where the infant could directly ingest the drugs from the skin. Only water-miscible cream or gel products should be applied to the breast because ointments may expose the infant to high levels of mineral paraffins via licking.[1] Any topical corticosteroid should be wiped off thoroughly prior to nursing if it is being applied to the breast or nipple area.

**Disclaimer:** Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site .

## Drug Levels

*Maternal Levels.* Relevant published information was not found as of the revision date.

*Infant Levels.* Relevant published information was not found as of the revision date.

## Effects in Breastfed Infants

Topical application of a corticosteroid with relatively high mineralocorticoid activity (isofluprednone acetate) to the mother's nipples resulted in prolonged QT interval, cushingoid appearance, severe hypertension, decreased growth and electrolyte abnormalities in her 2-month-old breastfed infant. The mother had used the cream since birth for painful nipples.[2]

## Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

## Alternate Drugs to Consider

(Topical) [Hydrocortisone](#), Topical

## References

1. Greenberger PA, Patterson R. The management of asthma during pregnancy and lactation. *Clin Rev Allergy.* 1987;5:317-24. PubMed PMID: 3319123.
2. Ellsworth A. Pharmacotherapy of asthma while breastfeeding. *J Hum Lact.* 1994;10:39-41. PubMed PMID: 7619245.
3. National Heart, Lung, and Blood Institute, National Asthma Education and Prevention Program Asthma and Pregnancy Working Group. NAEPP expert panel report. Managing asthma during pregnancy: recommendations for pharmacologic treatment-2004 update. 2004;1-57. Available at: <http://www.nhlbi.nih.gov/health/prof/lung/asthma/astpreg.htm>
4. Noti A, Grob K, Biedermann M et al. Exposure of babies to C(15)-C(45) mineral paraffins from human milk and breast salves. *Regul Toxicol Pharmacol.* 2003;38(3):317-25. PubMed PMID: 14623482.
5. De Stefano B, Bongo IG, Borgna-Pignatti C et al. Factitious hypertension with mineralocorticoid excess in an infant. *Helv Paediatr Acta.* 1983;38:185-9. PubMed PMID: 6874387.

## Substance Identification

### Substance Name

Alclometasone

### CAS Registry Number

67452-97-5

### Drug Class

Breast Feeding

Lactation

Corticosteroids, Topical

Glucocorticoids

## Anti-Inflammatory Agents