

NLM Citation: Drugs and Lactation Database (LactMed) [Internet]. Bethesda (MD): National Library of Medicine (US); 2006-. Psyllium.

[Updated 2019 Jan 7].

Bookshelf URL: https://www.ncbi.nlm.nih.gov/books/



Psyllium

Revised: January 7, 2019.

CASRN: 8063-16-9

Drug Levels and Effects

Summary of Use during Lactation

Few data are available on psyllium use during breastfeeding. However, the drug is not absorbed from the gastrointestinal tract, so it cannot enter the breastmilk. Psyllium is acceptable to use during breastfeeding.[1][2]

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Effects in Breastfed Infants

Twenty postpartum mothers were given a laxative containing 2.7 grams of plantango (psyllium) and senna equivalent to 15 mg of sennosides a and b daily on days 2 to 4 postpartum. Of the 11 infants who were breastfed (extent notstated), none had any loose stools.[3]

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Alternate Drugs to Consider

(Constipation) Bisacodyl, Docusate, Magnesium Hydroxide, Sodium Phosphate; (Irritable Bowel Syndrome) Linaclotide, Lubiprostone, Plecanatide, Prucalopride

References

- 1. Lewis JH, Weingold AB and the Committee of FDA-Related Matters, American College of Gastroenterology. The use of gastrointestinal drugs during pregnancy and lactation. Am J Gastroenterol. 1985;80:912-23. PubMed PMID: 2864852.
- 2. Hagemann TM. Gastrointestinal medications and breastfeeding. J Hum Lact. 1998;14:259-62. PubMed PMID: 10205441.

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site .

3. Faber P, Strenge-Hesse A. Relevance of rhein excretion into breast milk. Pharmacology. 1988;36 (Suppl 1):212-20. PubMed PMID: 336852.

Substance Identification

Substance Name

Psyllium

CAS Registry Number

8063-16-9

Drug Class

Breast Feeding

Lactation

Cathartics

Gastrointestinal Agents

Laxatives