




Sucralfate

Revised: October 31, 2018.

CASRN: 54182-58-0



Drug Levels and Effects

Summary of Use during Lactation

No data are available on sucralfate use during breastfeeding; however, it is virtually unabsorbed orally. Most authorities consider sucralfate acceptable to use during breastfeeding.[1][2][3][4] No special precautions are required.

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Alternate Drugs to Consider

Antacids, Cimetidine, Famotidine, Nizatidine, Omeprazole, Pantoprazole, Ranitidine

References

1. Lewis JH, Weingold AB and the Committee of FDA-Related Matters, American College of Gastroenterology. The use of gastrointestinal drugs during pregnancy and lactation. *Am J Gastroenterol.* 1985;80:912-23. PubMed PMID: 2864852.
2. Hagemann TM. Gastrointestinal medications and breastfeeding. *J Hum Lact.* 1998;14:259-62. PubMed PMID: 10205441.
3. Broussard CN, Richter JE. Treating gastro-oesophageal reflux disease during pregnancy and lactation: what are the safest therapy options? *Drug Saf.* 1998;19:325-37. PubMed PMID: 9804446.
4. Richter JE. Review article: the management of heartburn in pregnancy. *Aliment Pharmacol Ther.* 2005;22:749-57. PubMed PMID: 16225482.

Substance Identification

Substance Name

Sucralfate

CAS Registry Number

54182-58-0

Drug Class

Breast Feeding

Lactation

Anti-Ulcer Agents

Gastrointestinal Agents