

**NLM Citation:** Drugs and Lactation Database (LactMed) [Internet]. Bethesda (MD): National Library of Medicine (US); 2006-. Botulin B.

[Updated 2019 Jun 30].

Bookshelf URL: https://www.ncbi.nlm.nih.gov/books/



# **Botulin B**

Revised: June 30, 2019.

CASRN: 93384-44-2

## **Drug Levels and Effects**

## **Summary of Use during Lactation**

No data exist on the medical use of botulin B (botulinum toxin, rimabotulinumtoxinB) during breastfeeding. However, rimabotulinumtoxinB is not detectable systemically after intramuscular use, thus excretion into breast milk is considered unlikely. One infant was safely breastfed during maternal botulism and no botulinum toxin was detectable in the mother's milk or infant's blood or stools. Since the doses used medically are far lower than those that cause botulism, amounts ingested by the infant, if any, are expected to be small and not cause any adverse effects in breastfed infants.[1] No special precautions are required.

### **Drug Levels**

*Maternal Levels.* Published information on the medical use of botulin A during breastfeeding was not found as of the revision date.

Type A botulinum toxin was detected in the blood and stools of a nursing mother after ingesting fermented salmon eggs. She was given 2 vials of trivalent botulism antitoxin, 1 intravenously and 1 intramuscularly. A milk sample obtained 3 days after the onset of her illness and 4 hours after administration of botulinum antitoxin had no detectable botulinum toxin nor botulism organisms.[2]

*Infant Levels*. Type A botulinum toxin was detected in the blood and stools of a nursing mother after ingesting fermented salmon eggs. No botulinum toxin was detected in the breastfed infant's blood or stool on the day the mother was admitted to the hospital (3 days after the onset of illness) and no botulism organisms were detected in the infant's stools.[2]

### **Effects in Breastfed Infants**

Published information on the medical use of botulin B during breastfeeding was not found as of the revision date.

**Disclaimer:** Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site .

A woman developed botulism after ingesting fermented salmon eggs while breastfeeding her 8-month-old breastfed (extent not stated) infant. The infant developed no signs or symptoms of botulism even though she continued to nurse throughout the mother's hospitalization.[2]

#### **Effects on Lactation and Breastmilk**

Relevant published information was not found as of the revision date.

#### References

- 1. Lee KC, Korgavkar K, Dufresne RGJ et al. Safety of cosmetic dermatologic procedures during pregnancy. Dermatol Surg. 2013;39:1573-86. PubMed PMID: 24164677.
- 2. Middaugh J. Botulism and breast milk. N Engl J Med. 1978;298:343. PubMed PMID: 622098.

## **Substance Identification**

#### **Substance Name**

Botulin B

## **CAS Registry Number**

93384-44-2

## **Drug Class**

**Breast Feeding** 

Lactation

**Bacterial Toxins** 

Neuromuscular Agents

**Neurotoxins**