




Benzoyl Peroxide

Revised: October 31, 2018.

CASRN: 94-36-0



Drug Levels and Effects

Summary of Use during Lactation

Topical benzoyl peroxide has not been studied during breastfeeding. Because only about 5% is absorbed following topical application, it is considered a low risk to the nursing infant.[1][2] Ensure that the infant's skin does not come into direct contact with the areas of skin that have been treated. Only water-miscible cream or gel products should be applied to the breast because ointments may expose the infant to high levels of mineral paraffins via licking.[3]

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Alternate Drugs to Consider

(Acne) Azelaic Acid, Clindamycin, Erythromycin, Tretinoin

References

1. Leachman SA, Reed BR. The use of dermatologic drugs in pregnancy and lactation. *Dermatol Clin.* 2006;24:167-97. PubMed PMID: 16677965.
2. Zip C. Common sense dermatological drug suggestions for women who are breast-feeding. *Skin Therapy Lett.* 2002;7:5-7. PubMed PMID: 12007012.
3. Noti A, Grob K, Biedermann M et al. Exposure of babies to C(15)-C(45) mineral paraffins from human milk and breast salves. *Regul Toxicol Pharmacol.* 2003;38:317-25. PubMed PMID: 14623482.

Substance Identification

Substance Name

Benzoyl Peroxide

CAS Registry Number

94-36-0

Drug Class

Breast Feeding

Lactation

Dermatologic Agents

Peroxides