




## Butenafine

Revised: October 31, 2018.

CASRN: 101828-21-1



## Drug Levels and Effects

### Summary of Use during Lactation

Topical butenafine has not been studied during breastfeeding. Because it is poorly absorbed after topical application, it is not likely to reach the bloodstream of the infant or cause any adverse effects in breastfed infants. Ensure that the infant's skin does not come into direct contact with the areas of skin that have been treated. Only water-miscible cream or gel products should be applied to the breast because ointments may expose the infant to high levels of mineral paraffins via licking.[1]

### Drug Levels

*Maternal Levels.* Relevant published information was not found as of the revision date.

*Infant Levels.* Relevant published information was not found as of the revision date.

## Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

## Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

## Alternate Drugs to Consider

Clotrimazole, Miconazole, Terbinafine

## References

1. Noti A, Grob K, Biedermann M et al. Exposure of babies to C(15)-C(45) mineral paraffins from human milk and breast salves. Regul Toxicol Pharmacol. 2003;38:317-25. PubMed PMID: 14623482.

## Substance Identification

### Substance Name

Butenafine

### CAS Registry Number

101828-21-1

### Drug Class

Breast Feeding

Lactation

Anti-Infective Agents

Antifungal Agents