




Glycopyrrolate

Revised: December 3, 2018.

CASRN: 596-51-0



Drug Levels and Effects

Summary of Use during Lactation

No information is available on the use of glycopyrrolate during breastfeeding. Because glycopyrrolate is a quaternary ammonium compound, it is not likely to be absorbed and reach the bloodstream of the infant,[1][2][3] especially when given by inhalation or topically on the skin. Long-term oral use of glycopyrrolate might reduce milk production or milk letdown, but a single dose is unlikely to interfere with breastfeeding. During long-term use, observe for signs of decreased lactation (e.g., insatiety, poor weight gain).

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site .

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information in nursing mothers was not found as of the revision date. Anticholinergics can inhibit lactation in animals, apparently by inhibiting growth hormone and oxytocin secretion.[4][5][6][7][8] Anticholinergic drugs can also reduce serum prolactin in nonnursing women.[9] The prolactin level in a mother with established lactation may not affect her ability to breastfeed.

References

1. Hale TW. Anesthetic medications in breastfeeding mothers. *J Hum Lact.* 1999;15:185-94. PubMed PMID: 10578796.
2. Lee JJ, Rubin AP. Breast feeding and anaesthesia. *Anaesthesia.* 1993;48:616-25. PubMed PMID: 8346780.
3. Dalal PG, Bosak J, Berlin C. Safety of the breast-feeding infant after maternal anesthesia. *Paediatr Anaesth.* 2014;24:359-71. PubMed PMID: 24372776.
4. Aaron DK, Ely DG, Deweese WP et al. Reducing milk production in ewes at weaning using restricted feeding and methscopolamine bromide. *J Anim Sci.* 1997;75:1434-42. PubMed PMID: 9250502.
5. Powell MR, Keisler DH. A potential strategy for decreasing milk production in the ewe at weaning using a growth hormone release blocker. *J Anim Sci.* 1995;73:1901-5. PubMed PMID: 7592071.
6. Daniel JA, Thomas MG, Powell MR, Keisler DH. Methscopolamine bromide blocks hypothalamic-stimulated release of growth hormone in ewes. *J Anim Sci.* 1997;75:1359-62. PubMed PMID: 9159285.
7. Bizzarro A, Iannucci F, Tolino A et al. Inhibiting effect of atropine on prolactin blood levels after stimulation with TRH. *Clin Exp Obstet Gynecol.* 1980;7:108-11. PubMed PMID: 6788407.
8. Svennersten K, Nelson L, Juvnas-Moberg K. Atropinization decreases oxytocin secretion in dairy cows. *Acta Physiol Scand.* 1992;145:193-4. PubMed PMID: 1636447.
9. Masala A, Alagna S, Devilla L et al. Muscarinic receptor blockade by pirenzepine: effect on prolactin secretion in man. *J Endocrinol Invest.* 1982;5:53-5. PubMed PMID: 6808052.

Substance Identification

Substance Name

Glycopyrrolate

CAS Registry Number

596-51-0

Drug Class

Breast Feeding

Lactation

Adjuvants, Anesthesia

Bronchodilator Agents

Muscarinic Antagonists

Parasympatholytics