




Raltegravir

Revised: October 31, 2018.

CASRN: 518048-05-0



Drug Levels and Effects

Summary of Use during Lactation

In the United States and other developed countries, HIV-infected mothers should generally not breastfeed their infants. No published information is available on the use of raltegravir during breastfeeding. In countries in which no acceptable, feasible, sustainable and safe replacement feeding is available, World Health Organization guidelines recommend that all women with an HIV infection who are pregnant or breastfeeding should be maintained on antiretroviral therapy for at least the duration of risk for mother-to-child transmission. Mothers should exclusively breastfeed their infants for the first 6 months of life; breastfeeding with complementary feeding should continue through 12 months of life. The first choice regimen for nursing mothers is tenofovir, efavirenz and either lamivudine or emtricitabine. If these drugs are unavailable, alternative regimens include: 1) zidovudine, lamivudine and efavirenz; 2) zidovudine, lamivudine and nevirapine; or 3) tenofovir, nevirapine and

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site .

either lamivudine or emtricitabine. Exclusively breastfed infants should also receive 6 weeks of prophylaxis with nevirapine.[1][2]

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date. *Infant Levels.* Relevant published information was not found as of the revision date.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Alternate Drugs to Consider

Lamivudine, Nelfinavir, Nevirapine, Zidovudine

References

1. World Health Organization. HIV and infant feeding: update. 2007. Available at: http://whqlibdoc.who.int/publications/2007/9789241595964_eng.pdf
2. World Health Organization. Consolidated guidelines on the use of antiretroviral drugs for treating and preventing HIV infection. Geneva: World Health Organization. 2013. Available at: <http://www.who.int/hiv/pub/guidelines/arv2013/en/>

Substance Identification

Substance Name

Raltegravir

CAS Registry Number

518048-05-0

Drug Class

Breast Feeding

Lactation

Anti-Infective Agents

Anti-HIV Agents

Antiviral Agents

Anti-Retroviral Agents

HIV Integrase Inhibitors