

NLM Citation: Drugs and Lactation Database (LactMed) [Internet]. Bethesda (MD): National Library of Medicine (US); 2006-. Alendronate.

[Updated 2018 Dec 3].


Bookshelf URL: https://www.ncbi.nlm.nih.gov/books/



Alendronate

Revised: December 3, 2018.

CASRN: 66376-36-1



Drug Levels and Effects

Summary of Use during Lactation

Limited evidence indicates that breastfeeding after cessation of long-term bisphosphonate treatment appears to have no adverse effects on the infant. Because no information is available on the use of alendronate during breastfeeding, an alternate drug may be preferred, especially while nursing a newborn or preterm infant. However, absorption of alendronate by a breastfed infant is unlikely. If the mother receives a bisphosphonate during pregnancy or nursing, some experts recommend monitoring the infant's serum calcium during the first 2 months postpartum.[1]

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site .

Drug Levels

Alendronate is poorly absorbed orally (average in adults 1% on an empty stomach, negligible with food and calcium), so absorption of alendronate by a breastfed infant is unlikely. *Maternal Levels*. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Effects in Breastfed Infants

Because alendronate can persist in the body for years after long-term administration, the following cases may be relevant. A woman received alendronate for 6 months, then pamidronate every 4 months for 1 year prior to conception. Her infant was breastfed (extent not stated) for 3 months. The infant had mild hypocalcemia at 2 months of age, but a normal calcium level and normal long bone development at 5 months of age.[2]

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Alternate Drugs to Consider

Pamidronate

References

- 1. Stathopoulos IP, Liakou CG, Katsalira A et al. The use of bisphosphonates in women prior to or during pregnancy and lactation. Hormones. 2011;10:280-91. PubMed PMID: 22281884.
- 2. Hassen-Zrour S, Korbaa W, Bejia I et al. Maternal and fetal outcome after long-term bisphosphonate exposure before conception. Osteoporos Int. 2010;21:709-10. PubMed PMID: 19533208.

Substance Identification

Substance Name

Alendronate

CAS Registry Number

66376-36-1

Drug Class

Breast Feeding

Lactation

Bisphosphonates

Bone Density Conservation Agents

Diphosphonates