

U.S. National Library of Medicine National Center for Biotechnology Information **NLM Citation:** Drugs and Lactation Database (LactMed) [Internet]. Bethesda (MD): National Library of Medicine (US); 2006-. Measles-Mumps-Rubella-Varicella Vaccine. [Updated 2018 Dec 3]. **Bookshelf URL:** https://www.ncbi.nlm.nih.gov/books/



Measles-Mumps-Rubella-Varicella Vaccine

Revised: December 3, 2018.

Drug Levels and Effects

Summary of Use during Lactation

The Centers for Disease Control and Prevention and several health professional organizations state that vaccines given to a nursing mother do not affect the safety of breastfeeding for mothers or infants and that breastfeeding is not a contraindication to measles, mumps, rubella and varicella virus vaccine. Breastfed infants should be vaccinated according to the routine recommended schedules. Although rubella vaccine virus might be excreted into milk, the virus usually does not infect the infant. If an infection does occur, it is well tolerated because the viruses are attenuated.[1][2][3] No clear evidence exists of live attenuated measles or mumps vaccine virus excretion into breastmilk.[4]

Drug Levels

Maternal Levels. No studies have evaluated the effects of the combined measles, mumps, rubella and varicella (MMRV) vaccine during breastfeeding. A study of mothers vaccinated with the Cendehill strain of live, attenuated rubella virus found no transmission of the live virus to their breastfed infants.[5] However, rubella vaccine virus can appear in breastmilk and result in infections in some infants.[6][7][8][9] *See* "Reported Side Effects In Breastfed Infants" below.

In a prospective study, 169 mothers with low titers or either measles or rubella were given MR vaccine (Schwarz FF-8 strain/TO-336 strain; Takeda Pharmaceutical Co. Ltd, Osaka, Japan) at their 1-month postnatal checkup. Mothers provided 5 mL of breastmilk 2 weeks later. Rubella virus antigen was not detected in the breastmilk of any mothers. Measles virus RNA was isolated from the breastmilk of 2 vaccinated mothers.[10]

Infant Levels. Relevant published information was not found as of the revision date. Two postpartum women immunized with varicella vaccine provided preimmunization and serial postimmunization breastmilk samples (number not specified). One sample of colostrum contained detectable viral DNA and B-actin by polymerase chain reaction, but all other samples were negative for these species. No varicella gene sequences were found in any of the samples.[11]

One study of 12 women vaccinated with live, attenuated varicella vaccine found no evidence of varicella virus excretion into breastmilk.[12]

Infant Levels. Relevant published information was not found as of the revision date.

Disclaimer: Information presented in this database is not meant as a substitute for professional judgment. You should consult your healthcare provider for breastfeeding advice related to your particular situation. The U.S. government does not warrant or assume any liability or responsibility for the accuracy or completeness of the information on this Site .

Effects in Breastfed Infants

Limited data indicate that breastfeeding can enhance the response of the infant to certain vaccine antigens.[2][3] [13]

One 12-day-old breastfed infant developed a rubella infection 11 days after maternal vaccination with live rubella vaccine.[14] However, it is questionable if maternal vaccination was the cause of the infant's infection. [15]

Another breastfed infant had live rubella vaccine virus isolated from a throat swab after maternal immunization. The infant did not demonstrate seroconversion or adverse reactions.[6]

Some breastfed infants acquire passive immunity to rubella after maternal vaccination as do infants of mothers with natural rubella immunity. However, neither group of infants had a decreased response to rubella vaccine administered to the infant at 15 to 18 months of age.[7]

After immunization of their mothers with rubella vaccine, 25% of breastfed infants in one study showed transient seroconversion to rubella virus but without any clinical disease.[8]

In a prospective study of mothers given MR vaccine (Schwarz FF-8 strain/TO-336 strain; Takeda Pharmaceutical Co. Ltd, Osaka, Japan) at their 1-month postnatal checkup, measles virus RNA was isolated from the breastmilk of 2 vaccinated mothers. Neither of their breastfed infants had any clinical disease.[10]

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

References

- 1. Gruslin A, Steben M, Halperin S et al. Immunization in pregnancy: No. 220, December 2008. Int J Gynaecol Obstet. 2009;105:187-91. PubMed PMID: 19367691.
- 2. American Academy of Pediatrics Committee on Infectious Diseases, Kimberlin DW, Brady MT et al. Red Book: 2015 Report of the Committee on Infectious Diseases. 30th ed. Elk Grove Village, IL: American Academy of Pediatrics. 2015.
- 3. General recommendations on immunization --- recommendations of the Advisory Committee on Immunization Practices (ACIP). MMWR Recomm Rep. 2011;60 (RR-2):1-64. PubMed PMID: 21293327.
- 4. Alain S, Dommergues MA, Jacquard AC et al. State of the art: Could nursing mothers be vaccinated with attenuated live virus vaccine? Vaccine. 2012;30:4921-6. PubMed PMID: 22659446.
- 5. Farquhar JD. Follow-up on rubella vaccinations and experience with subclinical reinfection. J Pediatr. 1972;81:460-5. PubMed PMID: 5065683.
- 6. Buimovici-Klein E, Hite RL, Byrne T et al. Isolation of rubella virus in milk after pospartum immunization. J Pediatr. 1977;91:939-41. PubMed PMID: 925824.
- 7. Krogh V, Duffy LC, Wong D et al. Postpartum immunization with rubella virus vaccine and antibody response in breast-feeding infants. J Lab Clin Med. 1989;113(6):695-9. PubMed PMID: 2732617.
- 8. Losonsky GA, Fishaut JM, Strussenberg J. Effect of immunization against rubella on lactation products. II. Maternal-neonatal interactions. J Infect Dis. 1982;145:661-6. PubMed PMID: 7077090.
- 9. Losonsky GA, Fishaut JM, Strussenberg J, Ogra PL. Effect of immunization against rubella on lactation products. I. Development and characterization of specific immunologic reactivity in breast milk. J Infect Dis. 1982;145:654-60. PubMed PMID: 7077089.
- 10. Hisano M, Kato T, Inoue E et al. Evaluation of measles-rubella vaccination for mothers in early puerperal phase. Vaccine. 2016;34:1208-14. PubMed PMID: 26801065.
- 11. Dolbear GL, Moffat J, Falkner C, Wojtowycz M. A pilot study: Is attenuated varicella virus present in breast milk after postpartum immunization? Obstet Gynecol. 2003;101 (4 Suppl):47s. Abstract.

- 12. Bohlke K, Galil K, Jackson LA et al. Postpartum varicella vaccination: is the vaccine virus excreted in breast milk? Obstet Gynecol. 2003;102:970-7. PubMed PMID: 14672472.
- 13. Pabst HF. Immunomodulation by breast-feeding. Pediatr Infect Dis J. 1997;16:991-5. PubMed PMID: 9380478.
- 14. Landes RD, Bass JW, Millunchick EW et al. Neonatal rubella following postpartum maternal immunization. J Pediatr. 1980;97:465-7. PubMed PMID: 7191002.
- 15. Lerman SJ. Neonatal rubella following maternal immunization. J Pediatr. 1981;98:668-9. Letter. PubMed PMID: 7205504.

Substance Identification

Substance Name

Measles-Mumps-Rubella-Varicella Vaccine

Drug Class

Breast Feeding

Lactation

Vaccines