



Milk Thistle

Revised: May 1, 2019.

CASRN: 84604-20-6

Drug Levels and Effects

Summary of Use during Lactation

Milk thistle (*Silybum marianum*) contains silymarin which is a mixture of flavonolignans, mainly silibinin (also known as silybin), as well as silycristine, silydianin, quercetin and taxifolin.[1] Silymarin is a standardized preparation extracted from the fruits (seeds) of milk thistle. Milk thistle is a purported galactagogue,[2] and is included in some proprietary mixtures promoted to increase milk supply; however, no scientifically valid clinical trials support this use.[3][4][5] Although a study on the high potency purified milk thistle component, silymarin, and a phosphatidyl conjugate of silymarin indicated some galactagogue activity, this does not necessarily imply activity of milk thistle itself. Galactagogues should never replace evaluation and counseling on modifiable factors that affect milk production.[6]

Limited data indicate that the silymarin components are not excreted into breastmilk in measurable quantities. Additionally, because silymarin components are poorly absorbed orally, milk thistle is unlikely to adversely affect the breastfed infant. Milk thistle and silymarin are generally well tolerated in adults with only mild side effects such as diarrhea, headache, and skin reactions. It might increase the metabolism of some drugs. Rarely, severe allergies and anaphylaxis are reported. Avoid in patients with known allergy to members of the aster (Compositae or Asteraceae) family, such as daisies, artichokes, common thistle, and kiwi because cross-allergenicity is possible.

Dietary supplements do not require extensive pre-marketing approval from the U.S. Food and Drug Administration. Manufacturers are responsible to ensure the safety, but do not need to *prove* the safety and effectiveness of dietary supplements before they are marketed. Dietary supplements may contain multiple ingredients, and differences are often found between labeled and actual ingredients or their amounts. A manufacturer may contract with an independent organization to verify the quality of a product or its ingredients, but that does *not* certify the safety or effectiveness of a product. Because of the above issues, clinical testing results on one product may not be applicable to other products. More detailed information [about dietary supplements](#) is available elsewhere on the LactMed Web site.

Drug Levels

Silibinin has low and variable bioavailability of 23 to 47% after oral administration and is rapidly converted into glucuronide conjugates which have longer half-lives in plasma than the unconjugated forms.[7] *Maternal Levels.* Five mothers who had decided to stop nursing because their infants were 9 months old were given 600 mg of micronized silymarin (BIO-C brand) orally 3 times daily. After 5 days of therapy, milk was collected at unspecified times for analysis by HPLC. Silymarin flavonolignans were undetectable (<1 mcg/L) in milk samples; however, their conjugated forms might not have been detected by the assay method used.[8]

Silymarin metabolites (silychristin, silydianin, silibyn A, silibyn B, isosilybin A and isosilybin B) were not detected (<0.3 mg/kg) in breastmilk of any of 25 women studied on day 14 of treatment with micronized silymarin 252 mg (BIO-C) twice daily.[9]

Infant Levels. Relevant published information was not found as of the revision date.

Effects in Breastfed Infants

In a study of galactagogue containing 5 grams of a mixture of silymarin-phosphatidylserine and galega (goat's rue) in an unspecified proportion and from an unspecified source, none of the typical adverse effects of silymarin were noted in the breastfed infants.[10]

A study compared a commercial product containing silymarin 252 mg (BIO-C) to placebo every 12 hours in mothers of preterm (<32 weeks) infants. No adverse effects were observed in any of the infants.[9]

Effects on Lactation and Breastmilk

No human data are available on the effect of milk thistle or its components on serum prolactin. A study in gilts (female domestic pigs) found that silymarin 4 grams twice daily during pregnancy and lactation found that serum prolactin levels were increased compared to gilts given placebo. However, the slight increase in prolactin had no effect on mammary gland development, nor on plasma progesterone or estradiol.[11]

A study was performed on 50 medically normal postpartum mothers with milk production judged to be less than normal for patients in the hospital in Lima, Peru where the study was conducted. Mothers were divided non-randomly into 2 groups of 25 women who had identical ages, weights, number of children and newborn's age, although ages were not reported. The group that was given micronized silymarin (BIO-C brand) 420 mg daily for 63 days had a baseline milk production of 602 mL daily. The milk volumes and composition (water, fats, carbohydrate and protein) of the 2 groups were not significantly different on day 0. The group given an identical placebo had a baseline milk production of 530 mL daily. Milk production was measured on day 30 and day 63 by infant weighings before and after nursing followed by emptying the breasts with a breast pump. The composition of the milk was also determined. Statistically significant differences in average milk production were found on day 30 (990 grams in the silymarin group and 650 grams in the placebo group) and on day 63 (1119 grams in the silymarin group and 701 grams in the placebo group). Milk composition was not different between the groups at the two time points.[8] Deficiencies in this study include the lack of randomization, no investigator blinding, and no optimization of breastfeeding technique prior to study enrollment. Also, breastfeeding duration and long-term infant growth were not studied.

In a randomized, double blinded study, a placebo or galactagogue containing 5 grams of a mixture of silymarin-phosphatidylserine and galega (goat's rue) in a commercial product (Piulatte Plus, Milte) was given once daily to mothers of preterm infants. Phosphatidyl serine purportedly has improved bioavailability over silymarin alone. The placebo group received 5 grams of lactose once daily. The medication or placebo was given from day 3 to day 28 postpartum. Mothers pumped using a breast pump every 2 to 3 hours during the day and as desired at night. Milk production was measured on days 7, 14 and 28 postpartum. Daily milk production averaged 200 mL

in the treated group and 115 mL in the control group. The total amount of milk produced during the study period and the proportion of women producing more than 200 mL daily was greater in the treated group than controls on days 7 and 28.[10] Mothers were contacted at 3 and 6 months postpartum concerning breastmilk production. Of the 89 mothers who responded satisfactorily at 3 months, more mothers who had received silymarin-galega were exclusively breastfeeding than those who received placebo (22/50 vs 12/50). Also, more mothers were feeding more than 50% breastmilk to their infants in the treatment group than the placebo group (29/50 vs 18/50). At 6 months postpartum, more mothers were feeding more than 50% breastmilk to their infants in the treatment group than the placebo group (22/50 vs 12/50). These differences were statistically significant.[12]

A randomized study compared a commercial product containing silymarin 252 mg (BIO-C) to placebo every 12 hours in mothers of preterm (<32 weeks) infants, beginning at 10 days postpartum. Mothers used a breast pump 6 times daily and measured milk output before beginning, 5 times during the 28 days of treatment, and on days 36 and 45. No difference in milk production was observed between the two groups at any time point. The mothers' guesses of whether they had taken placebo or silymarin were no better than chance.[9]

In a survey of 188 nursing women from 27 states (52% from Louisiana), 24 had used fennel as a galactagogue. Of those who used it, 52% were not sure that it increased their milk supply and 4 reported (unspecified) side effects. [13]

References

1. Dietz BM, Hajirahimkhan A, Dunlap TL et al. Botanicals and their bioactive phytochemicals for women's health. *Pharmacol Rev.* 2016;68:1026-73. PubMed PMID: 27677719.
2. Jackson PC. Complementary and alternative methods of increasing breast milk supply for lactating mothers of infants in the NICU. *Neonatal Netw.* 2010;29:225-30. PubMed PMID: 20630837.
3. Mortel M, Mehta SD. Systematic review of the efficacy of herbal galactagogues. *J Hum Lact.* 2013;29:154-62. PubMed PMID: 23468043.
4. Zapantis A, Steinberg JG, Schilit L. Use of herbals as galactagogues. *J Pharm Pract.* 2012;25:222-31. PubMed PMID: 22392841.
5. Forinash AB, Yancey AM, Barnes KN, Myles TD. The use of galactagogues in the breastfeeding mother. *Ann Pharmacother.* 2012;46:1392-404. PubMed PMID: 23012383.
6. Brodribb W. ABM Clinical Protocol #9: Use of galactagogues in initiating or augmenting maternal milk production, second revision 2018. *Breastfeed Med.* 2018;13:307-14. PubMed PMID: 29902083.
7. He SM, Li CG, Liu JP et al. Disposition pathways and pharmacokinetics of herbal medicines in humans. *Curr Med Chem.* 2010;17:4072-113. PubMed PMID: 20939821.
8. Di Pierro F, Callegari A, Carotenuto D, Tapia MM. Clinical efficacy, safety and tolerability of BIO-C (micronized silymarin) as a galactagogue. *Acta Biomed.* 2008;79:205-10. PubMed PMID: 19260380.
9. Peila C, Coscia A, Tonetto P et al. Evaluation of the galactagogue effect of silymarin on mothers of preterm newborns (<32 weeks). *Pediatr Med Chir.* 2015;37:13-7. PubMed PMID: 26714778.
10. Zecca E, Zuppa AA, D'Antuono A et al. Efficacy of a galactagogue containing silymarin-phosphatidylserine and galega in mothers of preterm infants: A randomized controlled trial. *Eur J Clin Nutr.* 2016;70:1151-4. PubMed PMID: 27245206.
11. Farmer C, Lapointe J, Palin MF. Effects of the plant extract silymarin on prolactin concentrations, mammary gland development, and oxidative stress in gestating gilts. *J Anim Sci.* 2014;92:2922-30. PubMed PMID: 24504042.
12. Serrao F, Corsello M, Romagnoli C et al. The long-term efficacy of a galactagogue containing silymarin-phosphatidylserine and Galega on milk production of mothers of preterm infants. *Breastfeed Med.* 2018;13:67-9. PubMed PMID: 29148822.

13. Bazzano AN, Cenac L, Brandt AJ et al. Maternal experiences with and sources of information on galactagogues to support lactation: a cross-sectional study. *Int J Womens Health*. 2017;9:105-13. PubMed PMID: 28280392.

Substance Identification

Substance Name

Milk Thistle

Scientific Name

Silybum marianum

CAS Registry Number

84604-20-6

Drug Class

Breast Feeding

Lactation

Antioxidants

Complementary Therapies

Galactagogues

Phytotherapy

Plants, Medicinal

Protective Agents