




Indacaterol

Revised: December 3, 2018.

CASRN: 312753-06-3



Drug Levels and Effects

Summary of Use during Lactation

Although no published data exist on the use of indacaterol by mouth or inhaler during lactation, data from the related drug, terbutaline, indicate that very little is expected to be excreted into breastmilk.[1] The authors of several reviews agree that use of inhaled bronchodilators is acceptable during breastfeeding because of the low bioavailability and maternal serum levels after use.[2][3][4][5]

Drug Levels

Maternal Levels. Relevant published information was not found as of the revision date.

Infant Levels. Relevant published information was not found as of the revision date.

Effects in Breastfed Infants

Relevant published information was not found as of the revision date.

Effects on Lactation and Breastmilk

Relevant published information was not found as of the revision date.

Alternate Drugs to Consider

Formoterol, Salmeterol

References

- Lindberg C, Boreus LO, de Chateau P et al. Transfer of terbutaline into breast milk. *Eur J Resp Dis Suppl.* 1984;134:87-91. PubMed PMID: 6586490.
- McDonald CF, Burdon JG. Asthma in pregnancy and lactation. A position paper for the Thoracic Society of Australia and New Zealand. *Med J Aust.* 1996;165:485-8. PubMed PMID: 8937369.
- Ellsworth A. Pharmacotherapy of asthma while breastfeeding. *J Hum Lact.* 1994;10:39-41. PubMed PMID: 7619245.
- Nelson-Piercy C. Asthma in pregnancy. *Thorax.* 2001;56:325-8. PubMed PMID: 11254828.
- Taddio A, Ito S. Drugs and breast-feeding. In, Koren G, ed. *Maternal-fetal toxicology. A clinician's guide.* 3rd ed. New York. Marcel Dekker. 2001;177-32.

Substance Identification

Substance Name

Indacaterol

CAS Registry Number

312753-06-3

Drug Class

Breast Feeding

Lactation

Anti-Asthmatic Agents

Beta Adrenergic Agonists

Bronchodilator Agents