

**Table A.1.2 Unweighted Logistic Regression of Screener Contact: 2003**

<b>Variable</b>	<b>Coefficient</b>	<b>Standard Error</b>	<b>P Value</b>	<b>Odds Ratio</b>	<b>Lower 95% OR</b>	<b>Upper 95% OR</b>
<b>Intercept</b>	3.88	0.21	0.0000	48.21	31.95	72.76
<b>Interviewer Gender</b>						
Male (RC)	0.00	0.00		1.00	1.00	1.00
Female	-0.11	0.07	0.1333	0.90	0.78	1.03
<b>Interviewer Race/Ethnicity</b>						
White, Non-Hispanic (RC)	0.00	0.00		1.00	1.00	1.00
Black, Non-Hispanic	-0.27	0.11	0.0102	0.76	0.62	0.94
Other, Non-Hispanic	-0.31	0.21	0.1397	0.73	0.48	1.11
Hispanic	-0.35	0.19	0.0572	0.70	0.49	1.01
<b>Average Miles per Week during the Quarter</b>						
0 to 135 (RC)	0.00	0.00		1.00	1.00	1.00
136 to 250	-0.01	0.12	0.9260	0.99	0.78	1.25
251 to 415	-0.09	0.12	0.4545	0.91	0.72	1.16
416+	-0.35	0.16	0.0290	0.70	0.51	0.96
<b>Average Hours per Week during the Quarter</b>						
1 to 18 (RC)	0.00	0.00		1.00	1.00	1.00
18.25 to 26	0.12	0.11	0.3010	1.12	0.90	1.40
26.25 to 35.75	0.17	0.13	0.1928	1.18	0.92	1.52
36+	0.40	0.17	0.0212	1.49	1.06	2.09
<b>Cumulative Interview Count</b>						
1 to 19 (RC)	0.00	0.00		1.00	1.00	1.00
20 to 39	-0.11	0.20	0.5865	0.90	0.61	1.32
40 to 59	-0.46	0.18	0.0089	0.63	0.44	0.89
60 to 99	-0.06	0.15	0.6688	0.94	0.70	1.25
100 to 249	-0.08	0.13	0.5540	0.93	0.72	1.19
250 to 499	0.19	0.12	0.1151	1.21	0.95	1.54
500 to 749	0.04	0.16	0.8181	1.04	0.76	1.42
750 to 999	0.35	0.21	0.1074	1.41	0.93	2.15
1,000+	0.56	0.25	0.0235	1.76	1.08	2.86

RC = reference category.

Source: Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2003.