

**Table A.1.3 Unweighted Logistic Regression of Screener Contact: 2004**

<b>Variable</b>	<b>Coefficient</b>	<b>Standard Error</b>	<b>P Value</b>	<b>Odds Ratio</b>	<b>Lower 95% OR</b>	<b>Upper 95% OR</b>
<b>Intercept</b>	3.81	0.26	0.0000	45.30	27.32	75.11
<b>Interviewer Gender</b>						
Male (RC)	0.00	0.00		1.00	1.00	1.00
Female	-0.07	0.09	0.4565	0.93	0.78	1.12
<b>Interviewer Race/Ethnicity</b>						
White, Non-Hispanic (RC)	0.00	0.00		1.00	1.00	1.00
Black, Non-Hispanic	-0.46	0.15	0.0021	0.63	0.47	0.85
Other, Non-Hispanic	-0.34	0.27	0.2102	0.71	0.42	1.21
Hispanic	-0.42	0.17	0.0122	0.66	0.47	0.91
<b>Average Miles per Week during the Quarter</b>						
0 to 135 (RC)	0.00	0.00		1.00	1.00	1.00
136 to 250	0.16	0.14	0.2401	1.17	0.90	1.53
251 to 415	-0.05	0.14	0.7218	0.95	0.72	1.26
416+	-0.23	0.19	0.2277	0.80	0.55	1.15
<b>Average Hours per Week during the Quarter</b>						
1 to 18 (RC)	0.00	0.00		1.00	1.00	1.00
18.25 to 26	0.01	0.10	0.8882	1.01	0.83	1.23
26.25 to 35.75	0.10	0.12	0.4162	1.10	0.87	1.40
36+	0.09	0.14	0.5026	1.10	0.84	1.44
<b>Cumulative Interview Count</b>						
1 to 19 (RC)	0.00	0.00		1.00	1.00	1.00
20 to 39	-0.06	0.30	0.8361	0.94	0.53	1.68
40 to 59	-0.23	0.31	0.4528	0.79	0.43	1.45
60 to 99	0.05	0.18	0.7601	1.06	0.74	1.50
100 to 249	-0.18	0.14	0.2029	0.83	0.63	1.10
250 to 499	-0.11	0.15	0.4525	0.89	0.66	1.20
500 to 749	-0.00	0.13	0.9737	1.00	0.77	1.29
750 to 999	0.26	0.16	0.0909	1.30	0.96	1.77
1,000+	0.59	0.26	0.0231	1.81	1.08	3.01

RC = reference category.

Source: Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2004.