

**Table A.1.4 Unweighted Logistic Regression of Screener Contact: 2005**

<b>Variable</b>	<b>Coefficient</b>	<b>Standard Error</b>	<b>P Value</b>	<b>Odds Ratio</b>	<b>Lower 95% OR</b>	<b>Upper 95% OR</b>
<b>Intercept</b>	4.85	0.27	0.0000	127.59	75.59	215.37
<b>Interviewer Gender</b>						
Male (RC)	0.00	0.00		1.00	1.00	1.00
Female	-0.13	0.08	0.0933	0.87	0.75	1.02
<b>Interviewer Race/Ethnicity</b>						
White, Non-Hispanic (RC)	0.00	0.00		1.00	1.00	1.00
Black, Non-Hispanic	-0.55	0.11	0.0000	0.58	0.47	0.71
Other, Non-Hispanic	-0.13	0.24	0.6040	0.88	0.55	1.42
Hispanic	-0.47	0.13	0.0003	0.62	0.48	0.80
<b>Average Miles per Week during the Quarter</b>						
0 to 135 (RC)	0.00	0.00		1.00	1.00	1.00
136 to 250	0.05	0.09	0.5320	1.06	0.89	1.25
251 to 415	0.05	0.13	0.7010	1.05	0.82	1.35
416+	-0.08	0.15	0.5838	0.92	0.68	1.24
<b>Average Hours per Week during the Quarter</b>						
1 to 18 (RC)	0.00	0.00		1.00	1.00	1.00
18.25 to 26	0.03	0.10	0.7456	1.03	0.85	1.25
26.25 to 35.75	0.11	0.11	0.2981	1.12	0.90	1.39
36+	0.30	0.16	0.0637	1.34	0.98	1.84
<b>Cumulative Interview Count</b>						
1 to 19 (RC)	0.00	0.00		1.00	1.00	1.00
20 to 39	-0.27	0.16	0.0846	0.76	0.56	1.04
40 to 59	-0.27	0.21	0.1922	0.76	0.51	1.14
60 to 99	-0.38	0.25	0.1324	0.68	0.42	1.12
100 to 249	-0.11	0.14	0.4225	0.89	0.68	1.18
250 to 499	-0.31	0.14	0.0238	0.73	0.56	0.96
500 to 749	-0.03	0.14	0.8178	0.97	0.74	1.27
750 to 999	-0.35	0.21	0.0960	0.71	0.47	1.06
1,000+	0.27	0.17	0.1137	1.30	0.94	1.81

RC = reference category.

Source: Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2005.