

**Table C1.3 Adjusted Prevalence Estimates by Experience Measures for 2002 Cohort – Lifetime Use of Alcohol**

| Number of Interviews Completed at Time of Interview | Eventual Experience: Number of Interviews Completed by the End of 2007 |          |          |          |            |            |            |       |
|---|--|----------|----------|----------|------------|------------|------------|-------|
|   | 1 to 19  | 20 to 39 | 40 to 59 | 60 to 99 | 100 to 249 | 250 to 499 | 500 to 749 | 750+  |
| 1 to 19   | 0.734  | 0.723    | 0.701    | 0.726    | 0.738      | 0.711      | 0.699      | 0.699 |
| 20 to 39  | --   | 0.709    | 0.724    | 0.707    | 0.710      | 0.724      | 0.701      | 0.690 |
| 40 to 59  | --   | --       | 0.713    | 0.718    | 0.716      | 0.716      | 0.719      | 0.717 |
| 60 to 99  | --   | --       | --       | 0.724    | 0.718      | 0.719      | 0.728      | 0.703 |
| 100 to 249  | --   | --       | --       | --       | 0.729      | 0.703      | 0.716      | 0.722 |
| 250 to 499  | --   | --       | --       | --       | --         | 0.698      | 0.717      | 0.716 |
| 500 to 749  | --   | --       | --       | --       | --         | --         | 0.727      | 0.714 |
| 750+  | --   | --       | --       | --       | --         | --         | --         | 0.725 |

Source: Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002 to 2007.

**Table C1.4 Adjusted Prevalence Estimates by Experience Measures for 2002 Cohort – Past Year Use of Alcohol**

| Number of Interviews Completed at Time of Interview | Eventual Experience: Number of Interviews Completed by the End of 2007 |          |          |          |            |            |            |       |
|---|--|----------|----------|----------|------------|------------|------------|-------|
|   | 1 to 19  | 20 to 39 | 40 to 59 | 60 to 99 | 100 to 249 | 250 to 499 | 500 to 749 | 750+  |
| 1 to 19   | 0.627  | 0.636    | 0.569    | 0.607    | 0.631      | 0.615      | 0.569      | 0.619 |
| 20 to 39  | --   | 0.596    | 0.634    | 0.590    | 0.596      | 0.621      | 0.602      | 0.536 |
| 40 to 59  | --   | --       | 0.592    | 0.591    | 0.597      | 0.591      | 0.591      | 0.545 |
| 60 to 99  | --   | --       | --       | 0.616    | 0.598      | 0.596      | 0.599      | 0.583 |
| 100 to 249  | --   | --       | --       | --       | 0.616      | 0.589      | 0.606      | 0.603 |
| 250 to 499  | --   | --       | --       | --       | --         | 0.582      | 0.604      | 0.596 |
| 500 to 749  | --   | --       | --       | --       | --         | --         | 0.613      | 0.605 |
| 750+  | --   | --       | --       | --       | --         | --         | --         | 0.604 |

Source: Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002 to 2007.