

Table C1.7 Adjusted Prevalence Estimates by Experience Measures for 2002 Cohort – Lifetime Nonmedical Use of Analgesics

| Number of Interviews Completed at Time of Interview | Eventual Experience: Number of Interviews Completed by the End of 2007 | | | | | | | |
|---|--|----------|----------|----------|------------|------------|------------|-------|
| | 1 to 19 | 20 to 39 | 40 to 59 | 60 to 99 | 100 to 249 | 250 to 499 | 500 to 749 | 750+ |
| 1 to 19 | 0.174 | 0.177 | 0.161 | 0.199 | 0.178 | 0.209 | 0.177 | 0.171 |
| 20 to 39 | -- | 0.164 | 0.143 | 0.191 | 0.196 | 0.191 | 0.158 | 0.184 |
| 40 to 59 | -- | -- | 0.177 | 0.163 | 0.168 | 0.169 | 0.194 | 0.230 |
| 60 to 99 | -- | -- | -- | 0.164 | 0.170 | 0.182 | 0.160 | 0.167 |
| 100 to 249 | -- | -- | -- | -- | 0.178 | 0.177 | 0.170 | 0.186 |
| 250 to 499 | -- | -- | -- | -- | -- | 0.160 | 0.178 | 0.176 |
| 500 to 749 | -- | -- | -- | -- | -- | -- | 0.176 | 0.170 |
| 750+ | -- | -- | -- | -- | -- | -- | -- | 0.181 |

Source: Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002 to 2007.

Table C1.8 Adjusted Prevalence Estimates by Experience Measures for 2002 Cohort – Past Year Nonmedical Use of Analgesics

| Number of Interviews Completed at Time of Interview | Eventual Experience: Number of Interviews Completed by the End of 2007 | | | | | | | |
|---|--|----------|----------|----------|------------|------------|------------|-------|
| | 1 to 19 | 20 to 39 | 40 to 59 | 60 to 99 | 100 to 249 | 250 to 499 | 500 to 749 | 750+ |
| 1 to 19 | 0.084 | 0.083 | 0.064 | 0.104 | 0.078 | 0.103 | 0.089 | 0.085 |
| 20 to 39 | -- | 0.081 | 0.058 | 0.088 | 0.100 | 0.096 | 0.079 | 0.120 |
| 40 to 59 | -- | -- | 0.078 | 0.067 | 0.075 | 0.078 | 0.107 | 0.148 |
| 60 to 99 | -- | -- | -- | 0.067 | 0.084 | 0.075 | 0.066 | 0.090 |
| 100 to 249 | -- | -- | -- | -- | 0.081 | 0.088 | 0.082 | 0.078 |
| 250 to 499 | -- | -- | -- | -- | -- | 0.078 | 0.087 | 0.092 |
| 500 to 749 | -- | -- | -- | -- | -- | -- | 0.088 | 0.082 |
| 750+ | -- | -- | -- | -- | -- | -- | -- | 0.085 |

Source: Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002 to 2007.