

**Table C1.9 Adjusted Prevalence Estimates by Experience Measures for 2002 Cohort – Lifetime Use of Cocaine**

| Number of Interviews Completed at Time of Interview | Eventual Experience: Number of Interviews Completed by the End of 2007 |          |          |          |            |            |            |       |
|---|--|----------|----------|----------|------------|------------|------------|-------|
|   | 1 to 19  | 20 to 39 | 40 to 59 | 60 to 99 | 100 to 249 | 250 to 499 | 500 to 749 | 750+  |
| 1 to 19   | 0.145  | 0.127    | 0.138    | 0.142    | 0.147      | 0.144      | 0.125      | 0.100 |
| 20 to 39  | --   | 0.119    | 0.115    | 0.125    | 0.124      | 0.124      | 0.145      | 0.076 |
| 40 to 59  | --   | --       | 0.141    | 0.120    | 0.129      | 0.144      | 0.118      | 0.168 |
| 60 to 99  | --   | --       | --       | 0.112    | 0.119      | 0.143      | 0.121      | 0.141 |
| 100 to 249  | --   | --       | --       | --       | 0.124      | 0.123      | 0.138      | 0.135 |
| 250 to 499  | --   | --       | --       | --       | --         | 0.120      | 0.124      | 0.126 |
| 500 to 749  | --   | --       | --       | --       | --         | --         | 0.131      | 0.125 |
| 750+  | --   | --       | --       | --       | --         | --         | --         | 0.134 |

Source: Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002 to 2007.

**Table C1.10 Adjusted Prevalence Estimates by Experience Measures for 2002 Cohort – Positive Responses to Gate Items**

| Number of Interviews Completed at Time of Interview | Eventual Experience: Number of Interviews Completed by the End of 2007 |          |          |          |            |            |            |       |
|---|--|----------|----------|----------|------------|------------|------------|-------|
|   | 1 to 19  | 20 to 39 | 40 to 59 | 60 to 99 | 100 to 249 | 250 to 499 | 500 to 749 | 750+  |
| 1 to 19   | 4.037  | 3.732    | 3.551    | 3.936    | 3.953      | 3.807      | 3.526      | 3.256 |
| 20 to 39  | --   | 3.589    | 3.448    | 3.829    | 3.660      | 3.691      | 3.735      | 3.216 |
| 40 to 59  | --   | --       | 3.893    | 3.737    | 3.711      | 3.845      | 3.646      | 4.310 |
| 60 to 99  | --   | --       | --       | 3.867    | 3.703      | 3.765      | 3.640      | 3.697 |
| 100 to 249  | --   | --       | --       | --       | 3.800      | 3.590      | 3.748      | 3.659 |
| 250 to 499  | --   | --       | --       | --       | --         | 3.503      | 3.712      | 3.629 |
| 500 to 749  | --   | --       | --       | --       | --         | --         | 3.790      | 3.635 |
| 750+  | --   | --       | --       | --       | --         | --         | --         | 3.935 |

Source: Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002 to 2007.