

Table C2.7 Adjusted Prevalence Estimates by Experience Measures for 2003 Cohort – Lifetime Nonmedical Use of Analgesics

| Number of Interviews Completed at Time of Interview | Eventual Experience: Number of Interviews Completed by the End of 2007 | | | | | | | |
|---|--|----------|----------|----------|------------|------------|------------|-------|
| | 1 to 19 | 20 to 39 | 40 to 59 | 60 to 99 | 100 to 249 | 250 to 499 | 500 to 749 | 750+ |
| 1 to 19 | 0.139 | 0.175 | 0.187 | 0.197 | 0.182 | 0.159 | 0.167 | 0.186 |
| 20 to 39 | -- | 0.162 | 0.184 | 0.162 | 0.155 | 0.187 | 0.198 | 0.124 |
| 40 to 59 | -- | -- | 0.172 | 0.171 | 0.167 | 0.173 | 0.178 | 0.231 |
| 60 to 99 | -- | -- | -- | 0.186 | 0.161 | 0.167 | 0.166 | 0.160 |
| 100 to 249 | -- | -- | -- | -- | 0.161 | 0.170 | 0.159 | 0.146 |
| 250 to 499 | -- | -- | -- | -- | -- | 0.170 | 0.162 | 0.149 |
| 500 to 749 | -- | -- | -- | -- | -- | -- | 0.161 | 0.143 |
| 750+ | -- | -- | -- | -- | -- | -- | -- | 0.150 |

Source: Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2003 to 2007.

Table C2.8 Adjusted Prevalence Estimates by Experience Measures for 2003 Cohort – Past Year Nonmedical Use of Analgesics

| Number of Interviews Completed at Time of Interview | Eventual Experience: Number of Interviews Completed by the End of 2007 | | | | | | | |
|---|--|----------|----------|----------|------------|------------|------------|-------|
| | 1 to 19 | 20 to 39 | 40 to 59 | 60 to 99 | 100 to 249 | 250 to 499 | 500 to 749 | 750+ |
| 1 to 19 | 0.054 | 0.092 | 0.085 | 0.092 | 0.093 | 0.066 | 0.085 | 0.081 |
| 20 to 39 | -- | 0.061 | 0.063 | 0.059 | 0.069 | 0.085 | 0.113 | 0.053 |
| 40 to 59 | -- | -- | 0.069 | 0.068 | 0.077 | 0.084 | 0.074 | 0.138 |
| 60 to 99 | -- | -- | -- | 0.083 | 0.071 | 0.071 | 0.097 | 0.076 |
| 100 to 249 | -- | -- | -- | -- | 0.077 | 0.076 | 0.080 | 0.072 |
| 250 to 499 | -- | -- | -- | -- | -- | 0.080 | 0.076 | 0.074 |
| 500 to 749 | -- | -- | -- | -- | -- | -- | 0.074 | 0.061 |
| 750+ | -- | -- | -- | -- | -- | -- | -- | 0.060 |

Source: Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2003 to 2007.