

Table SU05.12 Needed and Received Treatment for an Illicit Drug or Alcohol Problem in the Past Year among Individuals Aged 12 or Older, by Demographic Characteristics: Comparison of Percentages and Standard Errors from Restricted-Use File and Public Use File Estimates, 2005 NSDUH

Demographic Characteristic	NEEDED TREATMENT FOR AN ILLICIT DRUG OR ALCOHOL PROBLEM IN THE PAST YEAR																		Percentage Who Received Treatment at a Specialty Facility among Individuals Who Needed Treatment											
	Total						Received Treatment at a Specialty Facility						Did Not Receive Treatment at a Specialty Facility						RUF Estimate		PUF Estimate		Ratio of Estimates		RUF SE		PUF SE		Ratio of SEs	
	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs						
TOTAL	9.5	9.4	0.99	0.18	0.19	1.02	0.9	1.0	1.01	0.06	0.07	1.12	8.6	8.4	0.99	0.17	0.17	1.00	10.0	10.1	1.02	0.63	0.71	1.12						
AGE																														
12-17	8.3	8.3	1.00	0.24	0.24	0.99	0.7	0.7	0.98	0.07	0.08	1.16	7.6	7.6	1.00	0.23	0.22	0.97	8.6	8.5	0.98	0.77	0.88	1.14						
18-25	22.2	22.1	1.00	0.41	0.41	1.01	1.6	1.6	1.02	0.10	0.13	1.30	20.6	20.5	0.99	0.40	0.41	1.03	7.2	7.4	1.02	0.46	0.60	1.30						
26 or Older	7.5	7.3	0.98	0.21	0.21	1.00	0.9	0.9	1.00	0.08	0.09	1.14	6.6	6.5	0.98	0.20	0.19	0.98	11.6	11.9	1.02	1.00	1.15	1.15						
GENDER																														
Male	12.6	12.6	1.00	0.30	0.32	1.07	1.3	1.3	1.03	0.11	0.12	1.09	11.3	11.3	0.99	0.28	0.29	1.06	10.2	10.5	1.03	0.84	0.90	1.08						
Female	6.6	6.4	0.97	0.19	0.20	1.03	0.6	0.6	0.96	0.06	0.06	0.95	6.0	5.8	0.97	0.18	0.19	1.04	9.5	9.4	0.99	0.91	0.90	0.99						
HISPANIC ORIGIN AND RACE																														
Not Hispanic or Latino	9.5	9.3	0.98	0.19	0.19	0.99	0.9	0.9	1.00	0.07	0.08	1.17	8.5	8.4	0.98	0.18	0.18	1.02	9.7	9.8	1.02	0.67	0.81	1.20						
White	9.6	9.5	0.98	0.21	0.21	1.02	0.8	0.8	1.01	0.06	0.08	1.27	8.8	8.6	0.98	0.20	0.20	0.98	8.5	8.7	1.02	0.63	0.79	1.26						
Black or African American	9.6	9.6	1.00	0.56	0.58	1.04	1.8	1.8	0.99	0.31	0.35	1.11	7.9	7.9	1.00	0.48	0.51	1.07	18.4	18.3	1.00	2.84	3.23	1.14						
American Indian or Alaska Native	21.5	23.9	1.11	3.64	4.42	1.21	1.9	1.5	0.81	0.49	0.52	1.05	19.7	22.4	1.14	3.64	4.51	1.24	8.7	6.3	0.73	2.65	2.58	0.98						
Native Hawaiian or Other Pacific Islander	11.0	10.0	0.91	2.76	2.50	0.91	0.3	0.4	1.06	0.22	0.22	1.01	10.7	9.7	0.91	2.71	2.42	0.89	*	3.6	*	*	2.01	*						
Asian	4.5	4.2	0.93	0.57	0.54	0.94	0.0	0.0	0.69	0.03	0.03	0.85	4.5	4.1	0.93	0.57	0.54	0.94	1.0	0.8	0.75	0.68	0.63	0.93						
Two or More Races	11.0	9.6	0.87	1.79	1.04	0.58	1.0	1.0	1.05	0.29	0.30	1.02	10.0	8.6	0.86	1.78	1.03	0.58	8.8	10.6	1.20	2.85	3.12	1.10						
Hispanic or Latino	9.9	10.0	1.01	0.54	0.57	1.05	1.2	1.2	1.03	0.22	0.22	0.96	8.8	8.8	1.00	0.48	0.52	1.07	11.7	12.0	1.02	2.04	1.97	0.97						

* Low precision; no estimate reported.

PUF = public use file, which refers to estimates produced from a NSDUH public use file (a PUF is a NSDUH restricted-use file that has undergone statistical disclosure limitation treatment); RUF = restricted-use file, which refers to estimates produced from a NSDUH RUF and published in NSDUH's detailed tables available at <http://www.samhsa.gov/data/>; SE = standard error that is associated with the estimate.

NOTE: $Ratio\ of\ estimates = PUF\ estimate \div RUF\ estimate$; $Ratio\ of\ SEs = PUF\ SE \div RUF\ SE$.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2005.