

Table SU07.12 Need and Received Treatment for an Illicit Drug or Alcohol Problem in the Past Year among Individuals Aged 12 or Older, by Demographic Characteristics: Comparison of Percentages and Standard Errors from Restricted-Use File and Public Use File Estimates, 2007 NSDUH

Demographic Characteristic	NEEDED TREATMENT FOR AN ILLICIT DRUG OR ALCOHOL PROBLEM IN THE PAST YEAR																		Percentage Who Received Treatment at a Specialty Facility among Individuals Who Needed Treatment					
	Total						Received Treatment at a Specialty Facility						Did Not Receive Treatment at a Specialty Facility						RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs
	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs
TOTAL	9.4	9.4	1.00	0.19	0.21	1.12	1.0	1.0	0.98	0.07	0.08	1.26	8.4	8.4	1.00	0.18	0.19	1.10	10.4	10.2	0.98	0.66	0.83	1.26
AGE																								
12-17	7.9	8.0	1.01	0.23	0.29	1.27	0.6	0.6	1.04	0.06	0.07	1.16	7.3	7.3	1.01	0.22	0.27	1.22	7.6	7.8	1.02	0.77	0.84	1.09
18-25	21.2	21.6	1.02	0.39	0.48	1.23	1.5	1.5	0.98	0.10	0.11	1.10	19.7	20.1	1.02	0.38	0.47	1.24	7.0	6.8	0.97	0.47	0.51	1.09
26 or Older	7.6	7.4	0.99	0.23	0.23	1.02	0.9	0.9	0.97	0.08	0.11	1.25	6.6	6.5	0.99	0.21	0.21	0.98	12.5	12.3	0.98	1.03	1.29	1.25
GENDER																								
Male	13.0	13.1	1.01	0.33	0.34	1.03	1.4	1.4	0.98	0.12	0.14	1.20	11.5	11.7	1.01	0.31	0.31	1.03	11.0	10.7	0.98	0.86	1.04	1.20
Female	6.0	5.9	0.98	0.18	0.26	1.42	0.6	0.5	0.96	0.06	0.08	1.31	5.4	5.3	0.98	0.17	0.23	1.36	9.3	9.1	0.98	0.93	1.19	1.28
HISPANIC ORIGIN AND RACE																								
Not Hispanic or Latino	9.5	9.5	1.00	0.20	0.24	1.21	1.1	1.0	0.97	0.08	0.09	1.25	8.5	8.5	1.00	0.18	0.22	1.18	11.1	10.8	0.98	0.74	0.92	1.24
White	9.8	9.8	1.00	0.23	0.28	1.24	1.0	0.9	0.96	0.08	0.09	1.25	8.8	8.8	1.01	0.21	0.26	1.21	10.0	9.5	0.96	0.74	0.91	1.23
Black or African American	9.3	9.0	0.97	0.57	0.72	1.26	1.7	1.7	1.01	0.32	0.36	1.10	7.6	7.3	0.96	0.48	0.58	1.21	18.2	18.8	1.04	3.01	3.27	1.09
American Indian or Alaska Native	14.5	14.2	0.98	2.90	2.70	0.93	3.4	3.2	0.94	1.32	1.31	0.99	11.1	11.0	0.99	2.61	2.42	0.93	*	22.4	*	*	8.27	*
Native Hawaiian or Other Pacific Islander	10.0	10.4	1.04	2.76	2.88	1.04	0.1	0.1	1.14	0.07	0.08	1.14	9.9	10.4	1.04	2.76	2.88	1.04	*	0.7	*	*	0.74	*
Asian	4.9	4.7	0.97	0.68	0.75	1.11	0.2	0.3	1.32	0.17	0.25	1.49	4.7	4.5	0.95	0.65	0.71	1.10	*	5.7	*	*	5.13	*
Two or More Races	11.7	11.3	0.97	1.35	1.45	1.07	1.8	1.8	1.00	0.67	0.75	1.12	9.9	9.5	0.96	1.19	1.26	1.06	*	16.2	*	*	5.86	*
Hispanic or Latino	8.6	8.6	1.00	0.49	0.47	0.96	0.5	0.5	1.00	0.10	0.10	0.99	8.1	8.1	1.00	0.48	0.45	0.94	6.0	6.0	0.99	1.15	1.10	0.96

* Low precision; no estimate reported.

PUF = public use file, which refers to estimates produced from a NSDUH public use file (a PUF is a NSDUH restricted-use file that has undergone statistical disclosure limitation treatment); RUF = restricted-use file, which refers to estimates produced from a NSDUH RUF and published in NSDUH's detailed tables available at <http://www.samhsa.gov/data/>; SE = standard error that is associated with the estimate.NOTE: Some 2007 RUF estimates and SEs may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).NOTE: $Ratio\ of\ estimates = PUF\ estimate \div RUF\ estimate$; $Ratio\ of\ SEs = PUF\ SE \div RUF\ SE$.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for illicit drug or alcohol use at a specialty facility (i.e., drug and alcohol rehabilitation facility [inpatient or outpatient], hospital [inpatient only], or mental health center). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2007.