

Table SU10.7 Types of Illicit Drug Use in the Past Month among Individuals Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Comparison of Percentages and Standard Errors from Restricted-Use File and Public Use File Estimates, 2010 NSDUH

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH																							
	Cigarettes and Alcohol						Cigarettes But Not Alcohol						Alcohol But Not Cigarettes						Neither Alcohol Nor Cigarettes					
	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs
ILLCIT DRUGS¹	27.7	27.8	1.01	0.66	0.74	1.11	12.6	12.7	1.00	0.81	0.95	1.17	7.6	7.8	1.03	0.31	0.28	0.93	2.4	2.4	1.01	0.14	0.16	1.13
Marijuana and Hashish	23.3	23.5	1.01	0.62	0.71	1.15	9.5	9.6	1.01	0.70	0.77	1.10	5.8	6.0	1.04	0.26	0.23	0.89	1.2	1.2	1.02	0.09	0.09	0.96
Cocaine	2.8	2.9	1.02	0.24	0.24	0.99	0.6	0.6	0.94	0.15	0.15	0.99	0.2	0.2	0.87	0.04	0.04	0.95	0.1	0.1	1.05	0.02	0.02	1.05
Crack	0.7	0.7	1.02	0.13	0.14	1.06	0.3	0.2	0.83	0.11	0.10	0.90	0.0	0.0	0.54	0.02	0.01	0.59	0.0	0.0	1.21	0.01	0.02	1.24
Heroin	0.4	0.5	1.07	0.09	0.10	1.11	0.2	0.2	0.94	0.08	0.09	1.02	*	0.0	*	*	0.00	*	0.0	0.0	0.90	0.02	0.02	1.02
Hallucinogens	2.2	2.1	0.98	0.18	0.20	1.10	0.3	0.4	1.10	0.06	0.07	1.12	0.3	0.2	0.98	0.03	0.03	0.99	0.1	0.1	1.07	0.02	0.02	0.99
LSD	0.3	0.3	1.01	0.04	0.05	1.15	0.0	0.1	1.19	0.02	0.02	1.31	0.0	0.0	0.88	0.01	0.01	0.99	*	0.0	*	*	0.00	*
PCP	0.0	0.0	1.08	0.01	0.01	1.35	0.0	0.0	1.13	0.00	0.00	1.09	*	0.0	*	*	0.00	*	0.0	0.0	1.11	0.02	0.02	1.04
Ecstasy	1.3	1.2	0.95	0.14	0.14	1.00	0.2	0.2	1.09	0.05	0.06	1.14	0.1	0.1	0.91	0.03	0.02	0.81	0.0	0.0	1.03	0.01	0.01	1.16
Inhalants	0.6	0.7	1.12	0.10	0.11	1.13	0.1	0.1	1.16	0.03	0.02	0.75	0.3	0.3	0.97	0.08	0.08	1.02	0.2	0.2	0.94	0.02	0.03	1.22
Nonmedical Use of Psychotherapeutics ^{2,3}	7.9	7.9	1.00	0.37	0.35	0.95	3.9	4.0	1.02	0.43	0.43	1.00	2.2	2.2	1.01	0.17	0.15	0.90	1.1	1.1	1.01	0.11	0.12	1.07
Pain Relievers	6.1	6.1	1.01	0.33	0.36	1.09	2.7	2.8	1.05	0.37	0.37	1.00	1.5	1.5	1.04	0.13	0.14	1.05	0.8	0.8	1.03	0.08	0.10	1.16
OxyContin [®]	0.9	0.9	0.95	0.11	0.11	0.96	0.7	0.7	1.06	0.24	0.24	0.98	0.1	0.1	1.14	0.02	0.03	1.25	0.0	0.0	0.95	0.00	0.01	1.13
Tranquilizers	2.8	2.8	1.00	0.22	0.21	0.93	1.4	1.3	0.90	0.25	0.23	0.89	0.6	0.6	1.00	0.08	0.08	0.98	0.2	0.2	1.04	0.05	0.05	1.07
Stimulants ³	1.5	1.5	1.00	0.16	0.14	0.87	0.6	0.6	1.00	0.13	0.14	1.08	0.3	0.3	1.00	0.04	0.05	1.09	0.1	0.1	0.73	0.03	0.02	0.78
Methamphetamine ³	0.5	0.5	1.08	0.09	0.10	1.19	0.3	0.3	1.14	0.10	0.13	1.29	0.1	0.1	1.04	0.03	0.03	1.25	0.0	0.0	1.02	0.02	0.02	1.02
Sedatives	0.3	0.3	0.96	0.08	0.09	1.08	0.1	0.1	1.17	0.07	0.08	1.13	0.1	0.1	0.58	0.06	0.03	0.49	0.1	0.1	1.02	0.04	0.04	1.09
ILLCIT DRUGS OTHER THAN MARIJUANA⁴	11.2	11.2	1.01	0.44	0.41	0.93	4.6	4.7	1.02	0.46	0.45	0.99	2.6	2.6	0.99	0.18	0.17	0.92	1.3	1.3	1.00	0.11	0.11	1.03

* Low precision; no estimate reported.

LSD = lysergic acid diethylamide; PCP = phencyclidine; PUF = public use file, which refers to estimates produced from a NSDUH public use file (a PUF is a NSDUH restricted-use file that has undergone statistical disclosure limitation treatment); RUF = restricted-use file, which refers to estimates produced from a NSDUH RUF and published in NSDUH's detailed tables available at <http://www.samhsa.gov/data/>; SE = standard error that is associated with the estimate.

NOTE: Some 2010 RUF estimates and SEs may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: $Ratio\ of\ estimates = PUF\ estimate \div RUF\ estimate$; $Ratio\ of\ SEs = PUF\ SE \div RUF\ SE$.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from new methamphetamine items added in 2005 and 2006. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from new methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in NSDUH reports prior to the 2007 national findings report. See Section B.4.8 in Appendix B of the *Results from the 2008 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010.