

Table SU10.10 Drove Under the Influence of Illicit Drugs or Alcohol in the Past Year among Individuals Aged 12 or Older, by Demographic Characteristics: Comparison of Percentages and Standard Errors from Restricted-Use File and Public Use File Estimates, 2010 NSDUH

| Demographic Characteristic | ILLCIT DRUGS | | | | | | ALCOHOL | | | | | | ILLCIT DRUGS OR ALCOHOL | | | | | |
|---|--------------|--------------|--------------------|--------|--------|--------------|--------------|--------------|--------------------|--------|--------|--------------|-------------------------|--------------|--------------------|--------|--------|--------------|
| | RUF Estimate | PUF Estimate | Ratio of Estimates | RUF SE | PUF SE | Ratio of SEs | RUF Estimate | PUF Estimate | Ratio of Estimates | RUF SE | PUF SE | Ratio of SEs | RUF Estimate | PUF Estimate | Ratio of Estimates | RUF SE | PUF SE | Ratio of SEs |
| TOTAL | 4.2 | 4.2 | 1.01 | 0.12 | 0.11 | 0.93 | 11.4 | 11.4 | 1.00 | 0.22 | 0.25 | 1.10 | 12.5 | 12.5 | 1.00 | 0.23 | 0.26 | 1.13 |
| AGE | | | | | | | | | | | | | | | | | | |
| 12-17 | 2.7 | 2.6 | 0.98 | 0.14 | 0.17 | 1.18 | 2.5 | 2.4 | 0.99 | 0.15 | 0.17 | 1.15 | 3.5 | 3.4 | 0.99 | 0.17 | 0.20 | 1.18 |
| 18-25 | 12.7 | 12.8 | 1.00 | 0.30 | 0.29 | 0.96 | 20.0 | 20.3 | 1.01 | 0.40 | 0.43 | 1.06 | 23.3 | 23.5 | 1.01 | 0.42 | 0.42 | 0.99 |
| 26 or Older | 2.9 | 2.9 | 1.01 | 0.14 | 0.15 | 1.04 | 11.0 | 11.0 | 1.00 | 0.27 | 0.31 | 1.11 | 11.7 | 11.7 | 1.00 | 0.28 | 0.32 | 1.16 |
| GENDER | | | | | | | | | | | | | | | | | | |
| Male | 5.8 | 5.9 | 1.01 | 0.20 | 0.22 | 1.08 | 15.1 | 15.2 | 1.01 | 0.37 | 0.46 | 1.26 | 16.5 | 16.6 | 1.01 | 0.38 | 0.50 | 1.32 |
| Female | 2.6 | 2.6 | 1.00 | 0.12 | 0.12 | 0.96 | 7.9 | 7.8 | 0.99 | 0.23 | 0.21 | 0.93 | 8.7 | 8.6 | 0.99 | 0.24 | 0.23 | 0.98 |
| HISPANIC ORIGIN AND RACE | | | | | | | | | | | | | | | | | | |
| Not Hispanic or Latino | 4.3 | 4.3 | 1.00 | 0.13 | 0.11 | 0.86 | 12.0 | 12.0 | 1.00 | 0.24 | 0.26 | 1.06 | 13.1 | 13.1 | 1.00 | 0.25 | 0.27 | 1.07 |
| White | 4.5 | 4.5 | 1.00 | 0.15 | 0.12 | 0.83 | 13.2 | 13.2 | 1.00 | 0.29 | 0.30 | 1.05 | 14.4 | 14.4 | 1.00 | 0.29 | 0.31 | 1.07 |
| Black or African American | 4.0 | 4.2 | 1.03 | 0.38 | 0.47 | 1.24 | 8.0 | 7.9 | 0.99 | 0.53 | 0.50 | 0.95 | 8.9 | 8.9 | 1.00 | 0.56 | 0.57 | 1.01 |
| American Indian or Alaska Native | 7.1 | 6.5 | 0.92 | 1.63 | 1.43 | 0.88 | 11.2 | 11.2 | 1.00 | 2.04 | 2.15 | 1.05 | 12.5 | 12.3 | 0.99 | 2.06 | 2.17 | 1.06 |
| Native Hawaiian or Other Pacific Islander | 2.2 | 1.9 | 0.84 | 0.88 | 0.88 | 0.99 | * | 3.9 | * | * | 1.27 | * | * | 3.9 | * | * | 1.27 | * |
| Asian | 1.8 | 1.9 | 1.05 | 0.42 | 0.46 | 1.11 | 5.2 | 5.3 | 1.03 | 0.69 | 0.67 | 0.97 | 5.8 | 6.0 | 1.03 | 0.74 | 0.77 | 1.05 |
| Two or More Races | 5.0 | 5.1 | 1.03 | 0.85 | 0.98 | 1.16 | 9.8 | 9.9 | 1.01 | 1.28 | 1.48 | 1.15 | 11.4 | 11.5 | 1.01 | 1.37 | 1.70 | 1.24 |
| Hispanic or Latino | 3.6 | 3.6 | 1.02 | 0.31 | 0.30 | 0.98 | 8.0 | 8.3 | 1.03 | 0.49 | 0.61 | 1.24 | 8.7 | 8.9 | 1.03 | 0.51 | 0.63 | 1.23 |

* Low precision; no estimate reported.

PUF = public use file, which refers to estimates produced from a NSDUH public use file (a PUF is a NSDUH restricted-use file that has undergone statistical disclosure limitation treatment); RUF = restricted-use file, which refers to estimates produced from a NSDUH RUF and published in NSDUH's detailed tables available at <http://www.samhsa.gov/data/>; SE = standard error that is associated with the estimate.

NOTE: Some 2010 RUF estimates and SEs may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the *Results from the 2011 National Survey on Drug Use and Health: National Findings*).

NOTE: $Ratio\ of\ estimates = PUF\ estimate \div RUF\ estimate$; $Ratio\ of\ SEs = PUF\ SE \div RUF\ SE$.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010.