150313

Table MH10.9 Substance Dependence or Abuse in the Past Year among Adults Aged 18 or Older, by Past Year Major Depressive Episode (MDE): Comparison of Percentages and Standard Errors from Restricted-Use File and Public Use File Estimates, 2010 NSDUH

| | TOTAL ¹ | | | | | | MDE | | | | | | NO MDE | | | | | |
|--|--------------------|-----------------|-----------------------|--------|--------|-----------------|-----------------|-----------------|-----------------------|--------|--------|-----------------|-----------------|-----------------|-----------------------|--------|--------|-----------------|
| Dependence or Abuse | RUF Estimate | PUF Estimate | Ratio of Estimates | RUF SE | PUF SE | Ratio of SEs | RUF Estimate | PUF Estimate | Ratio of Estimates | RUF SE | PUF SE | Ratio of SEs | RUF Estimate | PUF Estimate | Ratio of Estimates | RUF SE | PUF SE | Ratio of SEs |
| DEPENDENCE OR ABUSE | | | | | | | | | | | | | | | | | | |
| Illicit Drugs ² | 2.6 | 2.6 | 0.98 | 0.10 | 0.13 | 1.28 | 8.8 | 8.6 | 0.97 | 0.78 | 0.89 | 1.15 | 2.1 | 2.1 | 0.99 | 0.09 | 0.10 | 1.14 |
| Marijuana | 1.6 | 1.5 | 0.97 | 0.08 | 0.09 | 1.21 | 4.9 | 4.9 | 0.98 | 0.61 | 0.64 | 1.06 | 1.3 | 1.3 | 0.97 | 0.07 | 0.08 | 1.13 |
| Illicit Drugs Other Than Marijuana ² | 1.3 | 1.3 | 1.00 | 0.08 | 0.09 | 1.22 | 5.8 | 5.6 | 0.98 | 0.69 | 0.74 | 1.07 | 1.0 | 1.0 | 1.00 | 0.06 | 0.07 | 1.08 |
| Alcohol | 7.4 | 7.4 | 1.00 | 0.18 | 0.21 | 1.17 | 17.1 | 16.7 | 0.98 | 0.91 | 1.00 | 1.10 | 6.6 | 6.7 | 1.02 | 0.18 | 0.22 | 1.22 |
| Both Illicit Drugs and Alcohol ² | 1.1 | 1.0 | 0.95 | 0.05 | 0.07 | 1.19 | 3.9 | 3.7 | 0.93 | 0.44 | 0.45 | 1.01 | 0.8 | 0.8 | 0.97 | 0.05 | 0.06 | 1.15 |
| Illicit Drugs or Alcohol ² | 8.9 | 9.0 | 1.01 | 0.20 | 0.23 | 1.17 | 22.0 | 21.6 | 0.98 | 1.08 | 1.22 | 1.13 | 7.9 | 8.0 | 1.01 | 0.20 | 0.24 | 1.20 |
| DEPENDENCE | | | | | | | | | | | | | | | | | | |
| Illicit Drugs ² | 1.8 | 1.8 | 0.99 | 0.09 | 0.11 | 1.22 | 7.0 | 6.9 | 0.98 | 0.72 | 0.80 | 1.11 | 1.5 | 1.4 | 0.99 | 0.08 | 0.09 | 1.12 |
| Marijuana | 1.0 | 1.0 | 0.99 | 0.07 | 0.08 | 1.20 | 3.5 | 3.5 | 1.01 | 0.54 | 0.59 | 1.09 | 0.8 | 0.8 | 0.99 | 0.06 | 0.06 | 1.14 |
| Illicit Drugs Other Than Marijuana ² | 1.0 | 1.0 | 0.99 | 0.07 | 0.08 | 1.16 | 4.8 | 4.6 | 0.97 | 0.65 | 0.67 | 1.03 | 0.7 | 0.7 | 1.00 | 0.05 | 0.06 | 1.07 |
| Alcohol | 3.5 | 3.5 | 1.00 | 0.13 | 0.14 | 1.08 | 11.2 | 11.1 | 0.99 | 0.82 | 0.87 | 1.07 | 2.9 | 2.9 | 1.01 | 0.13 | 0.14 | 1.13 |
| Both Illicit Drugs and Alcohol ² | 0.5 | 0.4 | 0.96 | 0.04 | 0.04 | 1.10 | 1.9 | 1.9 | 0.96 | 0.28 | 0.27 | 0.97 | 0.3 | 0.3 | 0.96 | 0.03 | 0.04 | 1.12 |
| Illicit Drugs or Alcohol ² | 4.9 | 4.9 | 1.00 | 0.16 | 0.18 | 1.14 | 16.3 | 16.1 | 0.99 | 1.02 | 1.22 | 1.19 | 4.0 | 4.0 | 1.00 | 0.15 | 0.17 | 1.13 |

^{*} Low precision; no estimate reported.

PUF = public use file, which refers to estimates produced from a NSDUH public use file (a PUF is a NSDUH restricted-use file that has undergone statistical disclosure limitation treatment); RUF = restricted-use file, which refers to estimates produced from a NSDUH RUF and published in NSDUH's mental health detailed tables available at http://www.samhsa.gov/data/; SE = standard error that is associated with the estimate.

NOTE: Some 2010 RUF estimates and SEs may differ from previously published estimates due to updates (see Section B.3 in Appendix B of the Results from the 2011 National Survey on Drug Use and Health: Mental Health Findings).

NOTE: Ratio of estimates = PUF estimate \div RUF estimate; Ratio of SEs = PUF SE \div RUF SE.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV).

NOTE: Major Depressive Episode (MDE) is defined as in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV), which specifies a period of at least 2 weeks when a person experienced a depressed mood or loss of interest or pleasure in daily activities and had a majority of specified depression symptoms. Respondents with unknown past year MDE data were excluded.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2010.

¹ Estimates in the Total column represent adults aged 18 or older, including those with unknown past year MDE information.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.