

Table MH12.4 Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Adults Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Comparison of Percentages and Standard Errors from Restricted-Use File and Public Use File Estimates, 2012 NSDUH

Gender/ Age Group	TOTAL						ANY MENTAL ILLNESS						SERIOUS MENTAL ILLNESS					
	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs
TOTAL	8.8	8.8	1.00	0.20	0.20	1.01	19.2	19.4	1.01	0.65	0.69	1.07	27.3	27.4	1.00	1.54	1.56	1.01
18-25	18.9	19.0	1.00	0.39	0.38	0.99	34.5	34.8	1.01	1.00	0.91	0.91	39.9	39.6	0.99	2.10	2.17	1.03
26-49	10.5	10.4	0.99	0.33	0.35	1.05	22.6	22.6	1.00	0.98	1.01	1.03	29.4	29.4	1.00	2.08	2.33	1.12
50 or Older	3.8	3.8	1.01	0.26	0.30	1.15	8.6	9.0	1.05	0.95	1.27	1.34	18.0	18.4	1.03	3.09	3.92	1.27
MALE	12.2	12.0	0.99	0.34	0.31	0.92	27.7	27.8	1.00	1.24	1.19	0.96	38.1	37.8	0.99	2.98	3.33	1.12
18-25	22.7	22.8	1.01	0.61	0.59	0.96	42.9	43.3	1.01	1.73	1.60	0.93	44.8	45.3	1.01	3.81	3.52	0.93
26-49	14.3	14.0	0.98	0.54	0.53	0.98	32.4	32.3	1.00	1.73	1.90	1.10	39.8	39.8	1.00	3.82	4.39	1.15
50 or Older	6.2	6.1	0.99	0.51	0.53	1.05	14.7	15.1	1.03	2.03	2.31	1.14	*	31.5	*	*	8.21	*
FEMALE	5.7	5.8	1.02	0.20	0.21	1.06	13.9	14.2	1.02	0.65	0.70	1.08	20.7	21.1	1.02	1.57	1.53	0.98
18-25	15.1	15.1	1.00	0.45	0.50	1.11	29.2	29.4	1.01	1.24	1.30	1.05	37.0	36.4	0.98	2.60	2.63	1.01
26-49	6.8	6.9	1.02	0.36	0.39	1.09	16.1	16.3	1.01	1.02	1.05	1.02	22.7	23.3	1.03	2.19	2.42	1.11
50 or Older	1.7	1.8	1.07	0.23	0.24	1.05	5.0	5.3	1.06	0.90	0.99	1.10	9.9	10.0	1.01	2.86	2.75	0.96

See notes at end of table.

(continued)

Table MH12.4 Illicit Drug or Alcohol Dependence or Abuse in the Past Year among Adults Aged 18 or Older, by Past Year Level of Mental Illness, Gender, and Age Group: Comparison of Percentages and Standard Errors from Restricted-Use File and Public Use File Estimates, 2012 NSDUH (continued)

Gender/ Age Group	MODERATE MENTAL ILLNESS						MILD MENTAL ILLNESS						NO MENTAL ILLNESS					
	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs
TOTAL	18.9	18.9	1.00	1.13	0.96	0.86	15.9	16.3	1.02	0.82	1.02	1.24	6.4	6.3	0.99	0.18	0.20	1.11
18-25	36.0	38.3	1.06	1.93	1.88	0.98	31.6	31.1	0.99	1.41	1.37	0.97	15.1	15.1	1.00	0.39	0.39	1.02
26-49	23.3	22.9	0.98	1.82	1.68	0.92	18.7	19.1	1.02	1.32	1.65	1.25	7.2	7.0	0.97	0.30	0.32	1.07
50 or Older	6.5	6.4	0.99	1.37	1.52	1.11	6.3	7.0	1.11	1.07	1.46	1.36	2.9	2.9	0.99	0.27	0.30	1.12
MALE	26.7	26.1	0.97	2.27	2.27	1.00	23.9	24.6	1.03	1.53	1.72	1.13	9.4	9.2	0.97	0.33	0.31	0.94
18-25	48.4	48.9	1.01	3.26	3.47	1.06	39.7	40.2	1.01	2.32	2.36	1.02	19.0	19.1	1.00	0.60	0.55	0.92
26-49	32.4	31.4	0.97	3.42	3.81	1.11	28.7	29.4	1.02	2.39	2.90	1.21	10.5	10.2	0.97	0.51	0.51	0.99
50 or Older	11.4	10.9	0.96	3.05	3.43	1.12	10.2	11.2	1.10	2.16	2.33	1.08	5.0	4.8	0.97	0.52	0.54	1.04
FEMALE	13.9	14.3	1.03	1.17	1.02	0.87	11.0	11.1	1.01	0.84	1.11	1.32	3.3	3.4	1.01	0.16	0.17	1.03
18-25	28.9	32.4	1.12	2.37	2.80	1.18	25.9	24.6	0.95	1.75	1.85	1.06	10.7	10.5	0.99	0.43	0.46	1.07
26-49	17.3	17.0	0.98	1.96	1.47	0.75	12.1	12.4	1.02	1.34	1.68	1.25	3.7	3.7	1.02	0.30	0.28	0.96
50 or Older	3.3	3.6	1.09	1.06	1.14	1.07	4.1	4.5	1.11	1.17	1.50	1.28	0.9	1.0	1.07	0.19	0.22	1.16

* Low precision; no estimate reported.

PUF = public use file, which refers to estimates produced from a NSDUH public use file (a PUF is a NSDUH restricted-use file that has undergone statistical disclosure limitation treatment); RUF = restricted-use file, which refers to estimates produced from a NSDUH RUF and published in NSDUH's mental health detailed tables available at <http://www.samhsa.gov/data/>; SE = standard error that is associated with the estimate.

NOTE: $Ratio\ of\ estimates = PUF\ estimate \div RUF\ estimate$; $Ratio\ of\ SEs = PUF\ SE \div RUF\ SE$.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder that met the criteria found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes individuals in any of the three categories. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2012 National Survey on Drug Use and Health: Mental Health Findings*.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2012.