

**Table MH13.3 Substance Dependence or Abuse in the Past Year among Adults Aged 18 or Older, by Past Year Level of Mental Illness: Comparison of Percentages and Standard Errors from Restricted-Use File and Public Use File Estimates, 2013 NSDUH**

Dependence or Abuse Status	TOTAL						ANY MENTAL ILLNESS						SERIOUS MENTAL ILLNESS					
	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs
<b>TOTAL POPULATION</b>	100.0	100.0	1.00	0.00	0.00	.	100.0	100.0	1.00	0.00	0.00	.	100.0	100.0	1.00	0.00	0.00	.
<b>DEPENDENCE</b>																		
Both Illicit Drugs and Alcohol	0.4	0.4	1.04	0.04	0.04	1.17	1.5	1.5	0.99	0.16	0.18	1.13	2.4	2.4	0.98	0.40	0.48	1.20
Illicit Drugs or Alcohol	4.7	4.7	1.00	0.14	0.15	1.10	12.6	12.6	1.00	0.53	0.68	1.29	17.8	17.9	1.01	1.22	1.35	1.10
Illicit Drugs	1.9	1.9	1.02	0.09	0.10	1.20	5.8	5.7	0.98	0.34	0.37	1.09	8.7	8.6	0.98	0.74	0.88	1.18
Alcohol	3.3	3.2	0.99	0.12	0.13	1.04	8.3	8.5	1.02	0.44	0.59	1.34	11.4	11.7	1.02	1.08	1.44	1.33
<b>DEPENDENCE OR ABUSE</b>																		
Both Illicit Drugs and Alcohol	1.0	1.0	1.01	0.06	0.07	1.15	3.1	3.1	1.01	0.25	0.27	1.06	4.8	4.5	0.93	0.59	0.60	1.02
Illicit Drugs or Alcohol	8.5	8.6	1.00	0.19	0.21	1.09	17.5	18.0	1.03	0.61	0.78	1.28	23.1	24.0	1.04	1.40	1.43	1.03
Illicit Drugs	2.5	2.5	1.01	0.10	0.12	1.23	7.1	7.1	1.00	0.37	0.43	1.17	10.5	10.5	1.00	0.85	0.96	1.12
Alcohol	7.0	7.0	1.00	0.17	0.19	1.09	13.4	14.0	1.04	0.55	0.68	1.23	17.4	18.0	1.03	1.29	1.37	1.06

See notes at end of table.

(continued)

**Table MH13.3 Substance Dependence or Abuse in the Past Year among Adults Aged 18 or Older, by Past Year Level of Mental Illness: Comparison of Percentages and Standard Errors from Restricted-Use File and Public Use File Estimates, 2013 NSDUH (continued)**

Dependence or Abuse Status	MODERATE MENTAL ILLNESS						MILD MENTAL ILLNESS						NO MENTAL ILLNESS <sup>1</sup>					
	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs	RUF Estimate	PUF Estimate	Ratio of Estimates	RUF SE	PUF SE	Ratio of SEs
<b>TOTAL POPULATION</b>	100.0	100.0	1.00	0.00	0.00	.	100.0	100.0	1.00	0.00	0.00	.	100.0	100.0	1.00	0.00	0.00	.
<b>DEPENDENCE</b>																		
Both Illicit Drugs and Alcohol	2.2	2.1	0.93	0.42	0.41	0.99	0.8	0.8	1.08	0.16	0.18	1.15	0.2	0.2	1.12	0.03	0.04	1.40
Illicit Drugs or Alcohol	14.8	14.5	0.98	1.13	1.19	1.06	9.2	9.3	1.01	0.61	0.88	1.45	2.9	2.9	0.98	0.13	0.12	0.91
Illicit Drugs	7.1	7.1	1.00	0.84	0.81	0.97	3.8	3.6	0.96	0.38	0.36	0.95	1.0	1.0	1.06	0.08	0.08	1.06
Alcohol	9.9	9.5	0.96	0.88	0.89	1.01	6.1	6.5	1.06	0.52	0.76	1.47	2.1	2.0	0.96	0.11	0.10	0.90
<b>DEPENDENCE OR ABUSE</b>																		
Both Illicit Drugs and Alcohol	4.0	4.2	1.03	0.70	0.81	1.16	1.8	2.0	1.10	0.22	0.27	1.21	0.5	0.5	1.00	0.04	0.05	1.20
Illicit Drugs or Alcohol	19.6	20.1	1.02	1.29	1.45	1.12	13.9	14.2	1.03	0.75	1.07	1.42	6.5	6.4	0.99	0.19	0.18	0.93
Illicit Drugs	8.6	8.8	1.02	0.89	0.93	1.04	4.8	4.8	0.98	0.42	0.45	1.08	1.5	1.5	1.01	0.09	0.09	1.06
Alcohol	15.0	15.4	1.03	1.17	1.37	1.18	10.8	11.4	1.06	0.68	0.98	1.44	5.5	5.4	0.98	0.17	0.17	0.99

\* Low precision; no estimate reported.

PUF = public use file, which refers to estimates produced from a NSDUH public use file (a PUF is a NSDUH restricted-use file that has undergone statistical disclosure limitation treatment); RUF = restricted-use file, which refers to estimates produced from a NSDUH RUF and published in NSDUH's mental health detailed tables available at <http://www.samhsa.gov/data/>; SE = standard error that is associated with the estimate.

NOTE:  $Ratio\ of\ estimates = PUF\ estimate \div RUF\ estimate$ ;  $Ratio\ of\ SEs = PUF\ SE \div RUF\ SE$ .

NOTE: Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, including data from original methamphetamine questions but not including new methamphetamine items added in 2005 and 2006.

NOTE: Mental Illness is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder, assessed by the Mental Health Surveillance Study (MHSS) *Structured Clinical Interview for the Diagnostic and Statistical Manual of Mental Disorders—Fourth Edition—Research Version—Axis I Disorders* (MHSS-SCID), which is based on the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV). Three categories of mental illness severity are defined based on the level of functional impairment: mild mental illness, moderate mental illness, and serious mental illness. Any mental illness includes individuals in any of the three categories. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status. For details on the methodology, see Section B.4.3 in Appendix B of the *Results from the 2013 National Survey on Drug Use and Health: Mental Health Findings*.

<sup>1</sup> Respondents with no mental illness may still have received mental health treatment for a past disorder that is in partial or full remission or treatment for a disorder that was not included in the definition of mental illness or for other reasons.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2013.