

TAPS Tool Part I

Directions: The TAPS Tool Part 1 is a 4-item screening for tobacco use, alcohol use, prescription medication misuse, and illicit substance use in the PAST YEAR. Question 2 should be answered by males, and Question 3 should be answered by females. Each of the four multiple-choice items has five possible responses to choose from. Check the box to select your answer.

In the PAST 12 MONTHS:

1.	How often have you used any tobacco product (for example, cigarettes, ecigarettes, cigars, pipes, or smokeless tobacco)?				
	□ Never	\square Less than monthly	\square Monthly	☐ Weekly	☐ Daily or almost daily
2.	How often have you had 5 or more drinks containing alcohol in 1 day? One standard drink is about 1 small glass of wine (5 oz), 1 beer (12 oz), or 1 single shot of liquor. (Note: This question should only be answered by males.)				
	□ Never	\square Less than monthly	\square Monthly	☐ Weekly	\square Daily or almost daily
3.	How often have you had 4 or more drinks containing alcohol in 1 day? One standard drink is about 1 small glass of wine (5 oz), 1 beer (12 oz), or 1 single shot of liquor. (Note: This question should only be answered by females.)				
	□ Never	\square Less than monthly	\square Monthly	☐ Weekly	\square Daily or almost daily
4.	How often have you used any drugs including marijuana, cocaine or crack, heroin, methamphetamine (crystal meth), hallucinogens, or ecstasy/MDMA?				
	□ Never	\square Less than monthly	\square Monthly	☐ Weekly	\square Daily or almost daily
5.	How often have you used any prescription medications just for the feeling, more than prescribed, or that were not prescribed for you? Prescription medications that may be used this way include opiate pain relievers (for example, OxyContin, Vicodin, Percocet, or methadone), medications for anxiety or sleeping (for example, Xanax, Ativan, or Klonopin), or medications for ADHD (for example, Adderall or Ritalin).				
	□ Never	\square Less than monthly	\square Monthly	☐ Weekly	\square Daily or almost daily