

TAPS Tool Part 2

Directions: The TAPS Tool Part 2 is a brief assessment for tobacco use, alcohol use, illicit substance use, and prescription medication misuse in the PAST 3 MONTHS ONLY. Each of the following questions and subquestions has two possible answers, yes or no. Check the box to select your answer.

In the PAST 3 MONTHS:

1.	Did you smoke a cigarette containing tobacco?	☐ Yes	□ No
	If "Yes," answer the following questions:		
	Did you usually smoke more than 10 cigarettes each day?Did you usually smoke within 30 minutes after waking?	☐ Yes ☐ Yes	□ No □ No
2.	Did you have a drink containing alcohol?	☐ Yes	□ No
	If "Yes," answer the following questions:		
	 Did you have 4 or more drinks containing alcohol in a day?* (Note: This question should only be answered by females.) 	☐ Yes	□ No
	 Did you have 5 or more drinks containing alcohol in a day?* (Note: This question should only be answered by males.) 	☐ Yes	□ No
	Have you tried and failed to control, cut down, or stop drinking?Has anyone expressed concern about your drinking?	☐ Yes ☐ Yes	□ No □ No
3.	Did you use marijuana (hash, weed)?	☐ Yes	□ No
	If "Yes," answer the following questions:		
	 Have you had a strong desire or urge to use marijuana at least once a week or more often? Has anyone expressed concern about your use of marijuana? 	☐ Yes ☐ Yes	□ No □ No
4.	Did you use cocaine, crack, or methamphetamine (crystal meth)?	☐ Yes	□ No
	If "Yes," answer the following questions:		
	 Did you use cocaine, crack, or methamphetamine (crystal meth) at least once a week or more often? Has anyone expressed concern about your use of cocaine, crack, or methamphetamine (crystal 	☐ Yes	□ No
	meth)?	☐ Yes	☐ No
5.	Did you use heroin?	☐ Yes	□ No
	If "Yes," answer the following questions:		
	Have you tried and failed to control, cut down, or stop using heroin?Has anyone expressed concern about your use of heroin?	☐ Yes ☐ Yes	□ No □ No
6.	Did you use a prescription opiate pain reliever (for example, Percocet or Vicodin) not as prescribed or that was not prescribed for you?	☐ Yes	□ No
	If "Yes," answer the following questions:		
	Have you tried and failed to control, cut down, or stop using an opiate pain reliever?Has anyone expressed concern about your use of an opiate pain reliever?	☐ Yes ☐ Yes	□ No □ No

Continued on next page

^{*}One standard drink is about 1 small glass of wine (5 oz), 1 beer (12 oz), or 1 single shot of liquor.



TAPS Tool Part 2 (continued)

7.	Did you use medication for anxiety or sleep (for example, Xanax, Ativan, or Klonopin) not as prescribed or that was not prescribed for you?	☐ Yes	□ No
	If "Yes," answer the following questions:		
	 Have you had a strong desire or urge to use medications for anxiety or sleep at least once a week or more often? 	☐ Yes	□ No
	 Has anyone expressed concern about your use of medication for anxiety or sleep? 	☐ Yes	□ No
8.	Did you use medication for ADHD (for example, Adderall or Ritalin) not as prescribed or that was not prescribed for you?	☐ Yes	□ No
	If "Yes," answer the following questions:		
	• Did you use a medication for ADHD (for example, Adderall or Ritalin) at least once a week or more often?	☐ Yes	□ No
	 Has anyone expressed concern about your use of medication for ADHD (for example, Adderall or Ritalin)? 	☐ Yes	□ No
9.	Did you use any other illegal or recreational drugs (for example, ecstasy, molly, GHB, poppers, LSD, mushrooms, special K, bath salts, synthetic marijuana ["spice"], or whip-its)? If "Yes," answer the following question:	☐ Yes	□ No
	What were the other drug(s) you used? (write in response)		

The complete tool is available online (https://cde.drugabuse.gov/instrument/29b23e2e-e266-f095-e050-bb89ad43472f). Adapted from material in the public domain. 103