

Walking distance and HRQoL outcome measures used in included studies

| Trial name | Treatment and dose | Outcome measures for PFWD and MWD | Outcome measures for HRQoL |
|---------------------------------------|---|---|----------------------------|
| CASTLE, Hiatt 2008 ⁴⁸⁻⁵⁰ | Cilostazol 200 mg | NR | NR |
| O'Donnell 2009 ^{51,53-55,83} | Cilostazol 200 mg | <i>Treadmill with constant workload:</i> 3.2 km/hour (2 mph) 10% gradient | SF-36 VascuQoL |
| Strandness 2002 ^{56,57} | Cilostazol 200 mg | <i>Treadmill with constant workload:</i> 3.2 km/hour (2 mph) 12.5% gradient | SF-36 WIQ COM |
| Dawson 2000 ⁵⁸⁻⁶⁰ | Cilostazol 200 mg, pentoxifylline 1200 mg | <i>Treadmill with graded test:</i> 3.2 km/hour (2 mph) 0% gradient with a 3.5% increase in gradient every 3 minutes | SF-36 WIQ |
| Beebe 1999 ⁶¹ | Cilostazol 200 mg | <i>Treadmill with constant workload:</i> 3.2 km/hour (2 mph) 12.5% gradient | SF-36 WIQ COM |
| Otsuka 21-94-301 ³⁴ | Cilostazol 200 mg, pentoxifylline 1200 mg | <i>Treadmill with constant workload:</i> 3.2 km/hour (2 mph) 12.5% gradient | NR |
| Otsuka 21-98-213 ³⁴ | Cilostazol 200 mg, pentoxifylline 1200 mg | <i>Treadmill with constant workload:</i> 3.2 km/hour (2 mph) 12.5% gradient | SF-36 WIQ COM |
| Money 1998 ⁶² | Cilostazol 200 mg | <i>Treadmill with graded test:</i> 3.2 km/hour (2 mph) 0% gradient with a 3.5% increase in gradient every 3 minutes | SF-36 WIQ |
| Dawson 1998 ⁶³ | Cilostazol 200 mg | <i>Treadmill with constant workload:</i> 3.2 km/hour (2 mph) 12.5% gradient | NR |
| Elam 1998 ⁶⁴ | Cilostazol 200 mg | <i>Treadmill with graded test:</i> 3.2 km/hour (2 mph) 0% gradient with a 3.5% increase in gradient every 3 minutes | NR |
| Otsuka 21-95-201 ³⁴ | Cilostazol 200 mg | <i>Treadmill with constant workload:</i> 3.2 km/hour (2 mph) 12.5% gradient | SF-36 WIQ |
| INEXACT, Hobbs 2007 ⁸² | Cilostazol 200 mg, cilostazol 200 mg plus supervised exercise | <i>Treadmill with constant workload:</i> 3 km/hour 10% gradient | NR |
| Spengel 2002 ⁴⁷ | Naftidrofuryl oxalate 600 mg | Estimated by patient | CLAU-S |
| Kieffer 2001 ⁶⁵ | Naftidrofuryl oxalate 600 mg | <i>Treadmill with constant workload:</i> 3.2 km/hour (2 mph) 10% gradient | NR |
| Adhoute 1986 ⁶⁶ | Naftidrofuryl oxalate 600 mg | <i>Treadmill with constant workload:</i> 3.2 km/hour (2 mph) 10% gradient | NR |

| Trial name | Treatment and dose | Outcome measures for PFWD and MWD | Outcome measures for HRQoL |
|---|------------------------------|---|----------------------------|
| Trubestein 1984 ⁶⁷ | Naftidrofuryl oxalate 600 mg | <i>Treadmill with constant workload:</i> 5 km/hour 10% gradient, performed twice with at least 20 minutes interval | NR |
| Ruckley 1978 ⁶⁸ | Naftidrofuryl oxalate 300 mg | Unclear if treadmill used < 100 yards = severe 100–200 yards = moderate > 200 yards = mild | NR |
| Dettori 1989 ⁶⁹ | Pentoxifylline 1200 mg | <i>Treadmill with varied workload:</i> 3 km/hour. If PFW > 30 minutes, higher speed was used in the second test (5 km/hour) 10% gradient | NR |
| Creager 2008 ⁷⁰ | Pentoxifylline 1200 mg | <i>Treadmill with graded test:</i> 3.2 km/hour (2 mph) 0% gradient, increased by 2% every 2 minutes | SF-36 WIQ |
| Lindgarde 1989 ⁷¹ | Pentoxifylline 1200 mg | <i>Treadmill with constant workload:</i> 3.2 km/hour (2 mph) 12.5% gradient | NR |
| Porter 1982 ^{72,74} and Gillings 1987 ^{73,75} | Pentoxifylline 1200 mg | <i>Treadmill with constant workload:</i> 1.5 mph 7° gradient, two treadmill tests were performed at 30- to 60-minute intervals and the mean of the two tests used | NR |
| Gallus 1985 ⁷⁶ | Pentoxifylline 1200 mg | <i>Treadmill with constant workload:</i> 4 km/hour 10° gradient | NR |
| Di Perri 1983 ⁷⁷ | Pentoxifylline 1200 mg | Absolute distance covered by walking on horizontal level at metronome controlled speed of 120 steps/minute. Walking test was performed three times and a mean taken | NR |
| O'Hara 1988 ^{78,79} | Inositol nicotinate 4 g | Training device (pair of stirrups in a metal frame), which simulated box-stepping. Elapsed time and number of steps to claudication and time to recovery were recorded. Waist-band pedometer to record 'similar weekly walks' | NR |
| Kiff 1988 ⁸⁰ | Inositol nicotinate 4 g | <i>Treadmill with constant workload:</i> 10% gradient | NR |
| Head 1986 ⁸¹ | Inositol nicotinate 4 g | Time to claudication. Patients climbed up and down the first two steps of a standard ladder in time with a metronome set at 80 beats/minute leading with the worse leg | NR |

mph, miles per hour; NR, not reported.